

debted to Professor Yoshinori Kawano of Tohoku University for his kind advice and constant help with this work.

## REFERENCES

- AKIMOTO, S. AND T. KATSURA (1959) Magnetochemical study of the generalized titanomagnetite in volcanic rocks. *Geomag. Geoelectr.* **10**, 69–90.
- AOKI, K. (1959) Petrology of alkali rocks of the Iki Islands and Higashi-matsuura District, Japan. *Sci. Rep. Tohoku Univ. Ser. III*, **6**, 261–310.
- (1963a) The kaersutites and oxykaersutites from alkalic rocks of Japan and surrounding areas. *Jour. Petrology* **4**, 198–210.
- (1963b) On the anorthoclase feldspars of the alkalic rocks from the Iki and adjacent Islands. *Jour. Japan. Assoc. Mineral. Petrol. Econ. Geol.* **50**, 77–80 (in Japanese, Engl. abs.).
- (1964) Clinopyroxenes from alkaline rocks of Japan. *Am. Mineral.* **49**, 1199–1223.
- BASTA, E. Z. (1960) Natural and synthetic titanomagnetites (the system  $\text{Fe}_3\text{O}_4$ – $\text{Fe}_2\text{TiO}_4$ – $\text{FeTiO}_3$ ). *Neues Jb. Mineral., Abh., Bd.* **94**, 1017–1048.
- DEER, W. A., R. A. HOWIE AND J. ZUSSMAN (1962) *Rockforming Minerals*. Vol. 5, Non-silicates. Longmans Co. Ltd., London.
- KATSURA, T. AND I. KUSHIRO (1961) Titanomaghemite in igneous rocks. *Am. Mineral.* **46**, 134–145.
- KAWAI, N., S. KUME AND S. SASAJIMA (1954) Magnetism of rocks and solid phase transformation in ferromagnetic minerals. *Proc. Japan. Acad.* **30**, 588.
- MATSUI, K. (1958) Explanatory text of the geological map of Japan, Katsumoto, Gononra and Ashibe. *Geol. Surv. Japan* (in Japanese, Engl. abst.).
- VINCENT, E. A., J. B. WRIGHT, R. CHEVALLIER AND S. MATHIEU (1957) Heating experiments on some natural titaniferous magnetites. *Mineral. Mag.* **31**, 624–655.
- WRIGHT, J. B. (1959) Some further heating experiments on natural titaniferous magnetites. *Mineral. Mag.* **32**, 32–37.

THE AMERICAN MINERALOGIST, VOL. 51, NOVEMBER–DECEMBER, 1966

## CELL DIMENSIONS AND SPACE GROUP OF TAMARUGITE

PAUL D. ROBINSON, J. H. FANG AND F. D. BLOSS *Southern Illinois University, Carbondale, Illinois*

Tamarugite,  $\text{NaAl}(\text{SO}_4)_2 \cdot 6\text{H}_2\text{O}$ , is a secondary mineral formed from the oxidation of sulfides, usually under arid conditions. Optical examination of crystals found with sideronatrite in a sample from Mina de la Compania, Sierra Gorda, Chile (U. S. National Museum, Smithsonian Cat. No. R6287) revealed  $\alpha = 1.485$ ,  $\beta = 1.487$  and  $\gamma = 1.498$ , all  $\pm 0.002$ . These thus agree with the values,  $\alpha = 1.484$ ,  $\beta = 1.486$ ,  $\gamma = 1.497$ , all  $\pm 0.001$ —reported by Gordon (1940). Similarly, the morphological and physical properties observed in the tamarugite of this present study confirm those reported by Gordon (1940). Published x-ray data on tamarugite were not found by the writers. Thus it was necessary to determine

TABLE 1. INDEXED POWDER DATA FOR TAMARUGITE

| d(obs) | I/I' | d(calc) | hkl          | d(obs) | I/I' | d(calc) | hkl             |
|--------|------|---------|--------------|--------|------|---------|-----------------|
| 12.538 | 16   | 12.612  | 020          | 2.544  | 8    | 2.545   | 091             |
| 7.034  | 6    | 7.033   | 110          | 2.521  | 4    | 2.522   | 0·10·0          |
| 6.339  | 2    | 6.333   | 120          | 2.509  | 2    | 2.513   | 15 $\bar{2}$    |
| 5.530  | 8    | 5.523   | 130          | 2.462  | 9    | 2.462   | 062             |
| 4.923  | 6    | 4.923   | 031          | 2.350  | 2    | 2.350   | 23 $\bar{2}$    |
| 4.557  | 4    | 4.563   | 12 $\bar{1}$ | 2.333  | 4    | 2.330   | 0·10·1          |
| 4.424  | 3    | 4.413   | 111          | 2.329  | 6    | 2.328   | 31 $\bar{1}$    |
| 4.370  | 4    | 4.374   | 041          | 2.300  | 6    | 2.299   | 32 $\bar{1}$    |
| 4.223  | 100  | 4.223   | 121          | 2.281  | 3    | 2.281   | 24 $\bar{2}$    |
| 4.207  | 80   | 4.204   | 060          | 2.269  | 4    | 2.269   | 28 $\bar{1}$    |
| 4.157  | 12   | 4.155   | 150          | 2.228  | 4    | 2.226   | 290             |
| 3.964  | 32   | 3.955   | 131          | 2.189  | 7    | { 2.188 | 1·11·0          |
| 3.861  | 2    | 3.866   | 14 $\bar{1}$ |        |      | { 2.190 | 311             |
| 3.647  | 59   | 3.646   | 160          | 2.116  | 3    | 2.115   | 262             |
| 3.355  | 3    | 3.351   | 151          | 2.102  | 6    | 2.102   | 0·12·0          |
| 3.266  | 3    | 3.267   | 20 $\bar{1}$ |        |      | { 2.060 | 182             |
| 3.170  | 10   | 3.167   | 240          | 2.060  | 2    | { 2.059 | 092             |
| 3.162  | 12   | 3.163   | 22 $\bar{1}$ | 2.045  | 3    | 2.044   | 36 $\bar{1}$    |
| 3.153  | 21   | 3.153   | 080          |        |      | { 2.024 | 003             |
| 3.036  | 5    | 3.036   | 002          | 2.025  | 2    | { 2.024 | 272             |
| 2.949  | 2    | 2.952   | 022          |        |      | { 1.968 | 033             |
|        |      | { 2.896 | 180          | 1.969  | 1    | { 1.967 | 322             |
| 2.899  | 23   | { 2.901 | 24 $\bar{1}$ |        |      | { 1.904 | 143             |
|        |      | { 2.902 | 17 $\bar{1}$ | 1.904  | 3    | { 1.903 | 1·10· $\bar{2}$ |
| 2.876  | 5    | 2.879   | 11 $\bar{2}$ |        |      | { 1.902 | 113             |
| 2.823  | 2    | 2.824   | 12 $\bar{2}$ |        |      | { 1.876 | 1·13·0          |
| 2.766  | 5    | 2.761   | 260          | 1.876  | 2    | { 1.877 | 371             |
| 2.739  | 3    | 2.740   | 13 $\bar{2}$ |        |      | { 1.877 | 2·11· $\bar{1}$ |
| 2.733  | 4    | 2.736   | 042          |        |      | { 1.831 | 400             |
| 2.711  | 7    | 2.705   | 112          | 1.831  | 2    | { 1.830 | 0·11· $\bar{2}$ |
| 2.604  | 5    | 2.602   | 052          |        |      | { 1.702 | 1·12· $\bar{2}$ |
|        |      | { 2.580 | 26 $\bar{1}$ | 1.702  | 3    | { 1.703 | 083             |
| 2.582  | 5    | { 2.579 | 181          | 1.679  | 2    | { 1.679 | 460             |

the cell dimensions and space group of tamarugite as a first step in a contemplated crystal-structure analysis.

Weissenberg photographs of the h0l, h1l, h2l, and hk0 reflections for copper  $K\alpha$  radiation were made for two single crystals of tamarugite, both from specimen R6287. The indexing of these photographs showed that the only systematic extinctions were of the type  $k \neq 2n$  for (0k0). Accordingly the space group is  $P2_1$  or, if the crystal is centrosymmetrical,  $P2_1/m$ . The latter agrees with the  $2/m$  crystal class reported by Palache *et al.* (1951) on the basis of morphologic data.

Accurate lattice parameters were obtained by use of a least squares FORTRAN IV program written by Fang and Wolf following the scheme proposed by Main and Woolfson (1963) in which  $\alpha_1$ - $\alpha_2$  separations in the high angle regions of zero level Weissenberg photographs are utilized. The cell parameters were then slightly refined so as to minimize the difference between  $d(\text{calc})$  and  $d(\text{obs})$  for the powder diffraction data. The cell parameters thus obtained are:  $a = 7.35_3$ ,  $b = 25.22_5$ ,  $c = 6.09_7$  and  $\beta = 95^\circ 12'$ . Ungemach (1935) reports  $\beta$  to be  $94^\circ 50'$  on the basis of morphologic data.

The amount of tamarugite in specimen R6287 was too small for a powder pattern to be obtained other than the powder photograph earlier made by Robinson (1962). However, the U. S. National Museum kindly supplied two specimens of tamarugite—Nos. C4703 from Cerro Pintados, Chile and R9117 from Alcaparrosa, Chile—for which powder patterns were obtained in a Norelco wide-angle diffractometer using  $\text{CuK}\alpha$  radiation. The powder patterns for these two specimens appeared indistinguishable. The pattern obtained from a  $1/8^\circ$  ( $2\theta$ ) per minute scan of the Cerro Pintados material, internally calibrated with spinel, was satisfactorily indexed on the basis of the cell parameters reported above (Table 1). The observed  $d$  spacings for the Cerro Pintados pattern closely matched those measured by Robinson (1962) from a powder photograph of tamarugite collected from the Sierra Gorda specimen.

The axial ratios calculated from the lattice parameters here reported—that is, 0.2915:1:0.2417—compare with those of Ungemach (1935)—namely, 0.2919:1:0.2415.

The calculated density of tamarugite, assuming  $Z = 4$ , is  $2.066 \text{ gm/cm}^3$ . The measured density, by the flotation method, is  $2.06 \pm 0.01 \text{ gm/cm}^3$ . Gordon (1940) and Ungemach (1935) report the density of tamarugite to be  $2.07 \text{ gm/cm}^3$ .

The authors wish to thank Mr. Lloyd A. Wolf for his assistance in computer programming.

#### REFERENCES

- GORDON, S. G. (1940) The identity of lapparentite with tamarugite. *Ac. Nat. Sc. Philadelphia. Not. Nat.*, **57**, 1-9.
- MAIN, P. AND M. M. WOOLFSON (1963) Accurate lattice parameters from Weissenberg photographs. *Acta. Cryst.* **16**, 731-733.
- PALACHE, C., H. BERMAN AND C. FRONDEL (1951) *The System of Mineralogy*. Vol. 2, 7th ed. John Wiley and Sons, New York.
- ROBINSON, P. D. (1962) An  $x$ -ray study of sideronatrite. M.S. thesis, Southern Illinois Univ.
- UNGEMACH, M. H. (1935) Sur certains minéraux sulfatés du Chili. *Bull. Soc. Franc. Mineral.* **58**, 97-221.