

## ACKNOWLEDGMENTS

This work was supported by NSF grant GA 1270, grant SD-89-Research from the Advanced Research Projects Agency, and the Petroleum Research Fund of The American Chemical Society grant 12 PRF-1408-06. Grateful acknowledgment is made to the donors of that fund. We wish to thank Dr. O. J. Kleppa for his assistance in reading this paper.

## REFERENCES

- DACHILLE, F., P. Y. SIMONS, AND R. ROY (1968) Pressure-temperature studies of anatase, brookite, rutile and  $\text{TiO}_2(\text{II})$ . *Amer. Mineral.* **53**, 1929-1939.
- JAMIESON, J. C., AND B. OLINGER (1968) High-pressure polymorphism of titanium dioxide. *Science*, **161**, 893-895.
- ROBIE, R. A., P. M. BETHKE, M. S. TOULMIN, AND J. L. EDWARDS (1966) X-ray crystallographic data, densities and molar volumes of minerals. *Geol. Soc. Amer. Mem.* **97**, 42.
- NAVRÓTSKY, A., AND O. J. KLEPPA (1967) Enthalpy of the anatase-rutile transformation. *J. Amer. Ceram. Soc.*, **50**, 626.
- , J. C. JAMIESON, AND O. J. KLEPPA (1967) Enthalpy of transformation of a high-pressure polymorph of titanium dioxide to the rutile modification. *Science*, **158**, 388-389.
- RAO, C. N. (1961) Kinetics and thermodynamics of the crystal structure transformation of spectroscopically pure anatase to rutile. *Can. J. Chem.*, **39**, 498-500.
- , S. R. YOGANARASIMHAN, AND P. A. FAETH (1961) Studies on the brookite-rutile transformation. *Trans. Farad. Soc.*, **57**, 504-510.
- SHANNON, R. D., AND J. A. PASK (1965) Kinetics of the anatase-rutile transformation. *J. Amer. Ceram. Soc.*, **48**, 391-398.

THE AMERICAN MINERALOGIST, VOL. 54, SEPTEMBER-OCTOBER, 1969

PRESSURE-TEMPERATURE STUDIES OF ANATASE, BROOKITE,  
RUTILE AND  $\text{TiO}_2(\text{II})$ : A REPLY

FRANK DACHILLE, P. Y. SIMONS<sup>1</sup>, AND RUSTUM ROY, *Materials Research Laboratory and Department of Geochemistry and Mineralogy, The Pennsylvania State University, University Park, Pennsylvania 16802.*

Phase relationships among the  $\text{TiO}_2$  polymorphs have been the subject of experimental work and speculation for a long time. It is probably agreed upon by all, and certainly in this laboratory, that rutile is the only stable phase at or near one atmosphere. All approaches based on direct reversible phase equilibrium methods have so far failed to give definitive solutions to equilibrium among the various phases. Our recent study was concerned with kinetics (and mechanisms) and their possible relation to "equilibrium." (See also Simons, 1967.)

<sup>1</sup> Present address: E. I. du Pont de Nemours and Company, Parlin, N. J.

In the report in which the synthesis of TiO-II was described for the first time (Dachille and Roy, 1962) as well as in a later report (Simons and Dachille, 1968) it was clearly stated that there was insufficient evidence for any statement on equilibrium. In the present paper commented on by Jamieson and Olinger this is repeated at least three times and the point is emphasized by the explicit use of the term "reaction boundaries" in every relevant figure. There is no doubt whatsoever of the interesting  $p$ - $T$  dependence of certain reaction rates in the interconversion of the various TiO<sub>2</sub> phases. These are what are summarized in the "reaction boundaries" of the figures. Unfortunately, in discussing the results incorrect statements regarding the possible equilibrium relations among TiO<sub>2</sub> polymorphs crept in, in one or two places. Jamieson and Olinger have properly pointed out and provided a valuable comment on a possible TiO<sub>2</sub> stable equilibrium diagram. Indeed it will be seen that the latter is similar to our previous diagrams for isostructural phases: PbO<sub>2</sub> and MnF<sub>2</sub> undergoing analogous transitions (White, Dachille and Roy, 1961; Azzaria and Dachille, 1961).

Kinetic anomalies such as those observed by us (and not specifically treated by Jamieson and Olinger), their significance and possible relation to stable or metastable equilibria remain a subject for further study.

It should be mentioned that the Magnet Cover brookite used by us was processed mechanically to reduce the impurity level to 1% (Simons and Dachille, 1967; Simons, 1967).

#### REFERENCES

- AZZARIA, L., AND F. DACHILLE (1961). High-pressure polymorphism of manganous fluoride. *J. Phys. Chem.*, **65**, 889-890.
- DACHILLE, F., AND R. ROY (1962). A new high-pressure form of titanium dioxide (abstr.). *Amer. Ceram. Soc. Bull.*, **41**, 225.
- , P. Y. SIMONS, AND R. ROY (1968) Pressure-temperature studies of anatase, brookite, rutile and TiO<sub>2</sub>(II). *Amer. Mineral.* **53**, 1929-1939.
- SIMONS, P. Y. (1967) *Polymorphism of Titanium Dioxide Minerals*. PhD Thesis, Pennsylvania State University.
- , AND F. DACHILLE (1968) Pressure-temperature relationships among titanium dioxide minerals (abstr.). *Geol. Soc. Amer., Spec. Pap.* **101**, 000.
- , ——— (1967). The structure of TiO<sub>2</sub>II, a high-pressure phase of TiO<sub>2</sub>. *Acta Crystallogr.*, **23**, 334-334.
- WHITE, W. B., F. DACHILLE, AND R. ROY (1961). High-pressure high-temperature polymorphism of the oxides of lead. *J. Amer. Ceram. Soc.* **44**, 170-175.