

MOLYBDENITE POLYTYPES IN THEORY AND OCCURRENCE. II. SOME NATURALLY-OCCURRING POLYTYPES OF MOLYBDENITE

JUDITH W. FRONDEL, *Department of Geological Sciences, Harvard University, Cambridge, Massachusetts*

AND

FRANS E. WICKMAN, *Department of Geochemistry and Mineralogy, Pennsylvania State University, University Park, Pennsylvania*

108 specimens of naturally-occurring molybdenite, representing 83 localities throughout the world, were studied. About 80 percent of the specimens were the  $2H_1$  polytype, three were the  $3R$  polytype, and the remainder were mixtures of  $2H_1$  and  $3R$  in varying proportions. No new modifications of naturally-occurring molybdenite were found. Copper mineralization may play a role in the production of  $3R$  and mixtures of  $2H_1+3R$  polytypes, and a possible correlation between their occurrence and enrichment of rhenium is suggested.

INTRODUCTION

This study was made on 108 specimens of molybdenite from 83 localities throughout the world. This is, by no means, an exhaustive sampling of molybdenite, nor as detailed as some studies on specific geographical areas (e.g. Marmadov, 1965; Vorma *et al.*, 1966; Khurshudyan *et al.*, 1969), but may give a general idea of the great variety of geological occurrences of molybdenite. Some information about the geological environment was obtainable for a large percent of the localities.

THE MATERIAL<sup>1</sup>

<i>Specimen number</i>	<i>Locality</i>	<i>Polytype, and approximate ratio of mixtures</i>
------------------------	-----------------	--

A. *United States*

U.S.N.M. 94598	Copper Mt., Prince of Wales Is., Alaska	$2H_1$
----------------	---	--------

Rare molybdenite occurs in the contact zone around a granite, both in the altered granitic rock and in the skarn rocks, there associated with garnet, calcite, diopside and chalcopryrite (Wright, 1915).

U.S.N.M. 96611	Santa Niño Mine, Patagonia Mts.,	$2H_1$
U.S.N.M. 97491	Santa Cruz Co., Arizona	$2H_1$

<sup>1</sup> Specimens came from the mineralogical collections of Harvard University and the U. S. National Museum.

Molybdenite occurs here as "large bodies of fine grained massive material, as disseminated grains in quartz monzonite, and as good crystals in vein quartz with pyrite" (Galbraith, 1947). Chalcopyrite is found in massive ore (Kirkemo *et al.* 1965, pp. 14-16).

U.S.N.M. 97990      50 miles S.W. of Kingman, Hualpai Mts.,       $2H_1$   
Mojave Co., Arizona.

Many greisen-pegmatite deposits are known in this area (Wilson, 1941).

U.S.N.M. 91775      Helvetia, Pima Co., Arizona       $2H_1+3R\sim 9:1$

Around Helvetia there are many deposits containing molybdenite, which generally is associated with chalcopyrite.

Harvard 106525      Arizona       $2H_1$

The locality designation is too indefinite; therefore, nothing can be stated with certainty about this specimen.

Harvard unnumbered      Kenawyer Mine, King's River Canyon,       $2H_1$   
Fresno Co., Calif.

Molybdenite occurs with chalcopyrite in quartz veins in a granite (Horton, 1916, p. 62).

Harvard 4211      Mono Lake, California       $2H_1$

This particular specimen consists of flakes of molybdenite in quartz and has been mentioned by Horton (1916, p. 59), but nothing else is known about it.

U.S.N.M. 91868      Lake Tahoe, California       $2H_1$

Contact metamorphic occurrences of molybdenite are known from this region. This specimen is probably from the Alpine Mine, south of Lake Tahoe, where molybdenite occurs together with powellite and scheelite in tactite (King 1966).

Harvard 107537      311 Drift stope level, Climax, Colorado       $2H_1$

In the ore zone, which forms a dome-shaped cap of intensely silicified rocks over a quartz monzonite, there are three different kinds of veins; sericite-topaz veins, pyrite-topaz-quartz veins, and molybdenite-quartz veins. This third type also contains pyrite, sericite, orthoclase, and fluorite. The main Climax ore is an altered granite. The molybdenite occurs mostly in fine veinlets with molybdenite, wolframite, topaz, cassiterite, monazite, pyrite and chalcopyrite (Killefer and Linz, 1952, p. 5).

Harvard 81614 Fairplay, Park Co., Colorado  $2H_1$   
 This locality simply has been mentioned (Endlich, 1878).

Harvard unnumbered Mountain Lion Mine, Magnolia Dist.,  $2H_1$   
 Boulder Co., Colorado.

Molybdenite is found on the lower levels on the borders of a large pegmatite dike associated with massive white quartz and small amounts of chalcopyrite (Worcester, 1919).

Harvard 104570 Near St. Peters Dome, El Paso Co., Colo-  $2H_1$   
 rado.

The specimen indicates that molybdenite occurs here in pegmatite.

Harvard 98594 Middle Fork, south of Arkansas River,  $2H_1+3R\sim 1:1$   
 Chaffee Co., Colorado.

Many different kinds of molybdenite deposits are known from this area. Nothing definite can be stated about the genesis of this particular specimen.

Harvard 18212	Vinalhaven, Knox Co., Maine	$2H_1$
Harvard 18213		$2H_1$
Harvard 88638		$2H_1$

Molybdenite occurs here in quartz veins and in the rapakivi granite (Stewart, 1956, p. 36, pp. 62-63).

Harvard 98290 Sanford, York Co., Maine  $2H_1$

In this specimen molybdenite occurs together with vesuvianite.

Harvard unnumbered Tunk Pond, Cherryfield, Hancock Co.,  $2H_1$   
 Maine.

Molybdenite apparently accompanies a series of pegmatite dikes (Hussey, 1958; Horton, 1916, p. 86).

Harvard unnumbered Cooper, Washington Co., Maine  $2H_1$

Molybdenite occurs here in pegmatite dikes and disseminated in the surrounding granite (Kirkemo *et al.*, 1965, p. 61).

Harvard 90189 } Harvard 108167 }	Rockport, Massachusetts	$2H_1$ $2H_1$
-------------------------------------	-------------------------	------------------

Molybdenite is found in the granites and pegmatites of this Cape Ann locality (Warren and McKinstry, 1924).

Harvard 107870	Chesterfield, Massachusetts.	$2H_1+3R\sim 4:1$
----------------	------------------------------	-------------------

Molybdenite occurs in a granite pegmatite.

Harvard 101898	Near Sheridan, Madison Co., Montana.	$2H_1+3R\sim 1:1$
----------------	--------------------------------------	-------------------

In this specimen molybdenite occurs with sulfides in a quartz vein. No details about the occurrence at this locality are known.

Harvard 88912	Austin, Nevada	$2H_1$
---------------	----------------	--------

Small quantities of molybdenite occur with quartz, pyrite and some other sulfides (even proustite) in veins cutting a Paleozoic quartzite adjacent to a stock of quartz monzonite (Schilling, 1962).

Harvard 88909	Westmoreland, New Hampshire.	$2H_1+3R\sim 10:1$
---------------	------------------------------	--------------------

Molybdenite occurs together with muscovite and apatite at this old pegmatite locality (Jackson, 1844).

Harvard 88915	Ogden Mine, New Jersey	$2H_1$
---------------	------------------------	--------

Iron ores occur here in a gneiss penetrated by pegmatites (Bayley, 1910, p. 279). Associated with the ore are molybdenite, pyrite, chalcopyrite, garnet and apatite.

Harvard unnumbered	Hude Iron Mine, Netcong, New Jersey	$2H_1$
--------------------	-------------------------------------	--------

The study specimen is in an amphibolite.

Harvard 12741	Buckwheat Mine, Franklin Furnace, New Jersey.	$2H_1$
---------------	---	--------

The specimen is in scapolite, from limestone in the wallrock.

Harvard 105464	Franklin Furnace, New Jersey	$2H_1$
----------------	------------------------------	--------

The specimen is a cleavage piece from the shaft pillar.

Harvard unnumbered From the 1200' level of Sterling Mine,  $2H_1$   
Franklin, New Jersey.

In this specimen the molybdenite occurs with sphalerite in a dark silicate matrix.

Harvard unnumbered "N.E. New Mexico" The locality designa-  $2H_1$   
tion is indefinite.

This is possibly a "porphyry copper" specimen.

U.S.N.M. 62232 Tilly Foster Mine, Putnam Co., New York  $2H_1$

Molybdenite occurs sparingly as a contact mineral in the serpentine of a metamorphosed limestone (Newland, 1921, p. 163). Other minerals observed include magnetite, chondrodite, and clinohumite.

U.S.N.M. 83349 Demming, McDowell Co., North Carolina  $3R$

Molybdenite is found with quartz in a muscovite-biotite schist. (Horton, 1916, p. 88).

Harvard 88520 Easton, Pennsylvania  $2H_1$

Franklin limestone has been intruded by granite forming pegmatite veins in the limestone. Molybdenite occurs in the serpentine formed at the contacts.

Harvard 100826 Frankford, Philadelphia, Pennsylvania  $2H_1+3R\sim 2:1$

Molybdenite occurs in minor amounts in a granitized hornblende gneiss intrusive in high-grade metamorphosed Wissahickon Schist (J. W. Frondel, personal communication).

Harvard 88910 Chester, Pa.  $2H_1$

Molybdenite occurs in quartz veins in a hydrothermally emplaced granodiorite (Postel, 1940).

Harvard 97231 Little Cottonwood Canyon, Utah  $2H_1$

The intrusive rocks at this locality are jointed and mineralized along the joints (Butler

*et al.*, 1920, p. 265). Pyrite is the most common sulfide, but chalcopyrite, galena and molybdenite also occur.

Harvard 4341                      Cuttingsville, Vermont                       $2H_1+3R\sim 10:1$

Molybdenite occurs in syenite at its contact with gneiss on the crest of Granite Hill (Eggleston, 1918). "The molybdenite scales are of medium size, often thickly powdered, like the inclosing syenite, with yellow molybdenite."

Harvard 81612	}                      Okanogan Co., Washington	$2H_1$
Harvard 81613		$2H_1$
Harvard 81621		$2H_1$

In this area molybdenite occurs in a granite associated with pyrite (Horton, 1916, p. 83. U.S.G.S. 1966), but the genesis of the study specimens, which are mostly cleavage pieces, is not indicated.

U.S.N.M. 77517                      Mineral Hill, Conconully Dist., Okanogan Co., Washington.                       $2H_1$

Molybdenite occurs as coarse flakes in a pegmatite in a granite (Jones, 1917).

#### B. North America, exclusive of U.S.A.

The following is a short description of the Canadian occurrences (Vokes, 1963). Deposits of molybdenite occur in a belt from southern Gatineau and Pontiac Counties in Quebec through Renfrew and Haliburton Counties in Ontario in rocks, apparently of metasomatic origin, in the Grenville province of the Canadian shield. Molybdenite is associated with pyroxenite bodies, consisting mainly of diopside with some scapolite, plagioclase, and brown mica. The pyroxenite generally is included in or adjoins the Grenville limestone. Pegmatites sometimes occur with the pyroxenites. The genesis of the pyroxenite is disputed, some favoring a magmatic origin, others a metamorphic-metasomatic origin. It has not been proven that there is any genetic connection between the pyroxenites and the accompanying iron-sulfur-molybdenum mineralization. The molybdenite, the pyrite and/or pyrrhotite, and sometimes small amounts of chalcopyrite, occur as veins in breccia fillings in the pyroxenites.

Harvard 81617	Pontiac Co., Quebec	$2H_1$
Harvard 4333	}                      Renfrew Co., Ontario	$2H_1$
Harvard 4334		$2H_1$
Harvard 91499		$2H_1$
Harvard 95182		Ross Township, Renfrew Co., Ontario
Harvard 4651	Enterprise, Ontario	$2H_1$

It has not been possible to locate this mine or prospect. However, since in the study

specimen molybdenite occurs together with scapolite and pyrite, it may be assumed this is a "pyroxenite occurrence."

Harvard 92168            Spain Mine, Dacre, Ontario             $2H_1$

This deposit, treated in some detail by Vokes (1963) and Eardley-Wilmot (1925) belongs to the general pyrite-pyrrhotite-molybdenite mineralization of the region, but here the sulfides have been deposited as vertical veins in joints within the monzonite-gneiss country rock.

The following Canadian deposits represent other genetic types.

Harvard 90825            Don Rouyn Mine, Rouyn Township,     $2H_1$   
Quebec.

In this specimen molybdenite is associated with chalcopyrite. Molybdenite has been observed as vein filling in a fracture of the local Powell granite (Wilson, 1941, p. 73). No information was obtained "regarding the relationship of this occurrence of molybdenite to that of chalcopyrite, for the development of which the shaft was sunk."

Harvard 97213            Algonquin National Park, Ontario.     $2H_1$

This study specimen of molybdenite is a cleavage piece. No details concerning the locality are known.

Harvard unnumbered    Burnt Hill Brook, Miramichi River, New     $2H_1$   
Brunswick.

This is mainly a tungsten deposit of greisen type with cassiterite. Molybdenite is disseminated sparsely in both greisen and veins (Little, 1959).

Harvard 10193            Cananea, Sonora, Mexico             $2H_1$   
Harvard unnumbered    Sonora, Mexico             $2H_1+3R\sim 1:1$

In this porphyry copper deposit molybdenite occurs in three ways (Vanderwilt, 1942):  
(1) Ore consisting of copper sulfides with molybdenite in an altered porphyry.  
(2) Surrounding the ore, a zone of vein quartz with molybdenite and copper sulfides.  
(3) Below the ore, quartz with molybdenite and pyrite.

Specimen 101931 probably is from the third type of occurrence, while the unnumbered specimen may come from the first or second type.

Harvard 81618            Tortola Island, Virgin Islands, West Indies     $2H_1$

No occurrence of molybdenite is reported on Tortola Island (Martin-Kaye, 1959), though pegmatites and traces of copper mineralization have been observed. However, on the neighboring island, Virgin Gora, molybdenite, occurring in quartz veins with copper minerals, has been mined.

C. *South America*

Information on the Bolivian localities came from Ahlfeld and Schneider-Scherbina (1964).

Harvard 104637      Fenomenal, Larecaja, Bolivia.      2H<sub>1</sub>

Tin-bearing pegmatites occur here.

Harvard 91637      Challana, Cordillera Real, Bolivia.      2H<sub>1</sub>

Molybdenite occurs in this region with pegmatites, some containing muscovite, cassiterite, Li-phosphates and sulfides.

Harvard 103802      Murarata, Murillo y Sur Yungas, Bolivia.      2H<sub>1</sub>

Greisen and micropegmatites with wolframite and cassiterite occur here.

Harvard 98704      Mine Andina, Murillo, Bolivia      2H<sub>1</sub>

Quartz veins with tungsten and tin minerals occur here. Sulfides are found, also, together with fluorite.

Harvard 93068      Brazil      2H<sub>1</sub>

The study specimen is a cleavage piece. The locality designation is too indefinite to obtain information on genesis.

Harvard 81619	}      Santiago, Chile	3R
Harvard 4281		"Chile"

The exact localities are unknown. In both specimens molybdenite occurs with sulfides. They probably belong to the "porphyry copper" type, which is common in Chile.

Harvard unnumbered      Ricran, Jauja, Dept. Junin, Peru.      2H<sub>1</sub>

Quartz molybdenite veins occur in a granite (Petersen, personal communication).

D. *Europe*

Harvard 85064      Dalen Mine, Telemark, Norway.      2H<sub>1</sub>+3R~4:1



Molybdenite occurs in a quartz vein (Bugge, 1963, p. 92).

Harvard 85065            Bandakslı Mine, Telemark, Norway             $2H_1$

Molybdenite occurs in quartz veins and quartz-pegmatite veins, sometimes with fluorite (Bugge, 1963, pp. 90-92).

Harvard unnumbered    Frikstad, Iveland, Norway.             $2H_1$

Molybdenite occurs in a pegmatite.

Harvard 84610            Knaben Mine, Flekkefjord, Norway.             $2H_1$

Molybdenite is mined here and has many types of occurrences; in quartz veins, in pegmatites, and in impregnations of the country bed-rock (Bugge, 1963).

Harvard 91575            Norway             $2H_1$

The study specimen is only in crystals. The locality designation is too indefinite to obtain details of genesis.

Harvard 85066            Pielisjärvi, Finland             $2H_1$

Most probably the study specimen is from the Mätäsvaara Mine in the parish of Pielisjärvi (Kranck, 1945). The enrichment of molybdenite is connected closely with pegmatitic intrusions and quartz veins from a microcline granite, which has intruded a gneiss. The gneiss itself is silicified where molybdenite is concentrated. Other sulfides are present only in very small amounts.

U.S.N.M. R7478            Striegau, Silesia             $2H_1$

Molybdenite occurs in the granite of the Fuchsberge at Striegau. (Hintze, 1904).

Harvard 4222	} Altenberg	$2H_1+3R\sim 1:1$	
Harvard 81623		} Saxony	$2H_1+3R\sim 1:1$
Harvard unnumbered			$2H_1$

Small amounts of molybdenite, formed during the pneumatolytic stage, are found together with quartz, mica, topaz, cassiterite, arsenopyrite, native bismuth and wolframite (Cissarz, 1927-28).

Harvard 81622            Sadisdorf, Dippoldiswalde, Saxony             $2H_1$

Small amounts of pneumatolytic molybdenite occur together with cassiterite, Li-mica and arsenopyrite in quartz veins (Cissarz, 1927-1928).

Harvard 88914            Schlaggenwald, Bohemia             $2H_1$

Molybdenite is rare in the greisen (Cissarz, 1927-28).

Harvard 4291	} Zinnwald, Bohemia	$2H_1$
Harvard 4301		$2H_1+3R\sim 1:1$
Harvard 81624		$2H_1+3R\sim 1:1$
Harvard 105884		$2H_1$
Harvard unnumbered		$2H_1$

Molybdenite has not been found at this particular locality (Hintze, 1904 and Cissarz, 1927-28), but the study specimens probably come from some of the many localities in the neighborhood of Zinnwald and may all belong to the same group of greisen occurrences.

Harvard 4261            Schmirn, Tyrol, Austria.             $2H_1$

Molybdenite occurs in quartz veins in gneiss (granitic rocks?) in this valley (Hintze, 1904, p. 414).

U.S.N.M. 78904            Hallein, Austria             $2H_1$

There is no information in the literature that molybdenite has been found around this salt deposit, and the locality designation is, therefore, in doubt.

U.S.N.M. R 411            Sasca Montană, Romania             $2H_1+3R\sim 1:4$

The locality is given, also, as Szazka, Banat, Hungary.

Here the molybdenite occurs in both skarn and copper porphyry type deposits associated with a grandiorite (banatite) intrusion. Chalcopyrite is the dominant ore mineral with molybdenite, bornite, pyrite, digenite, tetrahedrite, pyrrotite, sphalerite and galena (Superceanu, 1969).

U.S.N.M. 94296	} Castangias, Sardinia	$2H_1$
Harvard 88814		Iglesias, Sardinia

The two specimens probably are from the same locality, since the name Iglesias often is used for the whole district. At Castangias molybdenite occurs in pegmatite veins in a granite about 20 km N.E. of Iglesias (Vardabasso, 1950, Fig. 64 and section in Fig. 65).

Harvard 103580	Serra de Gerez	$2H_1$
Harvard 103581	Portugal	$2H_1$

Nothing about the molybdenite of this district has been found in the literature, but from information regarding the tungsten mineralizations it is probable that molybdenite occurs in pegmatites or greisen.

#### E. Africa

Harvard 104686 } U.S.N.M. 112773 }	Azegour, Morocco	$2H_1$ $2H_1$
---------------------------------------	------------------	------------------

Molybdenite is found only in a metamorphosed garnet-rich limestone near a contact with granitic rocks (Von der Weid, 1941). In this skarn deposit accessory minerals are hematite, pyrite, pyrrhotite, galena and arsenopyrite.

Harvard 85173	"S.W. Africa"	$2H_1+3R\sim 9:1$
---------------	---------------	-------------------

The locality designation is indefinite and the study specimen is a crystal, so no information concerning genesis can be given.

Harvard 96916	Tsumeb, S.W. Africa	$2H_1+3R\sim 9:1$
---------------	---------------------	-------------------

In this well-known mine molybdenite is found disseminated in graphitized dolomite, as small particles together with green sphalerite, germanite, and renierite (Söhnge, 1964).

Harvard 93069	Olionsongate, S.W. Africa	$2H_1+3R\sim 9:1$
---------------	---------------------------	-------------------

"Olionsongate" is either a misspelling or a transliteration of the deposit Otjonzonjati (Westphal, 1914; Martin, 1965). Molybdenite occurs here sparsely, together with chalcopyrite and other sulfides in mineralized quartz and quartz feldspar breccia veins in biotite schists.

#### F. Asia

U.S.N.M. 76330	Colombo, Ceylon	$2H_1$
----------------	-----------------	--------

No information was obtained for this locality. This label may refer to some other locality in Ceylon.

U.S.N.M. 98099	Chicoi River, Transbaikalia, U.S.S.R.	$2H_1$
----------------	---------------------------------------	--------

Molybdenite occurs here in greisen and in quartz veins close to granites (Von zur Mühlen, 1926).

U.S.N.M. 88971      Chekiang, China       $2H_1+3R\sim 1:2$

Pegmatite veins containing molybdenite occur in great number in this region (Wong, 1920).

U.S.N.M. 105878      Mt. Kongo, Korea       $2H_1$

This locality is identical with the Kumgang molybdenum mine (Gallagher, 1963, p. 19), where molybdenite occurs in flakes up to 5 cm across in vertical quartz veins in a biotite granite. Wolframite also is found in the veins.

Harvard 108539      Hirasi mine, Gifu Prefecture, Japan.       $2H_1$

The deposit is in quartz veins ranging from the hydrothermal to the pegmatite stages; also, as dissemination in granitic rocks (Geol. Survey Japan, 1960). In addition to molybdenite the ore contains small amounts of pyrite and rare sphalerite and native bismuth.

U.S.N.M. 61448      Shirakawa, Hida, Japan.       $2H_1$

This is probably the same locality as that for specimen Harvard 108539, since the Hirase mine is situated on Shirakawa (Geol. Survey Japan, 1960).

U.S.N.M. 47143      Kosha, Japan.       $2H_1$

This may be a misspelling or a confused transliteration, for no mention of this locality has been found in the literature.

#### G. Australia

In New South Wales so-called pipes occur in many granite bodies (Garrety, 1953) and their chief minerals, besides quartz, are molybdenite, native bismuth, wolframite, arsenopyrite, chalcopyrite, cassiterite, and tetradymite.

U.S.N.M. 48788      New England District, N.S.W.       $2H_1+3R\sim 9:1$

The locality designation of this specimen is indefinite.

Harvard 88884	Kingsgate, New England District N.S.W.	$2H_1$
Harvard 4242		$2H_1$
Harvard 90314		$2H_1$






FIG. 1. Theoretical mixtures of the  $2H_1+3R$  polytypes of molybdenite.

the formation of  $3R$  could be promoted by a slower decrease of temperature or by the presence of mineralizers.

Of the 108 specimens in this investigation, about 80 percent were the  $2H_1$  polytype. The majority of  $2H_1$  samples came from quartz veins or pegmatites associated with granitic rocks or granite contacts. Minor amounts of sulfides, such as pyrite and chalcopyrite, accompanied the

molybdenite in some of these deposits.  $2H_1$  has been found, however, in every geological environment known for molybdenite. Three specimens were of the  $3R$  polytype, two of them from the same locality, probably a porphyry copper deposit. In the third specimen the molybdenite occurred with quartz in a muscovite biotite-schist. Nineteen specimens were mixtures of  $2H_1+3R$ . A ratio of  $2H_1$  to  $3R$  is given for each sample (see description of the localities). This proportion is only approximate and varies, though not greatly, in diffractometer runs on different samples of the same specimen. In Figure 2, X-ray diffraction patterns are shown for  $2H_1$ ,  $3R$ , and some mixtures of these two polytypes found in the natural material. By comparison of these patterns with the theoretical diagram in Part I of this study it is easy to distinguish between the mixtures of  $2H_1+3R$  and polytypes of higher symmetry. In half of the mixtures the  $2H_1$  polytype was dominant, and these specimens came generally from quartz veins, pegmatites, or granite contacts. The specimens with  $3R$  dominant, or present in a significant amount, came from a variety of geological environments. From the present study, then, a definite correlation between genesis and polytype occurrence can not be made, though copper mineralization seems to play an important role. No new modifications of naturally-occurring molybdenite were found, but study of more deposits may yield other polytypes.

The growing interest in molybdenite has been spurred by the search for new sources of the rare element rhenium, for which molybdenite serves, to date, as the most important host. Fleischer (1959) thought that "no generalizations are yet possible as to correlations of rhenium content with geological conditions of formation," but in 1960 he pointed out newer analyses (Zhirov and Ivanova, 1959) suggested "that the content of rhenium in molybdenites increases as one goes from high-temperature to medium-temperature deposits." There were, however, unexplained large variations of rhenium concentration from porphyry copper deposits. Since 1960 many investigations have been made (largely in U.S.S.R.) of rhenium enrichment in molybdenite. The results of these studies, summarized by Tischendorf *et al.* (1968), lead more and more to the conclusion that the enrichment of rhenium is greatest in copper-molybdenum formations. It is of interest to note that analyses of molybdenite from porphyry copper type deposits of Sasca-Montană, Romania (Superceanu, 1969) yielded as much as 0.13 percent rhenium, and this investigation's specimen from that locality (U.S.N.M.R 411) is a mixture of  $2H_1+3R \sim 1:4$  (Figure 2). A correlation between the presence of copper mineralizers with reference to rhenium enrichment and the occurrence of  $3R$  and  $2H_1+3R$  mixtures of molybdenite is




FIG. 2. X-ray diffraction patterns<sup>1</sup> of some naturally-occurring molybdenites.

- 1.) Harvard unnumbered, Mt. Lion Mine, Colorado;  $2H_1$
- 2.) Harvard 96916;  $2H_1:3R \sim 9:1$
- 3.) Harvard 4301;  $2H_1:3R \sim 1:1$
- 4.) U.S.N.M. R411;  $2H_1:3R \sim 1:4$
- 5.) Harvard 81619;  $3R$

<sup>1</sup> Copper radiation, with nickel filter.



suggested. This is supported by the recent work by Zelikman *et al.* (1970). It may well be that additional finds of 3R molybdenite will lead to new sources of rhenium.

## ACKNOWLEDGMENTS

This study was supported by A.E.C. Contract AT(301)-1403, at Harvard University and a grant-in-aid from the Joint Committee on Powder Diffraction Standards of the American Society for Testing and Materials to The Pennsylvania State University. Gratitude is extended to Professor Clifford Frondel of Harvard University for suggesting the problem and stimulating the research.

## REFERENCES

- AHLFELD, F., AND A. SCHNEIDER-SCHERBINA (1964) Los yacimientos minerales y de hidrocarburos de Bolivia. *Bolivia Dept. Nat. Geol. Bol.* 5.
- BAYLEY, W. S. (1910) Iron mines and mining in New Jersey. *New Jersey Geol. Surv. Final rep., series VII.*
- BUGGE, A. (1963) Norges molybdenforekomster. *Norges Geol. Undersokelse*, 217.
- BUTLER, B. S., G. F. LOUGHLIN, H. C. HEIKES, *et al.* (1920) The ore deposits of Utah. *U.S. Geol. Surv., Prof. Pap.* 111.
- CHUKHROV, F. W., B. B. ZVYAGIN, L. P. ERMILOVA, S. A. SOBELOVA, AND V. G. KHITROV (1968) Polytypes of molybdenite and their occurrence in pres. *Geol. Rud. Mestorozhd.* 10 (2), 12-26. [Transl. *Int. Geol. Rev.* 12, 74-86 (1970)]
- CISSARZ, A. (1927-28) Übergangslagerstätten innerhalb der intrusiv-magmatischen Abfolge. Teil I. Zinn-Wolfram-Molybdänformationen. *Neues Jahrb. Mineral. Beil.* 56A, 99-274.
- CREASY, S. C. (1957) Geology and resources. In W. McInnes, Molybdenum. A materials survey. *U.S. Bur. Mines Inform. Circ.* 7784.
- EARDLEY-WILMOT, V. L. (1925) Molybdenum. *Can. Dept. of Mines, Mine Branch Rep. No.* 592.
- EGGLESTON, J. W. (1918) Eruptive rocks at Cuttingsville, Vermont, *Amer. J. Sci. 4th Ser.* 45, 377-410
- ENDLICH, F. M. (1878) Catalogue of minerals found in Colorado. *U.S. Geol. and Geogr. Surv. Terr. (Hayden). Ann. Rep.* 10, 61-131.
- FLEISCHER, M. (1959) The geochemistry of rhenium, with special reference to its occurrence in molybdenite. *Econ. Geol.* 54, 1406-1413.
- (1960) The geochemistry of rhenium-addendum. *Econ. Geol.* 55, 607-609.
- GALBRAITH, F. W. (1947) Minerals of Arizona. 2nd ed. *Ariz. Bur. Mines Bull.* 153.
- GALLAGHER, D. (1963) *Mineral Resources of Korea. Vol. 5. Tungsten and Molybdenum.* Mining Branch, Industry and Mining Division, U. S. Operations Mission to Korea in cooperation with Geological Survey., Rep. of Korea.
- GARRETTY, M. D. (1953) Bismuth-molybdenite pipes of New England. In A. B. Edwards, ed., *Geology of Australian Ore Deposits.* Australian Institute of Mining and Metallurgy, Melbourne.
- GEOLOGICAL SURVEY OF JAPAN (1960) *Geology and Mineral Resources of Japan. 2nd. ed.* Dai-Nippon Printing Co., Ltd. Tokyo.
- HINTZE, C. (1904) *Handbuch der Mineralogie*, 1, Veit and Comp., Leipzig.
- HORTON, F. W. (1916) Molybdenum; its ores and their concentration. *U.S. Bur. Mines Bull.* 111.

- HUSSEY, A. M., II, *et al.* (1958) Maine metal mines and prospects. *Maine Geol. Surv. Minerals Res. Index*, 3.
- JACKSON, C. (1844) *Final Report on the Geology and Mineralogy of the State of New Hampshire*. Concord, N.H., 57 p.
- JONES, E. E. JR. (1917) Reconnaissance of the Conconully and Ruby Mining Districts, Washington, *U.S. Geol. Surv. Bull.* 640, 11-36.
- KHURSHUDYAN, E. KH. (1966) Conditions of formation of the rhombohedral modification of molybdenite. *Dokl. Akad. Nauk SSSR*, 171, 186-189.
- , L. A. ARUTYNNYAN AND B. M. MELIKSETYAN (1969) Genesis of molybdenite polytypes. *Geokhimiya*, 10, 1200-1209 [Transl. *Geochem. Intern.*, 6, 924-949.]
- KILLEFER, D. H., AND A. LINZ (1952) *Molybdenum Compounds*. Interscience Publishers, N.Y.-London.
- KING, R. V. (1966) Molybdenum. *Calif. Div. Mines and Geol. Bull.* 191, 264.
- KIRKEMO, H., C. A. ANDERSON, AND S. C. CREASY (1965) Investigations of the molybdenum deposits in the conterminous United States 1942-60. *U.S. Geol. Surv. Bull.* 1182-E.
- KRANCK, E. H. (1945) The molybdenum deposit at Mätäsvaara in Carelia (E. Finland). *Geol. Foren. Forh.* 67, 325-350.
- LITTLE, H. W. (1959) Tungsten Deposits of Canada. *Geol. Surv. Canada, Econ. Geol. Ser.* 17.
- MARMEDOW, Z. M. (1965) Geochemistry of rhenium in copper-molybdenum minerals of Paragachai. *Dokl. Akad. Nauk. Azerb. SSR*, 21, 24-27.
- MARTIN, H. (1965) *The Precambrian Geology of Southwest Africa and Namaqualand*. The Precambrian Research Unit. Univ. Cape Town.
- MARTIN-KAYE, P.H.A. (1959) *Reports on the Geology of the Leeward and British Virgin Islands*. Voice Publishing Co. Ltd. St. Lucia.
- NEWLAND, D. H. (1921) The mineral resources of the State of New York. *N. Y. State Mus. Bull.* 223 and 224.
- POSTEL, A. W. (1940) Hydrothermal emplacement of granodiorite near Philadelphia. *Proc. Acad. Natur. Sci. Philadelphia*, 123-152.
- SCHILLING, J. H. (1962) An Inventory of Molybdenum Occurrences in Nevada. *Nev. Bur. Mines Rep.* 2, 25.
- SÖHNGE, P. G. (1964) The geology of the Tsumeb Mine. *In Some Ore Deposits in Southern Africa* 2. Geol. Soc. S. Africa, Johannesburg, 367-382.
- STEWART, D. B. (1956) *Rapakivi Granite of the Deer Isle Region Maine*. Harvard Univ. Ph.D. Diss.
- SUPERCEANU, C. I. (1969) The copper ore deposits of Sasca Montană in south-western Banat and their position in the Alpine-Eastern Mediterranean copper-molybdenum Belt. *Geol. Rundsch.* 58, 798-861.
- TISCHENDORF, G., W. LIMPACK, AND H. UNGETHUEM (1968) the geochemistry of rhenium—progress and new results. *Geologie*, 17, 1208-1218.
- U.S. GEOL. SURV. (1966) *Mineral and Water Resources of Washington*. U.S. Senate; Committee on Interior and Insular Affairs, 89th Congress, 2nd Session.
- VANDERWILT, J. W. (1942) The occurrence and production of molybdenum. *Quart. Colo. Sch. Mines* 37, No. 4.
- VARDABASSO, S. (1950) Geologia dei giacimenti ercinici piombozinciferi della Sardegna. *Intern. Geol. Congress, 18th, Great Britain. Part VII, Sec. F.*, p. 257-265.
- VOKES, F. M. (1963) Molybdenum deposits of Canada. *Geol. Surv. Canada, Econ. Geol. Rep.* 20.
- VON DER WEID, F. (1941) *Le Gisement de Molybdénite d'Azégour et la Région des Guedmiousa*. Atlas de Marrakech, Maroc, George & Cie. Geneva.
- VON ZUR MÜHLEN, L. (1926) *Die Lagerstätten von Wolfram, Zinn und Molybdän in Russland*. E. Schweizerbart'sche Verlagsbuchhandlung. Stuttgart.

- VORMA, A., P. KALLIO, AND K. MERILÄINEN (1966) Molybdenite -3R from Inari, Finnish Lapland. *C. R. Soc. Geol. Finlande*, **38**, 67-68.
- WARREN, C. H., AND H. E. MCKINSTRY (1924) The granites and pegmatites of Cape Ann, Massachusetts. *Proc. Amer. Acad. Arts Sci.* **59**, 315-357.
- WESTPHAL, H. (1914) Die Kupfererz-Pegmatite von Otjonzonjati. (Deutsch-Südwest Afrika). *Z. Prakt. Geol.* **22**, 385-416.
- WILSON, E. D. (1941) Tungsten deposits of Arizona. *Univ. Arizona Bull.* **17**, No. 2.
- WILSON, M. E. (1941) Noranda District, Quebec. *Geol. Surv. Canada Mem* **229**.
- WONG, WEN-HAO (1920) Les provinces métallogéniques de Chine. *Bull. Geol. Surv. China*, **2**, 37-59.
- WORCESTER, P. C. (1919) Molybdenite deposits of Colorado. *Colo. Geol. Surv. Bull.* **14**, 32.
- WRIGHT, C. W. (1915) Geology and ore deposits of Copper Mountain and Kasaan Peninsula. *U.S. Geol. Surv. Prof. Pap.* **87**, 49.
- ZELIKMAN, A. N., G. V. INDENBAUM, M. V. TESLITSKAYA, AND V. P. SHALANKOVA (1969) Structural transformations in synthetic MoS<sub>2</sub>. [*Kristallografiya* **14** (1959) 795-799. Transl. *Soviet Physics-Crystallography* (1969) **14**, 687-691]
- ZHIROV, K. K., AND G. F. IVANOVA (1959) Distribution of rhenium in molybdenites from deposits in a number of genetic types. *Geokhimiya*, **1959**, 518-523.

*Manuscript received, May 22, 1970; accepted for publication, July 28, 1970.*