

RICKER, R. W., AND E. F. OSBORN (1954) Additional phase equilibrium data for the system CaO-MgO-SiO₂. *Amer. Ceram. Soc. J.* **37**, 133-139.

SIMKIN, T., AND J. V. SMITH (1970) Minor-element distribution in olivine. *J. Geol.* **78**, 304-325.

WARNER, R. (1971) *Experimental investigations in the sys-*

tem CaO-MgO-SiO₂-H₂O. Ph.D. Thesis, Stanford University. 165 pp.

YANG, H. Y., AND W. R. FOSTER (1972) Stability of iron-free pigeonite at atmospheric pressure. *Amer. Mineral.* **57**, 1232-1241.

American Mineralogist, Volume 58, page 345, 1973

Apatite Chemistry and Phosphorus Fugacity in a Differentiated Igneous Intrusion: Correction

W. P. NASH

*Department of Geological and Geophysical Sciences,
University of Utah, Salt Lake City, Utah 84112*

Values of $\Delta G_r^\circ/2.303 RT$ for two reactions given by Nash (1972) are incorrect.

For the reaction $\text{Ca}_5(\text{PO}_4)_3\text{OH} + 5/2 \text{Mg}_2\text{SiO}_4 + \text{KFe}_3\text{AlSi}_3\text{O}_{10}(\text{OH})_2 + 1/2 \text{KAlSi}_3\text{O}_8 + 15/2 \text{SiO}_2 = 5 \text{CaMgSi}_2\text{O}_6 + 3/2 \text{Fe}_2\text{SiO}_4 + 3/2 \text{KAlSi}_2\text{O}_6 + 15/4 \text{O}_2 + 3/2 \text{H}_2\text{O} + 3\text{P}$ the correct value is

$$\frac{\Delta G_r^\circ}{2.303 RT} = \frac{168,508}{T} - 57.87(\pm 7.2).$$

For the reaction $\text{Ca}_5(\text{PO}_4)_3\text{OH} + 3 \text{MgAl}_2\text{O}_4 + 57/4 \text{SiO}_2 + 15/2 \text{Fe} = 15/4 \text{Fe}_2\text{SiO}_4 + 1/2 \text{Mg}_2\text{SiO}_4 + 2\text{CaMgSi}_2\text{O}_6 + 3\text{CaAl}_2\text{Si}_2\text{O}_8 + 1/2 \text{H}_2\text{O} + 3\text{P}$ the correct value is

$$\frac{\Delta G_r^\circ}{2.303 RT} = \frac{51,529}{T} - 20.65(\pm 6.9).$$

The calculated phosphorus fugacities are shifted to higher values of f_p . The equilibrium curves for these two reactions (Nash, 1972, Fig. 1) are shifted, respectively, by the values $\Delta \log f_p = 5478/T +$

$5.26 (\pm 2.4)$ and $\Delta \log f_p = 2730/T + 7.93 (\pm 2.3)$. It is clear, upon recalculation, that the effects of oxidation state and silica activity on the equilibria are not immediately evident, as previously thought. Although the data of Stormer and Carmichael (1971) indicate a decrease of phosphorus fugacity with decreasing magmatic silica activity, no such simple behavior is indicated by the new curve for Shonkin Sag laccolith which crosses theoretical basalt and rhyolite curves. Contrary to prior conclusions, reduction in oxygen fugacity apparently increases phosphorus fugacity values, as indicated by the equilibria for a basaltic assemblage coexisting with metallic iron.

References

- NASH, W. P. (1972) Apatite chemistry and phosphorus fugacity in a differentiated igneous intrusion. *Amer. Mineral.* **57**, 877-886.
- STORMER, J. C., AND I. S. E. CARMICHAEL (1971) Fluorine-hydroxyl exchange in apatite and biotite: a potential igneous geothermometer. *Contrib. Mineral. Petrology*, **31**, 121-131.