

Visible-Region Absorption Spectra of a Ti^{3+} Fassaite from the Allende Meteorite: A Discussion

ROGER G. BURNS, AND FRANK E. HUGGINS

Department of Earth and Planetary Sciences, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139

Abstract

The spectral data of Dowty and Clark for the titaniferous pyroxene in the Allende meteorite have been reinterpreted. Arguments are presented in support of an alternative assignment of the two absorption bands at $21,000\text{ cm}^{-1}$ and $16,500\text{ cm}^{-1}$ as both being due to crystal field transitions in Ti^{3+} ions. Not only are two such bands expected for Ti^{3+} ions in distorted $M1$ coordination sites of clinopyroxenes, but the derived spectral parameters α and Δ_0 are consistent with other $Ti(III)$ compounds. The occurrence of Ti^{3+} ions in the clinopyroxene structure is attributed to the enhanced crystal field stabilization energy acquired in the $M1$ site.

Introduction

Dowty and Clark (1973) have described the crystal chemistry of an intriguing titaniferous pyroxene from the Allende meteorite, to which they ascribe the chemical formula:



Absorption bands in the optical spectra were observed at $21,000\text{ cm}^{-1}$ and $16,500\text{ cm}^{-1}$ in good agreement with results obtained for titaniferous pyroxenes from Apollo 11 rocks (Burns, Abu-Eid and Huggins, 1972). Burns *et al* (1972) assigned the two bands at $21,500$ and $16,000\text{ cm}^{-1}$ in the lunar pyroxenes to crystal field transitions in Ti^{3+} . However, Dowty and Clark (1973) assigned only one band ($21,000\text{ cm}^{-1}$) in their spectra to a Ti^{3+} crystal field transition. The other band ($16,500\text{ cm}^{-1}$) they attributed to a Ti^{3+} - Ti^{4+} charge-transfer transition.

We wish to prove that the spectral assignment proposed by Dowty and Clark (1973) is unlikely and to demonstrate that *both* bands are due to crystal field transitions in Ti^{3+} , the excited state of which is split into two energy levels in a distorted coordination site. We shall also indicate the significance of the results to the electronic structure and stability of Ti^{3+} ions in clinopyroxene structures.

Interpretation of the Spectra

To support our interpretation that titanium-related spectral features of the Allende and Apollo 11 py-

roxenes in the visible region are due only to crystal field transitions in Ti^{3+} ions, we cite arguments based on (a) correlations with the spectra of Fe^{2+} ions in calcic clinopyroxenes; (b) analogies with the spectra of $Ti(III)$ salts and complexes; (c) the nature of the $M1$ coordination site in calcic clinopyroxenes; (d) crystal field states of Fe^{2+} and Ti^{3+} ions in the $M1$ sites; (e) positions of absorption bands in relation to the spectro-chemical series; and (f) intensities of absorption bands for metal-metal charge transfer transition.

(a) Correlations with Fe^{2+} Spectra in Calcic Clinopyroxenes

Absorption spectra of clinopyroxene minerals are often complicated by contributions from Fe^{2+} ions in two coordination sites, $M1$ and $M2$, and by Fe^{2+} - Fe^{3+} charge transfer bands. The spectra of diopsides-hedenbergites containing negligible Fe^{3+} ions and with Ca^{2+} ions completely filling $M2$ positions show two absorption bands centered at $9,600\text{ cm}^{-1}$ and $8,500\text{ cm}^{-1}$. The most intense band occurs at $9,600\text{ cm}^{-1}$ in the $b = \text{beta}$ polarized spectrum and is distinctly polarization dependent. The spectra represent crystal field transitions in Fe^{2+} ions located in $M1$ positions of the clinopyroxene structure. In calcium deficient pyroxenes additional bands around 1000 nm ($10,000\text{ cm}^{-1}$) and 2200 nm ($4,500\text{ cm}^{-1}$) appear from Fe^{2+} ions in the $M2$ positions, while the presence of Fe^{3+} ions leads to a charge transfer band centered between 700 and 800 nm (approximately

12,000–14,000 cm^{-1}). The latter feature is demonstrated by the spectra (Fig. 1) of a hedenbergite, $\text{Ca}_{0.95}\text{Mg}_{0.20}\text{Fe}_{0.85}\text{Si}_2\text{O}_6$, containing about 5 percent ferric iron. The spectra show a prominent Fe^{2+} – Fe^{3+} charge transfer band around 12,200 cm^{-1} . The most significant features of the iron clinopyroxene spectra to the present discussion are: (1) the pronounced polarization dependence of the Fe^{2+} crystal field spectra; (2) the presence of the Fe^{2+} – Fe^{3+} charge transfer band in all three polarized spectra; and (3) the 1000–1500 cm^{-1} splitting in the crystal field spectra of Fe^{2+} ions in $M1$ positions. Similar split-




FIG. 1. Polarized absorption spectra of a hedenbergite. The calcic clinopyroxene has the formula $\text{Ca}_{0.95}\text{Mg}_{0.20}\text{Fe}_{0.85}\text{Si}_2\text{O}_6$ and contains 25.78 percent FeO and 1.53 percent Fe_2O_3 . It is specimen 2B in Mueller (1960) *Amer. J. Sci.* **258**, 449. α spectrum; — β spectrum; - - - γ spectrum (Optic orientation: $\alpha:a = 34^\circ$; $\beta:b$; $\gamma:c = 48^\circ$).

tings are observed in the spectra of Ti(III) compounds.

(b) Analogies with the Spectra of Ti(III) Compounds

In a classical study of several hydrated Ti(III) compounds, Hartmann and Schläfer (1951) demonstrated that the spectra of octahedral $\text{Ti(H}_2\text{O)}_6^{3+}$ ions consist of an asymmetric absorption band with a prominent shoulder on the long wavelength side. The spectrum of cesium titanium alum, for example, was resolved into two peaks at 19,900 cm^{-1} and 18,000 cm^{-1} (Schläfer, 1964), indicating a splitting of about 1,900 cm^{-1} in the high energy crystal field states of Ti^{3+} . Similar energy separations of up to 4000 cm^{-1} have been observed in the absorption spectra of a variety of Ti(III) compounds (Hartmann *et al.*, 1957; Hoff and Brubaker, 1971; Clark, 1968, p. 153). Clark (1968, p. 156) cites one exceptionally large splitting, involving the $[\text{Ti(H}_2\text{O)}_4\text{F}_2]^+$ ion, which appears to be as much as 8000 cm^{-1} . Other examples are given in Table 1. Similar energy separations have also been demonstrated in simple Fe(II) compounds (Cotton and Meyers, 1960). Thus, asymmetric or double-maxima absorption bands are prevalent in the crystal field spectra of octahedrally coordinated Fe^{2+} and Ti^{3+} ions.

(c) Nature of the Clinopyroxene $M1$ Coordination Site

Warren and Bragg (1928) originally demonstrated that the oxygen environment about the $M1$ position in diopside is appreciably distorted from octahedral symmetry (Figure 2). This has been confirmed and demonstrated in recent crystal structure refinements of a variety of clinopyroxenes (Prewitt and Burnham, 1966; Freed and Peacor, 1967; Peacor, 1967; Christensen and Hazell, 1967; Clark *et al.*, 1969; Schropfer, 1971). In diopside, for example, metal-oxygen distances in the $M1$ site range from 2.115 Å to 2.050 Å (Table 2), while there is considerable variation of oxygen-oxygen distances and O-M1-O bond angles around the $M1$ "octahedron" (Clark *et al.*, 1969). The structural data cited by Dowty and Clark (1973) for the titaniferous pyroxene in the Allende meteorite further demonstrate the asymmetry of the $M1$ coordination site. The range of metal-oxygen distances, 2.120 to 1.993 Å, is almost twice that observed in the diopside $M1$ site, suggesting increased distortion of the Allende pyroxene $M1$ site (containing Ti^{3+}) compared to

diopside. Such distortion profoundly affects the energy levels of Ti^{3+} ions in the M1 site.

(d) Crystal Field States of Ti^{3+} and Fe^{2+} in a Distorted Coordination Site

The field-free Fe^{2+} and Ti^{3+} ions both have D spectroscopic ground terms, which give rise to T_{2g} ground states and E_g excited states when the ions are in octahedral coordination in an oxide or silicate structure (Figure 3). The electronic configuration of the Ti^{3+} ion, $[A]3d^1$, is particularly simple in an octahedral crystal field. The ground state configuration, $(t_{2g})^1(e_g)^0$, is three-fold degenerate and is designated ${}^2T_{2g}$. The single 3d electron is excited into an e_g orbital by visible radiation, giving the excited state configuration $(t_{2g})^0(e_g)^1$ which is doubly degenerate and is designated by 2E_g . As a result of the dynamic Jahn-Teller effect (Cotton and Meyers, 1960), E_g excited states are resolved into two levels. Transitions to these levels produce asymmetric or two-peaked spectra in the visible region.

The crystal-field states are split further when the environment about the Ti^{3+} ions is distorted from octahedral symmetry. In low-symmetry environments, such as the clinopyroxene M1 site (Figure 2), the t_{2g} and e_g orbitals are resolved into a maximum of three and two energy levels, respectively (Figure 4). The representative spectral data of Ti(III) complexes summarized in Table 1 show typical splittings (α) for the e_g orbitals to be up to 4000 cm^{-1} . Splittings are smaller for the t_{2g} orbitals; magnetic data indicate β values (Figure 4) of up to 500 cm^{-1} . (Clark, 1968, p. 154).

The ground state configuration of Ti^{3+} corresponds to electron occupancy of the lowest energy t_{2g} -type orbital. Visible light excites this electron to each of the orbitals of the e_g -type orbital level, while transitions to the upper level produce the band at $21,000\text{ cm}^{-1}$ (Figure 4). The energy separation, $4,500\text{ cm}^{-1}$, is thus a direct consequence of the splitting of the e_g -type orbitals of Ti^{3+} in the distorted M1 site. Note that the baricenter of the e_g levels of Ti^{3+} ions in the Allende pyroxene is approximately $18,750\text{ cm}^{-1}$ above the ground state (Figure 4), which is identical to that derived for Ti^{3+} in the Apollo 11 clinopyroxenes (Table 1).

The electronic configurations and crystal field states of Fe^{2+} , $3d^6$, are similar to Ti^{3+} (Figure 3). The quintet ${}^5T_{2g}$ state corresponds to the ground-state configuration $(t_{2g})^4(e_g)^2$. The sixth 3d elec-




FIG. 2. (100) projection illustrating the distortion of the M1 coordination site in clinopyroxenes. Metal-oxygen and oxygen-oxygen distances are in Å. Data from Clark *et al* (1969).

tron of Fe^{2+} occupying a t_{2g} orbital behaves like the single 3d electron of Ti^{3+} , giving rise to analogous absorption bands around 1000 nm in the near infrared region (Figure 1). The smaller energy separation ($1000\text{--}1500\text{ cm}^{-1}$) between the split e_g -type orbital energy levels of Fe^{2+} in diopside is due partly to the reduced distortion of the diopside M1 site compared to that in the titanian fassaite and partly to the lower charge on the Fe^{2+} ion.

In summary, double-band crystal field spectra are predicted and observed for both Fe^{2+} and Ti^{3+} in




FIG. 3. Schematic energy level diagram for $Ti^{3+}(d^1)$ and $Fe^{2+}(d^6)$ in an octahedral crystal field. (a) Spectroscopic ground terms (b) Splitting of D terms in octahedral crystal field. (c) Crystal field states in octahedral coordination. (d) (e) Electronic configurations of the ground state and excited states of Ti^{3+} and Fe^{2+} , respectively.




FIG. 4. Energy levels of 3d orbitals of Ti^{3+} . (a) Typical values for $Ti(III)$ oxide compounds. (b) Values derived from absorption spectra of the titaniferous pyroxene in the Allende meteorite. All energies in cm^{-1} units.

distorted $M1$ sites of calcic clinopyroxenes. Thus, we believe that the bands at $21,000\text{ cm}^{-1}$ and $16,500\text{ cm}^{-1}$ reported by Dowty and Clark (1973) both represent crystal field transitions in Ti^{3+} .

(e) Positions of Absorption Bands in Relation to the Spectrochemical Series

The octahedral crystal field splitting parameter, Δ_o , for all transition metal ions depends on the type of anion or ligand coordinated to the cation. An abbreviated order of increasing Δ_o (that is, the spectrochemical series) is: halides $<$ H_2O and oxides $<$ NH_3 and amides $<$ cyanides. Values of Δ_o are derived experimentally from positions of absorption bands. Examples for $Ti(III)$ compounds are given in Table 1. Note that Δ_o for cyanide complexes is about $20,600\text{ cm}^{-1}$, which exceeds the values for oxygen complexes by about 2000 cm^{-1} .

Dowty and Clark (1973) assign only the $21,000\text{ cm}^{-1}$ band in the spectrum of the Allende pyroxene to a crystal field transition in Ti^{3+} ions. This implies that Δ_o for Ti^{3+} in this pyroxene is approximately $21,000\text{ cm}^{-1}$ or higher, which exceeds the value for $Ti(III)$ cyanide complexes. Clearly, this is at variance with trends shown in the spectrochemical series for other transition elements. If the band at $21,000$

cm^{-1} in the Allende pyroxene is assigned to a crystal field transition in Ti^{3+} , then there *must* (group theoretical selection rules permitting) be a second crystal field band located at lower energy, such as $16,500\text{ cm}^{-1}$, in order to lower the value of Δ_o to a value expected for Ti^{3+} coordinated to oxygen. Therefore, the spectrochemical series provides another argument in support of the assignment of the $16,500\text{ cm}^{-1}$ band in the Allende pyroxene to a crystal field transition in Ti^{3+} ions.

(f) Band Intensities for Metal-Metal Charge Transfer Transitions

Dowty and Clark (1973) argue from the polarization dependence of the band at $16,500\text{ cm}^{-1}$ that this band represents a charge transfer transition between Ti^{3+} and Ti^{4+} ions in the neighboring $M1$ positions of the Allende pyroxene. This assignment may be negated by reference to the $Fe^{2+}-Fe^{3+}$ charge transfer band in hedenbergite (Fig. 1). The chemical analysis of the hedenbergite shows 25.78 percent FeO and 1.53 Fe_2O_3 (J. H. Scoon, personal communication, in Bancroft *et al.*, 1967), giving a $Fe^{3+}:Fe^{2+}$ ratio of 1:20. In spite of the small proportion of Fe^{3+} ions, and therefore small probability of adjacent $Fe^{2+}-Fe^{3+}$ ion pairs, the charge-transfer band around $12,000\text{ cm}^{-1}$ has an intensity comparable to that of the Fe^{2+} crystal field transition around $10,000\text{ cm}^{-1}$. This conforms with group theory selection rules which predict greater probability of a charge transfer transition relative to a $d \rightarrow d$ crystal field transition. Note, also, that the charge transfer band in hedenbergite is not polarization dependent.

The chemical formula of the Allende fassaite derived by Dowty and Clark (1973) shows a $Ti^{4+}:Ti^{3+}$ ratio of approximately 2:7, indicating a considerably higher probability of Ti^{3+} and Ti^{4+} ion pairs in adjacent $M1$ positions than of $Fe^{2+}-Fe^{3+}$ ion pairs in the hedenbergite. If the $16,500\text{ cm}^{-1}$ band were a $Ti^{3+}-Ti^{4+}$ charge-transfer transition, its intensity would have to be considerably higher than the Ti^{3+} crystal field band proposed at $21,000\text{ cm}^{-1}$. Furthermore, by analogy with the $Fe^{2+}-Fe^{3+}$ charge transfer band in hedenbergite, the $16,500\text{ cm}^{-1}$ band should not show polarization dependence. However, by analogy with the crystal field spectra of Fe^{2+} ions shown in Figure 1, both the relative intensity and polarization dependence of the $16,500\text{ cm}^{-1}$ band of the Allende pyroxene may be readily explained as crystal field transitions in Ti^{3+} ions.

TABLE 1. Crystal Field Spectra of Ti(III) Compounds

| Compound | Absorption Bands (cm^{-1}) | | Splitting of e_g levels, (cm^{-1}) | Approx. Δ_o (cm^{-1}) | References |
|---|--|--------|--|--|------------|
| $\text{K}_3\text{Ti}(\text{CN})_6 \cdot 2\text{KCN}$ | 22,300 | 18,900 | 3,400 | 20,600 | 1 |
| Ti^{3+} -doped Al_2O_3 | 20,300 | 18,450 | 1,850 | 19,375 | 2 |
| $\text{CsTi}(\text{SO}_4)_2 \cdot 12\text{H}_2\text{O}$ | 19,900 | 18,000 | 1,900 | 18,950 | 3 |
| $\text{Ti}(\text{H}_2\text{O})_6\text{Cl}_3$ | 20,100 | 17,000 | 3,100 | 18,575 | 4 |
| $(\text{NH}_4)_3[\text{TiF}_6]$ | 19,000 | 15,100 | 3,900 | 17,050 | 5 |
| $\alpha\text{-TiCl}_3$ | 13,800 | 12,000 | 1,800 | 12,900 | 6 |
| Apollo 11 pyroxene (10047) | 21,500 | 16,000 | 5,500 | 18,750 | 7 |
| Allende fassaite | 21,000 | 16,500 | 4,500 | 18,750 | 8 |
| $\text{NaTiSi}_2\text{O}_6$ | $\approx(22,000)^*$ | 15,600 | $\approx(6,300)^*$ | (18,750)* | 9 |

* Inferred, see text.

1. H.L. Schläfer and R. Götz, *Z. anorg. allgem. Chem.*, **309**, 104 (1961).
2. D.S. McClure, *J. Chem. Phys.*, **36**, 2757 (1962).
3. H.L. Schläfer, *Proc. Sump. Theory Struct. Complex Comp.*, Wrocław, 1962 Pergamon, Press, p. 181.
4. H. Hartmann, H.L. Schläfer and K.H. Hansen, *A. anorg. allgem. Chem.*, **284**, 153 (1956).
5. H.D. Bedon, S.M. Horner and S.Y. Tyree, *Inorg. Chem.*, **3**, 647 (1964).
6. R.J.H. Clark, *J. Chem. Soc.*, 417 (1964).
7. R.G. Burns, R.M. Abu-Eid and F.E. Huggins, *Proc. 3rd Lunar Sci. Conf.*, Suppl. 3, vol. 1, 533-543 (1972).
8. E. Dowty and J.R. Clark, *Amer. Min.*, **58**, (1973).
9. C.T. Prewitt, R.D. Shannon and W.B. White, *Contr. Mineral. Petrol.*, **35**, 77 (1972).

Discussion

The arguments presented in the previous section not only demonstrate that two crystal-field absorption bands are to be expected for Ti^{3+} ions in distorted six-coordinate sites, but also invalidate the assignment of the band centered between 15,500–16,500 cm^{-1} in titaniferous pyroxenes to a Ti^{3+} – Ti^{4+} charge transfer transition. In support of their interpretation of the Allende fassaite spectra, Dowty and Clark (1973) cite diffuse reflectance spectral data for synthetic $\text{NaTi}^{3+}\text{Si}_2\text{O}_6$ (Prewitt *et al*, 1972). The spectrum illustrated by Prewitt *et al* (1972) shows a weak, broad band at 15,600 cm^{-1} and an intense absorption edge commencing around 500 nm (20,000 cm^{-1}) and obliterating the blue region of the spectrum. They interpreted the 15,600 cm^{-1} band as “. . . a charge transfer process between Ti^{3+} ions and some residual Ti^{4+} along the face-sharing¹

$M1$ octahedral chains.” No explanation was offered for the missing crystal-field features necessitated by the abundant Ti^{3+} ions in a stoichiometric $\text{NaTi}^{3+}\text{Si}_2\text{O}_6$ pyroxene, nor for how Ti^{4+} could be present in the $M1$ sites. To demonstrate that the 15,600 cm^{-1} band in the spectrum of synthetic $\text{NaTiSi}_2\text{O}_6$ is also a crystal-field transition in Ti^{3+} ions, we now cite structural data for several clinopyroxenes.

Selected interatomic distances for a variety of clinopyroxenes (Table 2) show for each structure a wide range of metal-oxygen and oxygen-oxygen distances in the coordination polyhedron about the $M1$ positions. This indicates considerable distortion of the $M1$ coordination sites from octahedral symmetry. Note that the larger distortion of the $M1$ site (based on ranges of $M1$ –O distances) in acmite compared to that in ureyite appears to be correlated with the larger ionic radius of Fe^{3+} (0.645 Å) relative to Cr^{3+} (0.615 Å) for these transition-metal-bearing sodic pyroxenes. Although crystal structural data for synthetic $\text{NaTi}^{3+}\text{Si}_2\text{O}_6$ are not yet available (Prewitt, personal communication), the larger ionic

¹ Presumably “edge-sharing $M1$ octahedral chains” is intended. Face sharing of octahedra does not occur in the pyroxene structure, in accord with Pauling’s third rule.

TABLE 2. Selected Bond Lengths in Certain Clinopyroxenes

| Bond and {multiplicity} | Diopside CaMgSi ₂ O ₆ (1) | Allende* Fassaite (2) | Jadeite NaAlSi ₂ O ₆ (3) | Ureyite NaCrSi ₂ O ₆ (4) | Acmite NaFeSi ₂ O ₆ (5) |
|----------------------------|---|-----------------------------|--|--|---|
| M1-01A1, B1 {2} | 2.115 | 2.120 | 1.996 | 2.039 | 2.109 |
| M1-01A2, B2 {2} | 2.065 | 2.064 | 1.933 | 2.009 | 2.029 |
| M1-02C1, D1 {2} | 2.050 | 1.993 | 1.856 | 1.947 | 1.936 |
| Mean M1-O | {6} 2.077 | 2.059 | 1.928 | 1.998 | 2.025 |
| Range | 0.065 | 0.127 | 0.140 | 0.092 | 0.173 |
| 01A1-01B1 | 2.781 | 2.78 | 2.726 | 2.775 | 2.798 |
| 02C1-02D1 | 2.981 | 2.98 | 2.790 | 2.897 | 2.941 |
| 01A1-02C1 {2} | 3.013 | 2.88 | 2.716 | 2.815 | 2.860 |
| 01A1-01A2 {2} | 3.051 | 3.10 | 2.918 | 2.975 | 2.985 |
| 01A2-02Ci {2} | 2.878 | 2.76 | 2.677 | 2.797 | 2.819 |
| 01A2-02D1 {2} | 2.979 | 2.91 | 2.818 | 2.904 | 2.964 |
| 01A1-01B2 {2} | 2.813 | 2.75 | 2.458 | 2.617 | 2.639 |
| Mean 0-0 | {12} 2.936 | 2.88 | 2.724 | 2.824 | 2.856 |
| Range | 0.270 | 0.35 | 0.460 | 0.358 | 0.346 |

* 0-0 distances calculated from data of Dowty and Clark (1973).

- (1) J.R. Clark, D.E. Appleman and J.J. Papike, *Min. Soc. Amer., Spec. Pap.*, 2, 31 (1969).
- (2) E. Dowty and J.R. Clark, *Amer. Min.*, 58 (1973).
- (3) C.T. Prewitt and C.W. Burnham, *Amer. Min.*, 51, 956 (1966); J.R. Clark *et al.*, loc. cit.
- (4) J.R. Clark *et al.*, loc. cit.
- (5) J.R. Clark *et al.*, loc. cit.

radius of Ti³⁺ (0.67) and the trends shown by the data in Table 2 suggest that considerable distortion of the M1 coordination site from octahedral symmetry is to be expected in NaTi³⁺Si₂O₆.

Therefore, the spectrum of NaTiSi₂O₆ should contain two absorption bands representing crystal field transitions in Ti³⁺, one at 15,600 cm⁻¹ and another at higher energy. Assuming the crystal field splitting parameter, Δ_o, to be comparable to the value 18,750 cm⁻¹ derived for Ti³⁺ ions in the Allende and Apollo 11 pyroxenes (Figure 4 and Table 1), the high energy band would be predicted to lie at about 22,000 cm⁻¹. In this location it would be obscured by the intense absorption edge beyond 500 nm observed by Prewitt *et al.* (1972).

The occurrence of Ti³⁺ ions in clinopyroxenes provides yet another example of the stabilizing influence of a distorted coordination site in a silicate structure on the crystal chemistry of certain transition metal ions (Burns, 1970, pp. 119-124). The importance of this site-distortion effect is demonstrated not only by the well documented fractionations of Fe²⁺ ions (which have an electronic configuration analogous to Ti³⁺, Figure 3) into the most distorted sites of silicate minerals, but also by the occurrence of unstable cations such as Mn³⁺ and Cr²⁺ in epidotes and lunar olivines, respectively. By analogy, the distorted M1 coordination site of clinopyroxenes not only enhances the crystal field stabil-

ization energy of Ti³⁺, but also leads to the fractionation of this relatively unstable cation into the clinopyroxene structure. Failure to take into account this relatively new principle of crystal chemistry inevitably leads to misinterpretation of experimental data, such as incorrect assignments of absorption bands in crystal-field spectra.

Note Added. Our discussion of the broad absorption band at 21,000 cm⁻¹ in the Allende titanian fassaite should not be confused with interpretations of the sharper peak at 22,000 cm⁻¹ observed in the absorption spectra of terrestrial titanogites. Current work on the Mössbauer and absorption spectra of several terrestrial titanogites suggests that the spectral feature found at 22,000 cm⁻¹ is due to absorption by Fe³⁺ ions in octahedral and tetrahedral coordination in the pyroxene structure (Burns, Abu-Eid, and Leung, 1972).

Acknowledgments

We thank Dr. Brian Mason for lending us a section of the Allende meteorite. Back-up spectral and universal stage measurements were performed by Mr. Rateb M. Abu-Eid and Professor Irene S. Leung, respectively. Mrs. Virginia Mee Burns helped in the bibliographic research, and Mrs. Sue Hall assisted in the preparation of the manuscript. We thank Professors E. Dowty and C. T. Prewitt for providing copies of manuscripts in advance of publication. Research on absorption spectral measurements of lunar and terrestrial minerals is supported by a grant from the National Aeronautics and Space Administration (NGR-22-009-551).

References

- BANCROFT, G. M., R. G. BURNS, AND A. G. MADDOCK (1967) Applications of the Mössbauer effect to silicate mineralogy. I. Iron silicates of known crystal structure. *Geochim. Cosmochim. Acta*, **31**, 2219–2246.
- BURNS, R. G. (1970) *Mineralogical Applications of Crystal Field Theory*. Cambridge University Press.
- , R. M. ABU-EID, AND F. E. HUGGINS (1972) Crystal field spectra of lunar pyroxenes. *Proc. Third Lunar Sci. Conf., Geochim. Cosmochim. Acta, Suppl. 3* (in press).
- , ———, AND IRENE S. LEUNG (1972) Titanaugites contain tetrahedral Fe^{3+} ions: A Mössbauer and electronic absorption study. *Geol. Soc. Amer. Abstr. Programs*, **4**, 463–464.
- CHRISTENSEN, A. N., AND R. G. HAZELL (1967) The crystal structure of $\text{NaIn}(\text{SiO}_3)_6$. *Acta Chem. Scand.* **21**, 1425–1429.
- CLARK, JOAN R., D. E. APPLEMAN, AND J. J. PAPIKE (1969) Crystal-chemical characterization of clinopyroxenes based on eight new structure refinements. *Mineral. Soc. Amer. Spec. Pap.* **2**, 31–50.
- CLARK, R. J. H. (1968) *The Chemistry of Titanium and Vanadium*. Elsevier Publishing Co.
- COTTON, F. A., AND M. D. MEYERS (1960) Magnetic and spectral properties of the spin-free $3d^8$ systems $\text{Fe}(\text{II})$ and $\text{Co}(\text{III})$ in cobalt (III) hexafluoride ion: probable observation of dynamic Jahn-Teller effects. *J. Amer. Chem. Soc.* **82**, 5023–5026.
- DOWTY, E., AND J. R. CLARK (1973) Crystal structure refinement and visible-region absorption spectra of a Ti^{3+} fassaite from the Allende meteorite. *Amer. Mineral.* **58**, 230–242.
- FREED, R. L., AND D. R. PEACOR (1967) Refinement of the crystal structure of johannsenite. *Amer. Mineral.* **52**, 709–720.
- HARTMANN, H., AND H. L. SCHLÄFER (1951) Über die Lichtabsorption komplexer Ionen des dreiwertigen Titans. *Z. Phys. Chem.* **197**, 116–142.
- , ———, AND K. H. HANSEN (1957) Über die Lichtabsorption von Dipolkomplexen des III-wertigen Titans vom Typ $[\text{TiA}_6]^{3+}$ mit $\text{A} = \text{H}_2\text{O}$, CH_3OH , $\text{C}_2\text{H}_5\text{OH}$ und $(\text{NH}_2)_2\text{CO}$. *Z. anorg. allgem. Chem.* **289**, 40–65.
- HOFF, G. R., AND C. H. BRUBAKER, JR. (1971) The preparation and characterization of some mixed-ligand complexes of titanium (III). *Inorg. Chem.* **10**, 2063–2066.
- PEACOR, D. R. (1967) Refinement of the crystal structure of a pyroxene of formula $M_1M_{11}(\text{Si}_{1.5}\text{Al}_{0.5})\text{O}_6$. *Amer. Mineral.* **52**, 31–41.
- PREWITT, C. T., AND C. W. BURNHAM (1966) The crystal structure of jadeite, $\text{NaAlSi}_2\text{O}_6$. *Amer. Mineral.* **51**, 956–975.
- PREWITT, C. T., R. D. SHANNON, AND W. B. WHITE (1972) Synthesis of a pyroxene containing trivalent titanium. *Contrib. Mineral. Petrology* **35**, 77–82.
- SCHLÄFER, H. L. (1964) Lichtabsorptionseigenschaften und Magnetische Momente von oktaedrischen komplexverbindungen des III-Wertigen Titans. In, B. Jezowska-Trzebiatowska, Ed. "Theory and Structure of Complex Compounds," *Proc. Symp. Theory Struct. Complex Comp., Wroclaw, 1962*, pp. 181–193.
- SCHROPFER, L. (1971) Über den gekoppelten Ersatz von $\text{Mg}^{2+}\text{Si}_2^{4+}$ durch $\text{Ti}^{4+}\text{Al}_2^{3+}$ im Diopsid. *Neues Jahrb. Mineral. Abh.* **116**, 20–40.
- WARREN, B. E., AND W. L. BRAGG (1928) The crystal structure of diopside, $\text{CaMg}(\text{SiO}_3)_2$. *Z. Kristallogr.* **69**, 168–193.