

## Ilmenite and Armalcolite in Apollo 17 Breccias

IAN M. STEELE

Department of the Geophysical Sciences,  
The University of Chicago, Chicago, Illinois 60637

### Abstract

Recrystallized Apollo 17 breccias contain high-Mg ilmenite, "Zr-", and "Cr-Zr-armalcolites"; textural relations indicate growth during recrystallization. The Mg/Fe ratios of both armalcolite types correlate positively with the Mg/Fe ratio of coexisting silicates; the distribution coefficient— $K_D = (\text{Mg/Fe})_{\text{Ti-oxide}}/(\text{Mg/Fe})_{\text{olivine}}$ —differs for the two armalcolite types ( $K_{\text{DZr-Arm}} \approx 0.29$ ;  $K_{\text{DCr-Zr-Arm}} \approx 0.12$ ), suggesting different structural control. Ilmenite from breccia is consistently high in MgO (range 4.6–7.6, average 6.5 wt percent MgO) compared to ilmenite from ilmenite basalt (range 0.8–3.5, average 1.8 wt percent MgO) reflecting the higher Mg/Fe ratio of the breccia;  $K_{\text{Dil}} = 0.125$  for recrystallized breccia. CaO in breccia ilmenite tends to be higher than in basaltic ilmenite.

Both armalcolite types coexist with high-Mg silicates [ $\text{Mg}/(\text{Mg} + \text{Fe}) > \sim 0.75$ ] while ilmenite occurs with silicates of lower Mg content. This relationship and the armalcolite → ilmenite reaction in ilmenite basalts suggest that armalcolite is a stable Ti-oxide phase coexisting with high-Mg silicates. Such an assemblage should be considered in lunar partial melting models especially because armalcolite can carry appreciable light rare-earth elements.


### Introduction

Ti-oxide phases are common in lunar rocks, and numerous studies have been made on their compositional variation (e.g., Haggerty, 1973a; El Goresy, Ramdohr, and Taylor, 1971; Brett *et al*, 1973; Taylor, McCallister, and Sardi, 1973). Few data exist regarding relationships between Ti-oxides and coexisting silicates. The most common observation is the positive correlation between the MgO content of ilmenite and the bulk rock Mg/Fe value (e.g., Powell and Weiblen, 1972; Drake *et al*, 1970). Anderson *et al* (1972) used the distribution of Mg between ilmenite and low-Ca pyroxene to estimate temperature of breccia equilibration, but calibration at low temperatures was poor. Akella and Boyd (1973) investigated experimentally the distribution of Ti and Al between coexisting silicates, oxides, and liquids. The basic problem in interpreting silicate–Ti-oxide data is whether equilibrium was attained, especially for igneous samples with zoning and incomplete reactions. From silicate compositions and textures of Apollo 14 samples, Warner (1972) showed that equilibrium is approached for fine-grained recrystallized breccia, a rock type common at the Apollo 16 and 17 sites (Cameron *et al*, 1973). Apollo 17 breccias were used in the present study


because both textures and silicate compositions indicate equilibrium, and because Ti-oxides are especially common with textures suggesting growth during recrystallization. Special attention was given to Ti-oxide compositions and their relation to coexisting silicate compositions.

### General Characteristics of Samples and Analytic Technique


The samples available for study comprised 1 to 2 mm fines 78462.2, 78442.2, 78482.5, 72502.19, 78422.2, and 73242.2. Thin sections were prepared and petrographic descriptions made for all fragments larger than ~ 0.5 mm. Fine-grained, holocrystalline breccia is common in all six, whether from the South Massif or Sculptured Hills stations. Calcium-rich plagioclase, low-Ca pyroxene, and high-Mg olivine dominate and minor opaque oxides are usually present. These breccias are always fine-grained (< 0.1 mm) although coarser clasts are sometimes present. Figures 1a–1d illustrate the textures of four fragments. Most typically, the Ti-oxides occur as grains much larger than the silicate grains which they enclose (Fig. 2a). This is interpreted as a result of late growth of the Ti-oxide phase. Less commonly, small Ti-oxide grains occur, but these



a



b




c

also tend to enclose silicate grains suggesting late growth (Fig. 2b).

Electron-microprobe analyses of Mg, Ca, and Fe were obtained for 10 to 25 silicate grains selected at random in each fragment. Silicon was determined by difference and Bence-Albee (1968) matrix corrections were applied. Nearly all fragments show very restricted ranges of silicate compositions dominated by low-Ca pyroxene but rarely by olivine (see Fig. 3). These restricted patterns indicate that equilibrium was attained during metamorphism (Warner, 1972), and enable assignment of a  $mg$  ( $= 100 \text{ Mg}/(\text{Mg} + \text{Fe})$ ) value to each fragment (Table 1). In addition several ilmenite basalt fragments are listed without  $mg$  values because the small sample size precluded obtaining accurate  $mg$  ranges; however, analyses of other ilmenite basalts indicate  $mg$  values less than  $\sim 70$ . The one plagioclase basalt in Table 1 has  $mg$  values ranging from  $\sim 95$  to 75, but these values probably do not represent olivines occurring with ilmenite in the mesostasis.

The fragments enumerated in Table 1 were chosen for detailed study because they had: (1) a narrow range of silicate compositions for any one fragment; (2) a wide range of silicate  $mg$  values for all fragments; (3) a texture suggesting late growth of Ti-oxides; and (4) reasonably large ( $>20 \mu\text{m}$ ) Ti-oxide grains suitable for microprobe analysis. In addition five ilmenite basalt fragments were selected at random as well as one plagioclase basalt so as to evalu-




d


FIG. 1. Photomicrographs illustrating textures of fragments studied. (a) 72502,19-3; (b) 73242,4-15; (c) 78422,2-15; (d) 78442,2-25. Scale: a,b,c—long dimension = 0.9 mm; d—0.45 mm. Transmitted plane-polarized light.

ate Ti-oxide compositions in relation to rock type.

Microprobe data were obtained for the Ti-oxide phases usually by analyzing two grains in each fragment. Standards were: Ti,Fe-A128 ilmenite; Ca,Si,Mg-diopside glass; Mn-Mn-hortonolite; Zr-baddeleyite; Al-anorthite glass; Cr-MB5 chromite. Backgrounds were collected for each mineral type, and all data were obtained without moving the sample with respect to the beam. Operating conditions were: 15 kV; 0.3  $\mu$ A beam current; focused beam ( $\sim 2 \mu$ m). Standard corrections were applied for deadtime, drift, and background. Matrix corrections



a



b

FIG. 2. Photomicrographs showing details of oxide-silicate intergrowth in two fragments illustrated in Figure 1. (a) 72502,19-3; (b) 78422,2-15. Scale: long dimension = 0.45 mm.




FIG. 3. Partial pyroxene quadrilateral showing ranges of pyroxene and olivine compositions for fragments illustrated in Figure 1. See text for details.

were taken from Albee and Ray (1970). Oxide sums for representative ilmenites (Table 2a) are quite acceptable ( $100 \pm 1.5$  wt percent), but armalcolite<sup>1</sup> sums (Table 2b) were low (94–99 percent). Because rare earths have been observed in “Cr-Zr-armalcolite,” wavelength scans were made. Lanthanum and the light rare earths (Ce,Pr,Nd, and Sm) were detected as were traces of Hf,Nb, and Y; however, all were below 0.1 oxide wt percent except  $\text{La}_2\text{O}_3$  ( $\cong 0.15$  wt percent) and  $\text{Ce}_2\text{O}_3$  ( $\cong 0.32$  wt percent). No other elements were detected. Low sums for “Cr-Zr-armalcolite” were also obtained by Steele and Smith (1972) using the same standards. Peckett, Phillips, and Brown (1972) noted the difficulty in obtaining proper oxide totals from other Zr-Ti oxides and suggested the presence of light elements, especially Li; however, Haggerty (1972, 1973a), Albee, Gancarz, and Chodos (1973) and Peckett, Phillips, and Brown (1972) did obtain good oxide totals for “Cr-Zr-armalcolite.” Several ilmenite analyses were rejected because of high oxide values for Si,Ca, and Al which probably came from silicate overlap. Rather high values for Si, Ca, and Al ( $>0.5$  oxide wt percent) recorded in the literature for ilmenites are questionable since overlap with a silicate is probable.

<sup>1</sup> The name armalcolite is used only to signify a similarity in composition; a structural similarity is not implied. Quotes are used for the specific terms “Zr-” and “Cr-Zr-armalcolite” to emphasize this point (Brown *et al.*, 1973).



TABLE 2b. Apollo 17 Cr-Zr- and Zr-Armacolite Analyses

	Cr-Zr-Armacolite			Zr-Armacolite						
	78442,2			78442,2		78422,2		78462,2		
	25a	25b	25c	3	12	16a	16b	7	11a	11b
MgO	2.79	2.65	3.05	9.17	9.28	11.4	12.1	7.25	7.80	7.44
Al <sub>2</sub> O <sub>3</sub>	1.46	1.05	.98	.59	.99	1.08	.90	.70	1.21	1.26
SiO <sub>2</sub>	.26	.11	.37	.11	.13	.11	.31	.20	.62	.29
CaO	3.30	3.19	3.18	.39	.46	.51	.45	.50	.72	.66
TiO <sub>2</sub>	61.4	62.8	62.2	71.6	70.9	71.0	72.8	69.6	66.6	68.0
Cr <sub>2</sub> O <sub>3</sub>	10.1	10.6	10.1	1.07	2.28	1.90	2.13	1.65	1.77	1.56
MnO	.08	.09	.08	.05	.03	.03	.04	.09	.05	.09
FeO	8.33	7.76	8.21	14.3	12.4	8.79	8.23	16.6	13.9	16.4
ZrO <sub>2</sub>	6.05	6.00	6.65	1.12	1.81	.52	.52	2.03	1.88	1.92
Σ	93.77	94.25	94.82	98.40	98.28	95.34	97.48	98.62	94.55	97.62
Cations based on 5(O)										
Mg	.17	.16	.18	.50	.51	.63	.65	.40	.45	.42
Al	.03	.03	.02	.01	.02	.02	.02	.01	.03	.03
Si	.02	.01	.02	.0	.01	.0	.01	.01	.02	.01
Ca	.15	.14	.14	.01	.02	.02	.02	.02	.03	.03
Ti	1.90	1.94	1.91	1.97	1.96	1.98	1.98	1.95	1.94	1.93
Cr	.17	.17	.16	.01	.03	.03	.03	.03	.03	.02
Mn	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0
Fe	.29	.27	.28	.44	.38	.27	.25	.52	.45	.52
Zr	.12	.12	.13	.02	.03	.01	.01	.03	.04	.04
<u>Mgx100</u> (Mg+Fe)	37	37	39	53	57	70	72	44	50	45

not confirm this result, and the present observation may result from the limited types of fragments studied. Alternatively, secondary fluorescence of Ca-rich phases may produce the high Ca values, but because most analyzed ilmenite grains were at least 0.1 mm in size and high Si values were not observed (although the secondary fluorescence effect would be less), the high-Ca values probably do not result from secondary fluorescence. The MgO-CaO plot (Fig. 4) shows distinct ranges for ilmenites from the three rock types; unfortunately ilmenites from plagioclase basalt are poorly represented. Brett *et al* (1973) did not find such distinct ranges for ilmenites from Luna 20 specimens; however, consistent differences were observed for ZrO<sub>2</sub> between ilmenites from "highland" rocks (low ZrO<sub>2</sub>) and mare basalts (high ZrO<sub>2</sub>). The present data for ZrO<sub>2</sub> are consistently low (<0.11 wt percent) and usually close to the detection limit (0.02). Values for ilmenites from Apollo 17 ilmenite basalts tend to be slightly higher than for breccia. Only the two ilmenites from the mesostasis of the plagioclase basalt show distinctly high values, but this is to be

expected because of Zr enrichment in late liquids. Prinz *et al* (1973) observed a weak correlation between MgO and Cr<sub>2</sub>O<sub>3</sub> in ilmenites but no correlation was found here.

Of the minor elements in ilmenite, Mg seems to best reflect the crystallization conditions. The




FIG. 4. CaO vs MgO in ilmenites from recrystallized breccia, ilmenite basalts, and plagioclase basalt.

Mg/Fe distribution between oxide and the silicate may depend mostly on temperature, and be almost independent of bulk composition. Anderson *et al* (1972) attempted to evaluate the temperature dependence of this distribution for coexisting pyroxene and ilmenite. For high temperatures ( $>900^{\circ}\text{C}$ ),  $K_D$  increases  $\sim 0.04$  for a  $100^{\circ}\text{C}$  rise in temperature, and the Mg/Fe fractionation between olivine and low-Ca pyroxene is close to unity (Ramberg and DeVore, 1951). The ilmenite-olivine data for recrystallized breccias (Fig. 5) form a tight grouping with  $K_D$  near 0.125: this corresponds to an estimated temperature near  $950^{\circ}\text{C}$ . Different  $K_D$  values for other lunar breccias should indicate different temperatures of equilibration relative to the Apollo

17 breccia. Presumably one could estimate the Mg/Fe value of olivine in equilibrium with ilmenite in an ilmenite basalt; this in turn would imply the Mg/Fe value of the bulk liquid at the time of crystallization, assuming equilibrium crystallization of ilmenite and olivine.

The combination of different bulk-rock  $mg$  values and equilibration temperatures could account for the distinct ranges of MgO (Brett *et al*, 1973; Fig. 4, this work) for ilmenites from mare-type basalts (low MgO in ilmenite, low bulk  $mg$ ) and plagioclase-rich rock types (high MgO in ilmenite, high bulk  $mg$ ). The generally higher temperature of crystallization for the basalt relative to the metamorphic temperatures of the plagioclase-rich breccia




FIG. 5. Plot of  $\text{Mg}/(\text{Mg} + \text{Fe})$  in armalcolite types vs  $\text{Mg}/(\text{Mg} + \text{Fe})$  in coexisting olivine. Points labeled Apollo 14 are from Steele and Smith (1972). Point labeled 14310 is from Peckett, Phillips, and Brown (1972); the  $mg$  value for a hypothetical olivine was inferred from the most Fe-rich of the low-Ca pyroxenes in 14310, and a large error should be expected. Points labeled Albee are from Albee, Gancarz, and Chodos (1973); point labeled Tarasov is from Tarasov *et al* (1973); point labeled Brown is from Brown *et al* (1973)—uncertainty in position is because of basaltic nature of this fragment. Ferropseudobrookite points were based on  $\text{Mg}/(\text{Mg} + \text{Fe}^{2+})$  values; one sample coexists with two olivines, hence the indicated range. Apollo 11 (A-11) armalcolite range was collected from many sources, but most values lie in cross-hatched rectangle. Horizontal lines from this range indicate range of olivine which might coexist with these armalcolites.

would tend to increase the MgO content of the basaltic ilmenite (higher  $K_D$ ), but this effect would be small (see before).

### Armalcolite

Two armalcolite types, "Zr-" and "Cr-Zr-armalcolite," were found in the recrystallized breccia; both types of armalcolite were recognized previously by Haggerty (1972) and the occurrence of "Cr-Zr-armalcolite" was discussed by Levy *et al* (1972), Peckett, Phillips, and Brown (1972), Steele and Smith (1972), and Haggerty (1973a).

The two armalcolite types coexist with silicates which have a rather wide range of  $mg$  values. Armalcolite compositions from Table 2b are plotted on Figure 5 as  $Mg/(Mg + Fe)$  in armalcolite vs  $Mg/(Mg + Fe)$  in coexisting olivine; some data were obtained from the literature (see caption for details). Figure 5 demonstrates the positive correlation between the  $mg$  value of olivine and the  $mg$  of both armalcolite types. Several interesting points can be made. (1) "Cr-Zr-armalcolites" and "Zr-armalcolites" have different distribution coefficients with respect to olivine  $K_{D_{Zr-arm}} \approx 0.29$ ,  $K_{D_{Cr-Zr-arm}} \approx 0.12$ . This may arise simply from the influence of appreciable  $Cr_2O_3$  and  $ZrO_2$  in the "Cr-Zr-armalcolite"; however, optical data (Levy *et al*, 1972) suggest that "Cr-Zr-armalcolite" is cubic rather than orthorhombic, and a real structural difference might well cause the difference in distribution coefficients. (2) Limited data for terrestrial ferro-pseudobrookite phenocrysts coexisting with olivine phenocrysts from Kilauean lava flows (Anderson and Wright, 1972) fall near the "Zr-armalcolite" trend on Figure 5 and thus suggest a similarity in stoichiometry as well as structure (Lind and Housley, 1971). (3) The Apollo 11 type armalcolites (see range at right of Figure 5), which are similar to "Zr-armalcolites" except mainly for the lower  $ZrO_2$  content, should be in equilibrium with olivines with an  $mg$  value of 0.74 to 0.78. This range corresponds to the most Mg-rich olivines found in ilmenite basalts, consistent with joint crystallization. Anderson (1971) argued for an exotic origin of some Apollo 11 armalcolite, but he gave no evidence from mineral compositions to support or refute this proposal. Based on the present data, Apollo 11 armalcolite could have crystallized from the Apollo 11 ilmenite basalt. (4) Anderson (1973) reported armalcolite in association with a Mg-rich assemblage in a peridotite 15445.10, but direct evidence of a

genetic association was not present. The  $mg$  value of this armalcolite ( $ZrO_2 < 0.1$  wt percent) suggests that it is in equilibrium with olivine of  $mg = 89$  (Fig. 5). Within experimental error this agrees with the value of  $mg = 91$  for the olivine of 15445. Hence the armalcolite could be genetically associated with the Mg-rich assemblage and need not be exotic. (5) Armalcolite and olivine from a terrestrial ultramafic nodule (Cameron and Cameron, 1973) plot on the "Zr-armalcolite" trend of Figure 5.

The armalcolite types described here and in the literature always seem to coexist with Mg-rich silicates.<sup>2</sup> When olivine has  $mg > \sim 75$ , armalcolite appears to be stable rather than ilmenite. This relationship was also seen in the armalcolites from mare-type basalts; initial crystallization yielded Mg-rich olivine ( $\sim Fo$  75-80) and armalcolite, but with further crystallization armalcolite reacted to give ilmenite. With olivine of  $\sim Fo$  75, both ilmenite and armalcolite coexist (*e.g.*, 78462.2-7; also see Albee, Gancarz, and Chodos, 1973). The Ti-oxide reactions as described by Haggerty (1973a) probably occur in assemblages with  $mg$  near 75; otherwise either ilmenite or armalcolite would be stable.

### General Discussion

The association of Mg-rich silicates and armalcolite has important implications regarding the chemistry of the Moon. If the mantle of the Moon formed during an initial differentiation and thus consists of Mg-rich silicates, one or more armalcolites would be expected as the coexisting Ti-oxide(s). The occurrence of an early crystallizing Ti-oxide phase may have effectively concentrated titanium in the mantle, and later partial melts derived from this region (*e.g.*, Fe-Ti-basalts?) could be expected to be Ti-rich. The upper limit of stability for armalcolite in ilmenite basalts is about 3 kbar (Ringwood and Essene, 1970), but this limit may not apply to Mg-rich assemblages. In addition, if "Cr-Zr-armalcolite" has a different crystal structure (see before), the phase relations might be quite different.

The REE-bearing "Cr-Zr-armalcolite" probably is not a primary liquidus phase because of its enrichment in trace elements; it may result only from

<sup>2</sup> The 14310 point on Figure 5 appears to be an exception, but this rock is probably hybrid, formed from melting of heterogeneous regolith (Morgan *et al*, 1972).

metamorphism of polymict breccia incorporating REE-rich material (KREEP component?). In any case, the partial-melt products, particularly the REE distribution, of a "Cr-Zr-armalcolite"-bearing rock would be dependent on the melting relations of the armalcolite. Nava and Philpotts (1973) have suggested that KREEP liquids result from partial melting of a feldspathic source in the lunar crust; the REE content of such a partial melt could possibly result from the extent to which armalcolite contributes to this liquid.

Armalcolite appears, at least in one case (excluding ilmenite basalts), to be a primary igneous phase; Anderson (1973) interpreted 15445 as an igneous assemblage. Possibly igneous "Zr-armalcolite" was more common, but reactions such as those suggested by Haggerty (1972, 1973a) have erased most evidence.

A detailed experimental examination of armalcolite phase relations probably would do much to establish the role of the various armalcolite types in the Moon.

### Acknowledgments

Helpful discussions with A. T. Anderson, J. V. Smith, A. J. Irving, and C. E. Nehru as well as reviews by L. A. Taylor and A. El Goresy provided many useful suggestions for the improvement of this manuscript. Financial support was provided by NASA grant NGL-14-001-171.

### References

- AKELLA, J., AND F. R. BOYD (1973) Partitioning of Ti and Al between coexisting silicates, oxides, and liquids. *Proc. Fourth Lunar Sci. Conf., Geochim. Cosmochim. Acta, Suppl. 1*, 1, 1049-1059. MIT Press.
- ALBEE, A. L. AND L. RAY (1970) Correction factors for electron probe microanalysis of silicates, oxides, carbonates, phosphates, and sulfates. *Anal. Chem.* **42**, 1408-1414.
- , A. J. GANCARZ, AND A. A. CHODOS (1973) Metamorphism of Apollo 16 and 17 and Luna 20 metaclastic rocks at about 3.95 AE: samples 61156, 64423, 14-2, 65015, 67483, 15-2, 76055, 22006, and 22007. *Proc. Fourth Lunar Sci. Conf., Geochim. Cosmochim. Acta, Suppl. 4*, 1, 569-595. MIT Press.
- ANDERSON, A. T. (1971) Exotic armalcolite and the origin of Apollo 11 ilmenite basalts. *Geochim. Cosmochim. Acta*, **35**, 969-973.
- (1973) The texture and mineralogy of lunar peridotite, 15445.10. *J. Geol.* **81**, 219-226.
- , AND T. L. WRIGHT (1972) Phenocrysts and glass inclusions and their bearing on oxidation and mixing of basaltic magmas, Kilauea volcano, Hawaii. *Am. Mineral.* **57**, 188-216.
- , T. F. BRAZIUNAS, J. JACOBY, AND J. V. SMITH (1972) Thermal and mechanical history of breccias 14306, 14063, 14270, and 14321. *Proc. Third Lunar Sci. Conf., Geochim. Cosmochim. Acta, Suppl. 3*, 1, 819-835. MIT Press.
- BENCE, A. E., AND A. L. ALBEE (1968) Empirical correction factors for the electron microanalysis of silicates and oxides. *J. Geol.* **76**, 382-403.
- BRETT, R., R. C. GOOLEY, E. DOWTY, M. PRINZ, AND K. KEIL (1973) Oxide minerals in lithic fragments from Luna 20 fines. *Geochim. Cosmochim. Acta*, **37**, 761-773.
- BROWN, G. M., A. PECKETT, R. PHILLIPS, AND C. H. EMELEUS (1973) Mineral-chemical variations in the Apollo 16 magnesian-feldspathic highland rocks. *Proc. Fourth Lunar Sci. Conf., Geochim. Cosmochim. Acta, Suppl. 4*, 1, 505-518. MIT Press.
- CAMERON, K. L., J. W. DELANO, A. E. BENCE, AND J. J. PAPIKE (1973) Petrology of the 2-4 mm soil fraction from the Hadley-Apennine region of the Moon. *Earth Planet. Sci. Lett.* **19**, 9-21.
- , AND M. CAMERON (1973) Mineralogy of ultramafic nodules from Knippa Quarry, near Uvalde, Texas (abstr.). *Geol. Soc. Am. Abstr. Programs*, **5**, 566.
- DRAKE, M. J., I. S. MCCALLUM, G. A. MCKAY, AND D. F. WEILL (1970) Mineralogy and petrology of Apollo 12 sample No. 12013: a progress report. *Earth Planet. Sci. Lett.* **9**, 103-123.
- EL GORESY, A., P. RAMDOHR, AND L. A. TAYLOR (1971) The geochemistry of the opaque minerals in Apollo 14 crystalline rocks. *Earth Planet. Sci. Lett.* **13**, 121-129.
- HAGGERTY, S. E. (1972) The mineral chemistry of some decomposition and reaction assemblages associated with Cr-Zr, Ca-Zr, and Fe-Mg-Zr titanates. In *The Apollo 15 Lunar Samples*, Eds. J. W. Chamberlain and C. Watkins, The Lunar Science Institute, p. 88-91.
- (1973a) Armalcolite and genetically associated opaque minerals in the lunar samples. *Proc. Fourth Lunar Sci. Conf., Geochim. Cosmochim. Acta, Suppl. 4*, 1, 777-797. MIT Press.
- (1973b) Luna 20: mineral chemistry of spinel, pleonaste, chromite, ulvöspinel, ilmenite and rutile. *Geochim. Cosmochim. Acta*, **37**, 857-867.
- LEVY, C., M. CHRISTOPHE-MICHEL-LEVY, P. PICOT, AND R. CAYE (1972) A new titanium and zirconium oxide from the Apollo 14 samples. *Proc. Third Lunar Sci. Conf., Geochim. Cosmochim. Acta, Suppl. 3*, 1, 1115-1120. MIT Press.
- LIND, M. D., AND R. M. HOUSLEY (1971) Crystallization studies of lunar igneous rocks: crystal structure of synthetic armalcolite. *Science*, **175**, 521-523.
- MORGAN, J. W., J. C. LAUL, U. KRÄHENBÜHL, R. GANAPATHY, AND E. ANDERS (1972) Major impacts on the Moon: characterization from trace elements in Apollo 12 and 14 samples. *Proc. Third Lunar Sci. Conf., Geochim. Cosmochim. Acta, Suppl. 3*, 2, 1377-1395. MIT Press.
- NAVA, D. F., AND J. A. PHILPOTTS (1973) A lunar differentiation model in light of new chemical data on Luna 20 and Apollo 16 soils. *Geochim. Cosmochim. Acta*, **37**, 963-973.
- PECKETT, A., R. PHILLIPS, AND G. M. BROWN (1972) New zirconium-rich minerals from Apollo 14 and 15 lunar rocks. *Nature*, **236**, 215-217.
- POWELL, B. N., AND P. W. WEIBLEN (1972) Petrology and



- origin of lithic fragments in the Apollo 14 regolith. *Proc. Third Lunar Sci. Conf., Geochim. Cosmochim. Acta, Suppl. 3, 1*, 837-852. MIT Press.
- PRINZ, M., E. DOWTY, K. KEIL, AND T. E. BUNCH (1973) Mineralogy, petrology and chemistry of lithic fragments from Luna 20 fines: origin of the cumulate ANT suite and its relationship to high-alumina and mare basalts. *Geochim. Cosmochim. Acta*, **37**, 979-1006.
- RAMBERG, H., AND G. DEVORE (1951) The distribution of Fe<sup>++</sup> and Mg<sup>++</sup> in coexisting olivines and pyroxenes. *J. Geol.* **59**, 193-210.
- RINGWOOD, A. E., AND E. ESSENE (1970) Petrogenesis of Apollo 11 basalts, internal constitution and origin of the Moon. *Proc. Apollo 11 Lunar Sci. Conf., Geochim. Cosmochim. Acta Suppl. 1*, 769-799. Pergamon.
- STEELE, I. M., AND J. V. SMITH (1972) Occurrence of diopside and Cr-Zr-armalcolite on the Moon. *Nature Phys. Sci.*, **237**, 105-106.
- TARASOV, L. S., M. A. NASAROV, I. D. SHEVALEEVSKY, E. S. MAKAROV, AND V. I. IVANOV (1973) Mineralogy of anorthositic rocks from the region of the crater Apollonius C (Luna 20). *Proc. Fourth Lunar Sci. Conf., Geochim. Cosmochim. Acta, Suppl. 4, 1*, 333-349. MIT Press.
- TAYLOR, L. A., R. H. MCCALLISTER, AND O. SARDI (1973) Cooling histories of lunar rocks based on opaque mineral geothermometers. *Proc. Fourth Lunar Sci. Conf., Geochim. Cosmochim. Acta, Suppl. 4, 1*, 819-828. MIT Press.
- WARNER, J. L. (1972) Metamorphism of Apollo 14 breccias. *Proc. Third Lunar Sci. Conf., Geochim. Cosmochim. Acta, Suppl. 3, 1*, 623-643. MIT Press.

*Manuscript received, January 10, 1974; accepted for publication, April 4, 1974.*