

Experimental evidence at high pressure for potassic metasomatism in the mantle of the Earth¹

I. D. RYABCHIKOV

*Institute of Geology of Ore Deposits
35 Staromonetny
Moscow, 109017, U.S.S.R.*

AND A. L. BOETTCHER

*Institute of Geophysics and Planetary Physics
and Department of Earth and Space Sciences
University of California, Los Angeles
Los Angeles, California 90024*

Abstract

To assess the concentration of potassium in aqueous fluids as a model for metasomatism of the mantle of the Earth, we experimentally determined the solubility of potassium in an aqueous vapor in equilibrium with phlogopite and other phases to pressures of 30 kbar at 1100°C. The concentration of potassium (in grams of K₂O per 100 g H₂O) is 4 at 11 kbar, 7 at 20 kbar, and 25 at 30 kbar. Such values are sufficient to transfer from the interior of the Earth all of the K₂O to the continental crust during formation of the hydrosphere by outgassing. However, it is likely that most of this transport was by silicate-rich liquids.

Introduction

A number of investigators have suggested that primary alkaline magmas and particularly highly potassic melts are formed by partial melting of mantle material that was previously enriched in potassium and other components by precursory metasomatic processes (Sobolev, 1977; Ryabchikov and Green, 1979; Lloyd and Bailey, 1975; Boettcher and O'Neil, 1980). The present work was designed to provide a quantitative check on the possibility of such processes by experimentally measuring K₂O concentrations in aqueous vapors in equilibrium with phlogopite-bearing mineral assemblages typical of the mantle at high pressures and temperatures.

Earlier studies (Wendlandt, 1977; Yoder and Kushiro, 1969) revealed that subsolidus experiments at 20 kbar for the composition joins phlogopite-H₂O and phlogopite-H₂O-CO₂ quenched to an assemblage containing phlogopite plus forsterite. This demonstrates that incongruent solubility of phlo-

pite in an aqueous vapor under these conditions is significant and that K₂O and Al₂O₃ dissolve in the vapor in excess of that in phlogopite composition. In this work we extended the investigation of phlogopite-containing joins to higher H₂O concentrations and determined the H₂O contents at various pressures where phlogopite became unstable, dissolving entirely in the vapor, enabling us to determine K₂O contents of fluids in equilibrium with phlogopite, olivine, and some other phases typical of mantle assemblages.

Experimental procedures

Starting materials

A mixture of oxides of anhydrous phlogopite composition was prepared by sintering and partial melting at about 1100°C. Fired gels of pyrope and forsterite compositions were added to anhydrous phlogopite powder for some of our experiments.

Experiments were performed in sealed platinum capsules on mixtures of known proportions of silicate components and water.

¹ Institute of Geophysics and Planetary Physics Contribution No. 1916.

Apparatus

All the experiments were in piston-cylinder apparatus similar to that described by Boyd and England (1960). The experimental method was described earlier (Allen and Boettcher, 1978).

Identification

After sealing, the capsules were folded in such a way that the starting mixtures were contained in one half of each of the capsules. During the run, however, the second half of each capsule was filled with vapor phase, which precipitated quench material during fast isobaric cooling at the end of an experiment (see Fig. 1). Comparison of the material from both halves of the capsules was very useful for the distinction of primary and quench phlogopite crystals.

The nature of quench material changed considerably with pressure. Between 10 and 14 kbar, it was mainly an amorphous, glass-like phase forming spheres, rods, and coatings on primary crystals (Fig. 2). At these pressures, quench phlogopite occurred in minor amounts as small ($<5 \mu\text{m}$), thin flakes.

At 20 kbar the proportion of quench phlogopite was noticeably greater, forming large ($>20 \mu\text{m}$), thin




Fig. 1. Photomicrographs of products of run 33; (a) quench vapor in folded part of capsule, (b) forsterite + quench vapor in main part of capsule. Bars are $100 \mu\text{m}$.




Fig. 2. Photomicrograph of quench vapor in products of run 26. Bar is $100 \mu\text{m}$.

plates with uneven extinction and some skeletal morphology. The spheres of amorphous material were typically situated along the growth figures of quench phlogopite. At 30 kbar, large ($700 \mu\text{m}$), euhedral (but thin) phlogopite plates were part of the quench material and were overgrown by irregular aggregates of phlogopite and amorphous material (Fig. 2). Primary phlogopites at all pressures were euhedral, relatively thick crystals (Fig. 3).

Olivines occurred in all our run products. They were typically euhedral, tabular grains whose average size increased with increasing pressure, ranging from $10 \mu\text{m}$ at 10 kbar to $500 \times 150 \mu\text{m}$ at 30 kbar. Some olivines contained fluid inclusions.

The run products for the join phlogopite-pyroxene-water contained orthopyroxene and spinel in addition to phlogopite and olivine.

Experimental results

The results of our experiments are summarized in Table 1. Their interpretation is based on the following reasoning. In the runs in which the products did not include primary phlogopite, all the K_2O was incorporated into the fluid phase, inasmuch as the co-




Fig. 3. Photomicrograph of thin plates of quench phlogopite and other quench products in run 24. Bar is $200 \mu\text{m}$.

Table 1. Experimental data

Run	Pres (kbars)	Temp (°C)	Bulk composition	$\frac{g \text{ K}_2\text{O}}{100 \text{ g H}_2\text{O}}$	Time (hrs)	Run products
32	10	1100	Aph ₁₇ Fo ₃₅ W ₄₈	4.2	9	Fo+Ph+Q
33	12	1100	Aph ₁₈ Fo ₃₆ W ₄₆	4.7	11.5	Fo+Q
30	14	1100	Aph ₁₇ Fo ₃₅ W ₄₈	4.3	9	Fo+Q
23	20	1100	Aph ₃₃ W ₆₇	5.8	13	Fo+Q
26	20	1100	Aph ₃₇ W ₆₃	7.0	9	Fo+(Ph)+Q
3	20	1100	Aph ₄₂ W ₅₈	8.7	8	Fo+Ph+Q
14	20	1100	Aph ₄₂ W ₅₈	9.6	10	Fo+Ph+Q
11	20	1100	Aph ₇₆ W ₂₄	36.6	8	Fo+Ph+Q
22	20	1050	Aph ₁₀ Py ₄₀ W ₅₀	2.2	9	Opx+Sp+Q
25	20	1050	Aph ₂₀ Py ₃₀ W ₅₀	4.8	11	Fo+Opx+Sp+Q
29	20	1050	Aph ₂₄ Py ₂₄ W ₅₂	5.6	12	Fo+Opx+Sp+Q
27	20	1050	Aph ₃₀ Py ₂₀ W ₅₀	7.2	12	Fo+(Opx)+Sp+Ph+Q
21	30	1100	Aph ₁₇ W ₈₃	10.3	12	Fo+Q
19	30	1100	Aph ₅₄ W ₄₆	13.8	8	Fo+Q
24	30	1100	Aph ₅₇ W ₄₃	15.8	8.5	Fo+Q
31	30	1100	Aph ₅₅ W ₄₅	21.9	10	Fo+Q
37	30	1100	Aph ₇₁ W ₂₉	28	8.5	Fo+Ph+Q

Abbreviations: Aph is anhydrous phlogopite composition ($\text{KMg}_3\text{AlSi}_3\text{O}_{11}$); Fo is forsterite (Mg_2SiO_4); W is water (H_2O); Py is pyrope composition ($\text{Mg}_3\text{Al}_2\text{Si}_2\text{O}_{12}$); Ph is phlogopite; Opx is orthopyroxene; Sp is spinel; Q is quench material (amorphous substance and quench mica); parentheses indicate minor amount.

existing crystalline phases (olivine or olivine + orthopyroxene + spinel) contain only negligible amounts of this element. Interpolating between the compositions of two adjacent runs with and without primary phlogopite, we deduce the $\text{K}_2\text{O}/\text{H}_2\text{O}$ ratio in the fluid coexisting with the traces of phlogopite in addition to forsterite or forsterite + orthopyroxene + spinel. In this manner, we estimated the values of K_2O concentrations (in grams of K_2O per 100 g of H_2O) in equilibrium with assemblages containing phlogopite and forsterite in 1100°C to be as follows: 4 at 11 kbar, 7 at 20 kbar, and 25 at 30 kbar.

The run products for the join phlogopite–pyrope–water at 20 kbar and 1100°C were buffered with respect to the chemical potential of SiO_2 (μ_{SiO_2}) by the presence of both forsterite and orthopyroxene and with respect to $\mu_{\text{Al}_2\text{O}_3}$ by olivine, orthopyroxene, and spinel. These compositions are more nearly representative of the chemistry of mantle material than are the joins phlogopite–water and phlogopite–forsterite–water. The concentration of K_2O in the vapor in equilibrium with phlogopite, forsterite, orthopyroxene, and spinel at 1050°C and 20 kbar was estimated from these experiments as 6 g K_2O per 100 g H_2O , only slightly lower than that for phlogopite + forsterite phase assemblages at the same pressure but at 1100°C.

Thus our experimental data permit us to evaluate $\text{K}_2\text{O}/\text{H}_2\text{O}$ ratios in aqueous fluids. The same reason-

ing is also applicable to Al_2O_3 concentrations, inasmuch as this component also occurs in olivines only in trace amounts. It is also clear that the $\text{K}_2\text{O}/\text{Al}_2\text{O}_3$ ratios of our solutions were always close to unity (the same as in phlogopite). Unfortunately, it is not possible to estimate MgO and SiO_2 concentrations in aqueous solutions from our experimental data alone. However, from the results of Modreski and Boettcher (1973), who analyzed quenched vapor in equilibrium with forsterite, enstatite, and phlogopite at 1050°C and 10 kbar, we conclude that the MgO content of aqueous fluid at these conditions is negligible. Therefore, it is rather safe to assume that concentrations of magnesium in the fluid in our runs at 1100°C and 11 kbar are also very low, because the pressure–temperature conditions of these runs and those of Modreski and Boettcher (1973) are similar. From this we estimate the K_2O (3.4 wt%), Al_2O_3 (3.7 wt%), and SiO_2 (6.6 wt%) concentrations in the aqueous phase in equilibrium with forsterite and phlogopite at 1100°C and 11 kbar, with the total content of dissolved oxides being approximately 14 wt%. At higher pressures MgO concentrations in fluids coexisting with phlogopite-containing phase assemblages must be substantial, judging from the increasing amount of quench phlogopite.

Eggler and Rosenhauer (1978) determined the solubility of diopside in an aqueous fluid at 20 and 30 kbar and approximately 1200°C to be approximately 10 wt%, which corresponds to about 2.5 wt% of MgO in solution. We assume that in our experiments at the same pressures the MgO content of fluid was the same or higher because the chemical potential of MgO for a forsterite-containing phase assemblage must be higher than in the presence of diopside alone. Assuming 2.5 wt% as a minimum value, we estimate the concentrations of other components, and the results of these calculations show that the total





Fig. 4. Photomicrograph of primary phlogopite and forsterite in the products of run 32. Bar is 100 μm .

contents of dissolved silicate components in aqueous fluids in equilibrium with phlogopite and forsterite at 1100°C and pressures of 20 and 30 kbar are about 25 wt% and 50 wt%, respectively. It is clear that at 30 kbar fluids in equilibrium with phlogopite-containing phase assemblages are near critical conditions.

Table 2 illustrates that the proportion of solute in the vapor is a function of pressure, ranging from about 14 wt% at 11 kbar to about 55 wt% at 30 kbar. At 30 kbar, our vapors contain approximately 25 wt% dissolved solids. This is about twice the value for the solubility of diopside in a vapor containing 95 wt% H₂O and 5 wt% CO₂, determined by Egger (1975). Experiments by Nakamura and Kushiro (1974) at 15 kbar reveal that aqueous vapor coexisting with forsterite and enstatite dissolves about 18 wt% SiO₂ at 1280°C and 22 wt% SiO₂ at 1310°C; the vapor coexisting with only enstatite contains up to 40 wt% SiO₂ at 1280°C and 15 kbar.

Petrological applications

Our experimental data were obtained for compositions with H₂O as the only volatile component. However, vapors in the mantle contain other volatile species, and experiments have shown that addition of components such as CO₂ and H₂ that lower the fugacity of H₂O will reduce the solubility of silicates in an aqueous vapor (for example, Shettel, 1973; Nakamura, 1974). Comparison of *P-T* coordinates for continental shield geotherms with the positions of vapor-phase isopleths for divariant assemblages corresponding to the following reactions (in the presence of H₂O and CO₂) shows that under these conditions H₂O/CO₂ values in vapors are high, particularly at depths below those of the stability of amphibole:



(Newton and Sharp, 1975; Egger, 1977; Wyllie, 1979)

Our results show that under these conditions fluids may be very efficient agents for transporting K₂O. For example, temperatures of about 1050°C are reached under continental shields at depths of about 130 km corresponding to pressures of 40 kbar, from the geotherm of Clark and Ringwood (1964). Extrapolation of our results to 1050°C and 40 kbar clearly shows that the vapors in equilibrium with phlogopite would contain in excess of 30 g K₂O per 100 g of H₂O. If we accept 10 g K₂O/100 g H₂O as a conservative estimate, and if we assume that all the water in the present hydrosphere was transferred from

the deep levels of the mantle of the Earth, then vapor could have transported $1.4 \cdot 10^{23}$ g of K₂O, which is approximately the amount in the continental crust. However, the shield geotherm intersects the peridotite-H₂O solidus at about 100 km, and it is more likely that much of the K₂O was transported in a silicate liquid.

It may be noted that the high mobility of K₂O-rich fluids under conditions of continental shield geotherms is consistent with the fact that only on continents do we find evidence for enrichment of mantle-derived magmatic rocks that commonly bring to the surface xenoliths of phlogopite-bearing ultramafic rocks. Under the higher temperature conditions of the deep oceanic lithosphere, vapor may have a higher value of CO₂/H₂O if carbonates are unstable (Egger, 1978). Mysen *et al.* (1978) have experimentally investigated the mobility of H₂O in peridotite under mantle conditions. These preliminary results do suggest that the rate of migration of H₂O-rich fluids is sufficiently rapid to permit mobilization of K₂O and other components. However, high mobility of K₂O in aqueous fluids is necessary but not sufficient for intensive mantle metasomatism. There must also exist certain mechanisms for fixation of dissolved K₂O. One obvious method to precipitate K₂O is to decrease the pressure and temperature of the ascending vapor. The change in the CO₂/H₂O ratio of the vapor during ascent (Wyllie, 1979) would also affect the solubility of K₂O and other solutes. Crystallization of amphiboles at depths of about 60 km would certainly increase CO₂/H₂O, and the K₂O, Al₂O₃, and other components previously dissolved in the fluid would take part in metasomatic reactions.

Some K₂O is fixed in the upper levels of the mantle at depths of about 50 km, as evidenced by metasomatism of lherzolite xenoliths in alkali basalts (Boettcher and O'Neil, 1980; Stewart and Boettcher, 1977; Wilshire *et al.*, 1980). However, the abundance of phlogopite in kimberlites and in xenoliths in kimberlites suggests that considerable amounts of K₂O reside at depths greater than 50 km.

Acknowledgments

This research was supported by NSF grant EAR78-16413 to ALB. Chris West and Dave Klimberg provided valuable technical assistance in the laboratory; Harriett Arnoff typed the manuscript. The reviews by D. H. Egger and B. O. Mysen are appreciated.

References

- Allen, J. C. and A. L. Boettcher (1978) Amphiboles in andesites and basalt: II. Stability as a function of $P-T-f_{\text{H}_2\text{O}}-f_{\text{O}_2}$. *Am. Mineral.*, 63, 1074-1087.

- Boettcher, A. L. and J. R. O'Neil (1980) Stable isotope, chemical, and petrographic studies of high-pressure amphiboles and micas: evidence for metasomatism in the mantle source regions of alkali basalts and kimberlites. *Am. J. Sci.*, in press.
- Boyd, F. R. and J. L. England (1960) Apparatus for phase-equilibrium measurements at pressures up to 50 kilobars and temperatures up to 1750°C. *J. Geophys. Res.*, 65, 741-748.
- Clark, S. P. and A. E. Ringwood (1964) Density distribution and constitution of the mantle. *Rev. Geophys.*, 2, 35-88.
- Eggler, D. H. (1975) CO₂ as a volatile component of the mantle: the system Mg₂SiO₄-SiO₂-H₂O-CO₂. In L. H. Ahrens *et al.*, Eds., *Physics and Chemistry of the Earth*, 9, p. 869-881. Pergamon Press, New York.
- (1977) The principle of the zone of invariant vapor composition: an example in the system CaO-MgO-SiO₂-CO₂-H₂O and implications for the mantle-solidus. *Carnegie Inst. Wash. Year Book*, 76, 428-435.
- (1978) Stability of dolomite in a hydrous mantle, with implications for the mantle solidus. *Geology*, 6, 397-400.
- and M. Rosenhauer (1978) Carbon dioxide in silicate melts: II. Solubilities of CO₂ and H₂O in CaMgSi₂O₆ (diopside) liquids and vapors at pressures to 40 kb. *Am. J. Sci.*, 278, 64-94.
- Lloyd, F. E. and D. K. Bailey (1975) Light element metasomatism of the continental mantle: the evidence and the consequences. In L. H. Ahrens *et al.*, Eds., *Physics and Chemistry of the Earth*, 9, p. 389-416. Pergamon Press, New York.
- Modreski, P. J. and A. L. Boettcher (1973) Phase relationships of phlogopite in the system K₂O-MgO-CaO-Al₂O₃-SiO₂-H₂O to 35 kilobars: a better model for micas in the interior of the Earth. *Am. J. Sci.*, 273, 385-414.
- Mysen, B. O., I. Kushiro and T. Fujii (1978) Preliminary experimental data bearing on the mobility of H₂O in crystalline upper mantle. *Carnegie Inst. Wash. Year Book*, 77, 793-797.
- Nakamura, Y. (1974) The system SiO₂-H₂O-H₂ at 15 kbar. *Carnegie Inst. Wash. Year Book*, 73, 259-263.
- and I. Kushiro (1974) Composition of the gas phase in Mg₂SiO₄-SiO₂-H₂O at 15 kbar. *Carnegie Inst. Wash. Year Book*, 73, 255-258.
- Newton, R. C. and W. E. Sharp (1975) Stability of forsterite + CO₂ and its bearing on the role of CO₂ in the mantle. *Earth Planet. Sci. Lett.*, 26, 239-243.
- Ryabchikov, I. D. and D. H. Green (1979) A study of the origin of potassic magmas. *Contrib. Mineral. Petrol.*, in press.
- Shettel, D. L. (1973) Solubility of quartz in H₂O-CO₂ fluids at 5 kb and 500°-900°C. *EOS*, 54, 480.
- Sobolev, N. V. (1977) Deep-seated inclusions in kimberlites and the problem of the composition of the upper mantle. *Am. Geophys. Union*, Washington, D. C.
- Stewart, D. C. and A. L. Boettcher (1977) Chemical gradients in mantle xenoliths (abstr.). *Geol. Soc. Am. Abstracts with Programs*, 9, 1191-1192.
- Wendlandt, R. F. (1977) The system K₂O-MgO-Al₂O₃-SiO₂-H₂O-CO₂: stability of phlogopite as a function of vapor composition and high pressures and temperatures. *Carnegie Inst. Wash. Year Book*, 76, 441-448.
- Wilshire, H. G., J. E. N. Pike, C. E. Meyer and E. C. Schwarzman (1980) Kaersutite veins in lherzolite xenoliths, Dish Hill and Deadman Lake, California. *Am. J. Sci.*, in press.
- Wyllie, P. J. (1979) Magmas and volatile components. *Am. Mineral.*, 64, 469-500.
- Yoder, H. S., Jr. and I. Kushiro (1969) Melting of a hydrous phase: phlogopite. *Am. J. Sci.*, 267-A, 558-582.

Manuscript received, June 26, 1979;
accepted for publication, March 3, 1980.