

Margarite pseudomorphs after kyanite in Glen Esk, Scotland

EMM. BALTATZIS

*Institute of Mineralogy and Petrology
The National University of Athens
Panepistimiopolis, Ano Ilissia
Athens 621, Greece*

AND C. KATAGAS

*Department of Geology
University of Patras
Patras, Greece*

Abstract

This paper describes a new occurrence of margarite in a graphitic gneiss near Fettercairn village, Glen Esk, Scotland. Margarite occurs as pseudomorphs after kyanite and contains considerable paragonite component in solid solution (~25.5 mole %). The data presented enable establishment of another tie line on the white-mica plane of the system $\text{Al}_2\text{O}_3\text{-CaO-Na}_2\text{O-K}_2\text{O-SiO}_2\text{-H}_2\text{O}$.

Introduction

Recognition of margarite as a common rock-forming white mica resulted from the application of X-ray diffraction and electron probe analysis techniques to petrological problems. The number of reported margarite occurrences is continuously increasing and margarite is now considered as a widespread mineral, encountered in various types of low to medium grade metamorphic rocks (Hietanen, 1963, p. B24–B25; Sagon, 1967, 1970; Frey and Niggli, 1972; Ackermann and Morteani, 1973; Höck, 1974; Chinner, 1974; Fox, 1975; Guidotti and Cheney, 1976; Frey, 1978; Teale, 1979; Guidotti *et al.*, 1979).

Difficulties in interpretation of the various reported margarite-bearing assemblages led in turn to the application of experimental methods to these problems. Experimental data on margarite and its phase relations are becoming increasingly available (Velde, 1971; Wall and Essene, 1972; Storre and Nitsch, 1974; Chatterjee, 1974, 1976).

This paper describes a new occurrence of margarite in Glen Esk, Scotland, as pseudomorphs after kyanite, and presents data which establish another tie line on the white-mica plane of the system $\text{Al}_2\text{O}_3\text{-Na}_2\text{O-K}_2\text{O-CaO-SiO}_2\text{-H}_2\text{O}$.

Geologic setting

The margarite-bearing rock was collected 8 km NW of Fettercairn village (Fig. 1). The area studied lies within the region made classic by Barrow's work (1893, 1912) on metamorphic zones and falls within parts of Barrow's original staurolite, kyanite, and sillimanite zones. A more recent account on the metamorphic history of Dalradian rocks in Glens Clova, Esk and Lethnot, Angus, Scotland has been presented by Harte and Johnson (1969).

Petrography

Margarite was found in a kyanite gneiss which contains the assemblage kyanite–muscovite–chlorite (retrograde)–plagioclase–quartz–graphite–tourmaline–apatite and opaques. The kyanite crystals are commonly twinned and have been partially replaced by margarite (Fig. 2). Fine-grained plagioclase is sometimes present as an additional mineral in the pseudomorphs. Porphyroblasts of plagioclase, muscovite, retrograde chlorite (after biotite), and quartz are developed at the border of the pseudomorphs and form the main groundmass of the rock. Graphite, tourmaline, apatite, and opaques are common accessories. Careful microscopic observation, X-ray dif-




Fig. 1. Map showing the locality (□) of the collected sample.

fractograms, and microprobe analyses failed to reveal the presence of paragonite.

Mineral data

Mineral compositions were determined with a Link Systems Model 290-2KX energy-dispersive spectrometer, attached to a Cambridge Geoscan. The spectra were processed using a commercial version of the program developed by P. J. Statham, Department of Mineralogy and Petrology, Cambridge.




Fig. 2. Margarite pseudomorphs after kyanite (Glen Esk); K = kyanite, M = Margarite; (x4) (crossed polars).

Table 1. Microprobe analyses of coexisting margarite and muscovite from Glen Esk, Scotland

	Margarite	Muscovite
SiO ₂	32.59	45.73
TiO ₂	—	0.29
Al ₂ O ₃	50.33	34.20
FeO	0.24	2.77
MnO	—	—
MgO	—	0.73
CaO	9.79	—
Na ₂ O	1.96	1.81
K ₂ O	0.48	8.75
Total	95.39	94.29
Ions to 22(0).		
Si	4.292	6.167
Al ^{iv}	3.708	1.833
Z	8.000	8.000
Al ^{vi}	4.104	3.604
Ti	—	0.030
Fe ⁺²	0.026	0.312
Mn	—	—
Mg	—	0.147
Y	4.130	4.093
Ca	1.382	—
Na	0.501	0.473
K	0.081	1.507
X	1.964	1.980
Paragonite	25.50	23.88
Mol% Margarite	70.36	—
Muscovite	4.14	76.12

Representative analyses of coexisting margarite and muscovite are given in Table 1. Margarite contains negligible muscovite and muscovite does not contain any margarite component in solid solution. On the other hand, the margarite–muscovite pair contains substantial paragonite component in solid solution; nearly equal amounts (~25 mole %) of paragonite solid solution have been detected in each phase (Table 1). Similar observations have been also reported by other workers (Höck, 1974; Fox, 1975; Guidotti and Cheney, 1976; Teale, 1979; Guidotti *et al.*, 1979). Figure 3 is a Na–Ca–K plot of the margarite–muscovite pair from the sample analyzed, with SiO₂ in excess and a H₂O-rich fluid assumed present.

Discussion

The tie lines of the margarite–muscovite pairs reported by Höck (1974) and Teale (1979) (H-148/70 and 454/382) respectively, as plotted on the Na–Ca–K diagram, fall outside the three-phase field depicted by Guidotti *et al.* (1979). In contrast, the tie line derived from our study and one of Teale's (1979) tie lines (454/1301) fall within this field (Fig. 3). The




Fig. 3. Coexisting margarite–muscovite pair from this study plotted on the Na–Ca–K (end-member white micas) diagram; also shown are margarite–muscovite pairs from Höck (1974) (H-148/70), Teale (1979) (454/382, 454/1301) and Guidotti and Cheney (1976) (Ra-a86'-66). Dashed lines show average end-member values for margarite, paragonite, and muscovite pairs reported by Guidotti *et al.* (1979).

margarite–muscovite tie line (H-148/70) given by Höck (1974) is of nearly the same orientation as that (Ra-a86'-66) given by Guidotti and Cheney (1976). The latter, however, crosses the Guidotti *et al.* (1979) muscovite–margarite tie line (Fig. 3) for coexisting muscovite–paragonite–margarite, implying that the amount of paragonite solid solution in margarite of the three-mica assemblage is lower than that in the muscovite–margarite assemblage.

The data are conflicting and no firm conclusion can be reached on the amount of paragonite solid solution in margarite and muscovite from paragonite-bearing and paragonite-free rocks. These amounts probably depend on specific conditions of metamorphism. Our results, however, suggest a more extensive paragonite solid solution in coexisting margarite–muscovite pairs than that found in the three-mica specimens reported by Guidotti *et al.* (1979).

The fact that in most of the margarite occurrences reported so far, margarite appears as a pseudomorph replacement of aluminosilicate polymorphs (Chinner, 1974; Lanphere and Albee, 1974; Guidotti and Cheney, 1974; Teale, 1979; Guidotti *et al.*, 1979)

is of great interest. The replacement of the aluminosilicate by margarite is by no means puzzling, inasmuch as both minerals have the same Al:Si ratio and only CaO and H₂O need to be added, as suggested by Burt in Guidotti *et al.* (1979). The CaO may well be contributed by the plagioclase (Ab₇₈An₂₂) in the groundmass (Guidotti *et al.*, 1979). Nevertheless, difficulties arise in the interpretation of the replacement textures. The common occurrence of margarite associated with a pseudomorph-producing event suggests that, in some cases, margarite may not be a primary phase of the prograde sequence and is instead related to polymetamorphism.

Further data are needed on the phase relations of margarite in the system Al₂O₃–CaO–Na₂O–K₂O–SiO₂–H₂O. Precise data on the metamorphic histories and the metamorphic conditions of areas where margarite is developed will likewise be needed before an understanding of the petrologic implications of margarite appearance can be reached.

Acknowledgments

This work was carried out at the Geology Department, University of Manchester. E. B. would like to thank Dr. B. J. Wood for

his advice and Professor J. Zussman for allowing the use of analytical facilities.

References

- Ackermand, D. and Morteani, G. (1973) Occurrence and breakdown of paragonite and margarite in the Greiner Schiefer series (Zillertal Alps, Tyrol). *Contributions to Mineralogy and Petrology*, 40, 293–304.
- Barrow, G. (1893) On an intrusion of muscovite biotite gneiss in the southeast Highlands of Scotland and its accompanying metamorphism. *Quarterly Journal of the Geological Society of London*, 49, 330–358.
- Barrow, G. (1912) On the geology of lower Deeside and the southern Highland Border. *Proceedings of the Geological Association*, 23, 268–273.
- Chatterjee, N. D. (1974) Synthesis and upper thermal stability limit of 2M-margarite, $\text{CaAl}_2(\text{Al}_2\text{SiO}_{10})(\text{OH})_2$. *Schweizerische Mineralogische und Petrographische Mitteilungen*, 54, 753–767.
- Chatterjee, N. D. (1976) Margarite stability and compatibility relations in the system $\text{CaO}-\text{Al}_2\text{O}_3-\text{SiO}_2-\text{H}_2\text{O}$ as a pressure-temperature indicator. *American Mineralogist*, 61, 699–709.
- Chinner, G. A. (1974) Dalradian margarite: a preliminary note. *Geological Magazine*, 111, 75–78.
- Fox, J. S. (1975) Three-dimensional isograds from the Lukmanier Pass, Switzerland, and their tectonic significance. *Geological Magazine*, 112, 547–564.
- Frey, M. (1978) Progressive low-grade metamorphism of a black shale formation, Central Swiss Alps, with special reference to pyrophyllite and margarite bearing assemblages. *Journal of Petrology*, 19, 93–135.
- Frey, M. and Niggli, E. (1972) Margarite, an important rock-forming mineral in regionally metamorphosed low-grade rocks. *Naturwissenschaften*, 59, 214–215.
- Guidotti, C. V. and Cheney, J. T. (1976) Margarite pseudomorphs after chiastolite in the Rangeley area, Maine. *American Mineralogist*, 61, 431–434.
- Guidotti, C. V., Post, J. L. and Cheney, J. T. (1979) Margarite pseudomorphs after chiastolite in the Georgetown area, California. *American Mineralogist*, 64, 728–732.
- Harte, B. and Johnson, M. R. W. (1969) Metamorphic history of Dalradian rocks in Glens Clova, Esk and Lethnot, Angus, Scotland. *Scottish Journal of Geology*, 5, 54–80.
- Hietanen, A. (1963) Anorthosite and associated rocks in the Boehls Butte quadrangle and vicinity, Idaho. U.S. Geological Survey Professional Paper, 344-B, B1–B78.
- Höck, V. (1974) Coexisting phengite, paragonite and margarite in metasediments of the Mittlere Hohe Tauern, Austria. *Contributions to Mineralogy and Petrology*, 43, 262–273.
- Lanphere, M. A. and Albee, A. L. (1974) $^{40}\text{Ar}/^{39}\text{Ar}$ age measurements in the Worcester Mountains: evidence of Ordovician and Devonian metamorphic events in northern Vermont. *American Journal of Science*, 274, 545–555.
- Sagon, J. P. (1967) Le métamorphisme dans le Nord-Est du bassin de Châteaulain découverte de chloritoïde et de margarite dans les schistes Dévonien. *Comptes Rendus de la Société géologique de France*, 5, 206–207.
- Sagon, J. P. (1970) Mineralogie des schistes paléozoïques du bassin de Châteaulain (Massif armoricain): distribution de quelques minéraux phylliteux de métamorphisme. *Comptes Rendus de l'Académie des Sciences (Paris)*, 270, 1853–1856.
- Storre, B. and Nitsch, K. H. (1974) Zur Stabilität von Margarit im System $\text{CaO}-\text{Al}_2\text{O}_3-\text{SiO}_2-\text{H}_2\text{O}$. *Contributions to Mineralogy and Petrology*, 43, 1–24.
- Teale, G. S. (1979) Margarite from the Olary Province of South Australia. *Mineralogical Magazine*, 43, 433–435.
- Velde, B. (1971) The stability and natural occurrence of margarite. *Mineralogical Magazine*, 38, 317–323.
- Wall, V. J. and Essene, E. J. (1972) Subsolidus equilibria in $\text{CaO}-\text{Al}_2\text{O}_3-\text{SiO}_2-\text{H}_2\text{O}$. (abstr.) *Geological Society of America Abstracts with Programs*, 4, 700.

*Manuscript received, April 15, 1980;
accepted for publication, July 16, 1980.*