

Ilvaite, an alteration product replacing olivine in the Skaergaard intrusion

H. R. NASLUND, J. M. HUGHES,¹ AND R. W. BIRNIE

Department of Earth Sciences, Dartmouth College
Hanover, New Hampshire 03755

Abstract

Ilvaite, $\text{CaFe}_2^{2+}\text{Fe}^{3+}(\text{OH})\text{Si}_2\text{O}_8$, is an abundant mineral in the upper part of the Skaergaard intrusion of East Greenland. It occurs in these rocks as an alteration product replacing fayalitic olivine. Analyses of unaltered olivine and ilvaite demonstrate that the replacement process requires the removal of FeO and MgO from the olivine and the addition of CaO, H₂O, and O₂. Single crystal X-ray diffraction studies indicate that the Skaergaard ilvaite is the orthorhombic polymorph. It is suggested that the ilvaite crystallized under conditions of high $f\text{H}_2\text{O}$, low $f\text{CO}_2$, low $f\text{O}_2$, and moderate T during the subsolidus cooling of the intrusion.

Introduction

Numerous detailed studies have been made of olivine alteration products (Wilshire, 1958; Gay and LeMaitre, 1961; Haggerty and Baker, 1967; Baker and Haggerty, 1967; Champness, 1970; Ganyuchenkov and Barabanov, 1972; Kohlstedt and Vander Sande, 1975; Delvigne *et al.*, 1979; and Moseley, 1981). Although these studies have described several phases which occur as deuteric alteration products of olivine, none of them has reported ilvaite. The only reported occurrences of ilvaite in altered olivine are in the Skaergaard intrusion, in olivine diabase from Elfdalen, Sweden, and in the Basistoppen sill which has intruded the upper part of the Skaergaard intrusion (Ramdohr, 1969; Douglas, 1964). Ilvaite, a relatively rare mineral, typically occurs as a late forming mineral in Ca-Fe-Si skarn deposits (Burt, 1971). The ilvaite in the Skaergaard intrusion is believed to have formed from the interaction of olivine and a fluid phase and is probably the result of an unusual set of conditions which operated during the cooling of the pluton.

Occurrence

Ilvaite is a common alteration product of olivine in the Upper Border Group γ (UBG γ) unit of the Skaergaard intrusion. This unit has been described as a mixture of "ferrodiorite" and "melanogranophyre" (Wager and Brown, 1967), but most samples would be classified as quartz diorite or tonalite using the system of Streckeisen (1976). UBG γ rocks are medium to coarse-grained with rare finer-grained layers and are composed primarily of plagioclase (andesine to oligoclase), clinopyroxene (ferroaugite to ferrohedenbergite), fayalitic olivine, quartz,

magnetite-ulvöspinel, ilmenite-hematite, apatite, and micropegmatite (Naslund, 1980). Zircon and titanite are accessory minerals; ilvaite, chlorite, serpentine, sericite, epidote, titanite, rutile, hornblende, and calcite are present as alteration minerals.

The Skaergaard ilvaite is opaque in thin section and occurs as anhedral patches around the margins and along fractures in olivine crystals (Fig. 1). The ilvaite forms patches up to 1.5 mm across and in some samples replaces up to 75% of individual olivine crystals. Serpentine is commonly associated with ilvaite in Skaergaard olivines, but the proportions of ilvaite and serpentine vary from sample to sample. Hematite was detected in the X-ray diffraction mount of one sample, but it is not an abundant phase in these rocks. Olivine grains adjacent to plagioclase, apatite, pyroxene, and/or quartz may contain ilvaite. No correlation was observed between the adjacent mineral type and the amount of ilvaite present in the olivine. In some samples, olivine grains adjacent to plagioclase were surrounded by a rim of ferroaugite 0.1 to 0.5 mm in thickness. In these samples, contacts between olivine and plagioclase have developed the sequence plagioclase, ferroaugite, ilvaite, olivine or plagioclase, ferroaugite, serpentine, ilvaite, olivine (Fig. 2).

X-ray crystallography

The ilvaite from the Skaergaard intrusion was first identified by comparison of its diffraction pattern with that of JCPDS #25-149. The diffraction pattern was obtained from a polycrystalline grain mount in a 114.6 mm Gandolfi camera, using $\text{CrK}\alpha$ in a helium atmosphere.

In a detailed crystallographic study of the compound, Dietrich (1972) noted that there are two polymorphs of ilvaite. The orthorhombic polymorph occurs in a columnar and more coarsely-crystalline form. The monoclinic

¹ Present address: Department of Geology, Miami University, Oxford, Ohio 45056.





Fig. 1. Olivine grain in sample KG-247 75% replaced by ilvaite. Photo includes: grains of olivine (ol), ilvaite (il), plagioclase (pl), and apatite (ap); and patches of serpentine (se) and myrmekite (my).

Fig. 2. Margin of olivine grain in sample KG-131 with the contact sequence plagioclase (pl)–ferroaugite (f)–serpentine (se)–ilvaite (il)–olivine (ol).

form (Dietrich, 1972) crystallizes in a fine felt-like aggregate. Dietrich noted that both polymorphs occur together in the contact metamorphic ore deposits at Gruba, Oberhalbstein, Switzerland; and he distinguished the monoclinic form by its β angle of 90.22° and the concomitant separation of hkl and $hk\bar{l}$ diffraction peaks.

Three lines of evidence suggest that the Skaergaard ilvaite is the orthorhombic polymorph. First, precise lattice constants were determined by a least-squares refinement of 12 reflections obtained from the Gandolfi pattern. The refinement used monoclinic cell constraints. The β angle was refined to 90.10 (17°) which deviates by less than 0.6σ from 90° . Second, no diffraction lines on the Gandolfi pattern showed hkl – $hk\bar{l}$ splitting. Third, the coarsely crystalline equant habit of the grains is consistent with Dietrich's orthorhombic polymorph.

The space group of orthorhombic ilvaite is $Pbnm$ (#62) (Dietrich, 1972)². The space group of the monoclinic polymorph has not been reported. Single crystal precession photographs of the Skaergaard ilvaite were taken

using $MoK\alpha$ radiation. The precession suite showed diffraction symmetry consistent with space group $Pbnm$ (#62) or $P2_1nb$ (#33). The centrosymmetric space group $Pbnm$ was chosen because it was used in the structure determinations of Takeuchi (1948) and Belov and Mokejeva (1951), and we had no evidence to indicate that the crystal is not centrosymmetric.

Chemical composition

Major element analyses of coexisting ilvaite and olivine in three Skaergaard samples were performed using a MAC electron microprobe with Krisel automation and

² Dietrich actually assigned space group $Pbmn$ (#53) to Strunz's (1937) cell parameters, but this space group is not equivalent to $Pcmm$ (#62) reported by Strunz (1944) by any axis transformation of Strunz's original parameters. We presume this is an oversight caused by transposing the glide and mirror plane symbols. The same error is present in Takeuchi's (1948) study where the correct Schoenflies notation is given for space group $Pbnm$ (#62), but the space group symbol is given as $Pbmn$ (#53).

Table 1. Electron microprobe chemical analyses of Skaergaard ilvaite

	71-07	131-14	247-05	Average	Theoretical
†Oxide Weight Percent					
SiO ₂	28.48	28.91	28.13	28.51	29.40
TiO ₂	0.05	0.12	0.06	0.08	—
Al ₂ O ₃	0.06	0.06	0.63	0.25	—
*Fe ₂ O ₃	19.37	19.04	19.38	19.26	19.53
*FeO	34.87	34.30	34.87	34.68	35.15
MnO	1.08	0.59	0.93	0.87	—
MgO	0.20	0.38	0.32	0.30	—
CaO	12.54	12.96	13.35	12.95	13.72
**H ₂ O	2.18	2.22	2.19	2.20	2.20
Total	98.83	98.58	99.86	99.10	100.00
Cations per 9 Oxygens					
Si	1.97	1.99	1.93	1.96	2.00
Ti	0.00	0.00	0.00	0.00	—
Al	0.00	0.00	0.05	0.02	—
Fe ³⁺	1.01	0.99	1.00	1.00	1.00
Fe ²⁺	2.02	1.97	2.00	2.00	2.00
Mn	0.06	0.02	0.06	0.05	—
Mg	0.02	0.02	0.02	0.02	—
Ca	0.93	0.96	0.98	0.96	1.00
H	1.00	1.00	1.00	1.00	1.00
†Cr ₂ O ₃ , V ₂ O ₅ , P ₂ O ₅ , Na ₂ O, and K ₂ O <.05% each					
*Fe ²⁺ /Fe ³⁺ assumed to be 2/1					
**H ₂ O calculated based on 1 H ⁺ per 9 oxygen					

Bence-Albee on-line data reduction (Finger, 1974; Bence and Albee, 1968). The accelerating voltage was 15 kV, the sample current was 500 mA, and the counting limits were set at 30 seconds or 30,000 counts. The analyses are

Table 2. Electron microprobe chemical analyses of Skaergaard olivines

	KG-71	KG-131	KG-247
†Oxide Weight Percent			
SiO ₂	30.05	31.89	29.58
TiO ₂	0.09	0.06	0.10
Al ₂ O ₃	0.02	0.01	0.01
FeO	65.98	57.90	62.97
MnO	1.24	0.93	1.45
MgO	2.65	9.68	3.07
CaO	0.36	0.25	0.49
Total	100.39	100.72	97.67
Cations per 4 Oxygen			
Si	0.994	0.997	0.999
Ti	0.002	0.001	0.003
Fe ²⁺	1.825	1.514	1.779
Mn	0.035	0.025	0.041
Mg	0.131	0.451	0.155
Ca	0.013	0.008	0.018
End member calculations			
FO%	6.56	22.67	7.83
FA%	93.44	77.33	92.17
†Na ₂ O, K ₂ O, and P ₂ O ₅ <.05% each			

presented in Tables 1 and 2. The proportions of FeO and Fe₂O₃ in ilvaite were calculated from ΣFeO assuming Fe²⁺/Fe³⁺ = 2.0. The proportion of H₂O in ilvaite was calculated assuming one hydrogen per nine oxygen in the unit cell.

Discussion

Studies by Burt (1971) on the stability of ilvaite in Ca-Fe-Si skarns indicate that ilvaite is stable under conditions of relatively high $f\text{H}_2\text{O}$, low $f\text{CO}_2$, moderate to low temperature, and low $f\text{O}_2$. Available data suggest that these conditions occurred during the subsolidus cooling of the Skaergaard intrusion. Cooling models based on oxygen isotope data (Taylor and Forrester, 1979; Norton and Taylor, 1979) indicate that the Skaergaard intrusion was in contact with circulating groundwater during its post-magmatic cooling. These models suggest that the solidified Skaergaard intrusion cooled from 1000°C to 600°C during a period of approximately 70,000 years, cooled to 310°C after an additional 100,000 years, and cooled to near ambient temperature (230°C) during the next 200,000 years. Oxygen fugacity during the cooling of the Skaergaard rocks was probably buffered by reactions between quartz, olivine, magnetite, and ilmenite. Calculations based on analyzed UBG oxide minerals (Naslund, 1980) and experimental data (Buddington and Lindsley, 1964) suggest that the UBG rocks cooled under relatively reducing conditions near the quartz-fayalite-magnetite temperature- $f\text{O}_2$ curve (Fig. 3). Calcite is not common in UBGγ samples suggesting that $f\text{CO}_2$ was low during the hydrothermal alteration of these rocks. Estimates of total





Fig. 3. Calculated temperatures and oxygen fugacities from exsolved magnetite-ulvöspinel and ilmenite-hematite in UBG samples (Naslund, 1980). Calculations are based on the experimental data of Buddington and Lindsley (1964). The buffer curves for magnetite-hematite (MH), quartz-fayalite-magnetite (QFM), and quartz-fayalite-iron (QFI) are from Eugster and Wones (1962). UBG oxide minerals are presumed to have reequilibrated during subsolidus cooling.


pressure at the UBG γ horizon during crystallization range from 500 bars to 1000 bars (Lindsley, *et al.*, 1969; Williams, 1971).

The UBG γ unit of the Skaergaard intrusion has been intruded by the post Skaergaard Basistoppen intrusion, a strongly differentiated, gabbroic sill approximately 500 m thick. Several lines of evidence indicate that the Basistoppen magma was emplaced before the Skaergaard intrusion had completely cooled (Douglas, 1964; Taylor and Forrester, 1979; Schwarz *et al.*, 1979). Heat from the cooling Basistoppen magma may have contributed to the hydrothermal alteration of the Skaergaard olivines. Olivine from the Basistoppen intrusion also contains ilvaite as an alteration mineral (Douglas, 1964).


The composition of ilvaite and unaltered olivine indicate that the alteration process requires the addition of CaO, H₂O, and O₂ and the removal of MgO and FeO as follows:




The reaction could also be written as:



If serpentine is involved, the reaction could proceed as follows:



or:



Free oxygen in these reactions could also be expressed by the addition of H₂O and the removal of H₂, and variations in the Fe–Mg ratios of the olivine and serpentine will affect the quantities of the reactants and products. In any case, however, CaO, H₂O, and O₂ must have been added to the olivine during the alteration process. The distribution of ilvaite around the margins of olivine grains suggest that the ilvaite grew from the margins inward. There is no evidence, however, to suggest that CaO was derived from any immediately adjacent CaO-bearing mineral phases. Chemical components were probably transported to and away from the site of alteration in a fluid phase traveling along grain boundaries.

Ilvaite is not a common mineral phase, and its occurrence as an alteration product of olivine is particularly rare. The conditions which led to the formation of ilvaite in the Skaergaard intrusion probably included: 1) emplacement of the magma at a depth sufficient to allow relatively slow subsolidus cooling, but shallow enough to

allow the circulation of ground water; 2) emplacement into permeable country rock; and 3) the development of extreme iron-enrichment and the crystallization of iron-rich olivine (fayalite 80–100%). It is believed that the transfer of material into and away from the site of alteration was facilitated by the circulation of hydrothermal fluids. Parts of the Skaergaard intrusion which contain fayalitic olivine, but which are adjacent to relatively impermeable gneissic country rock, do not contain ilvaite. Likewise, parts of the Skaergaard intrusion which are adjacent to relatively permeable basaltic country rock, but which contain more magnesium-rich olivine, contain serpentine but do not contain ilvaite.

Acknowledgments

The chemical analyses were performed at the Geophysical Laboratory of the Carnegie Institution of Washington, D.C., while H. R. Naslund was a predoctoral fellow. The X-ray crystallography was supported by the Committee on Research and the Department of Earth Sciences, Dartmouth College. G. D. Johnson helped with the preparation of the photomicrographs at Dartmouth. The manuscript benefited from a critical review by D. M. Burt.

References

- Baker, I. and Haggerty, S. E. (1967) The alteration of olivine in basaltic and associated lavas; Part II: Intermediate and low temperature alteration. *Contributions to Mineralogy and Petrology*, 16, 258–273.
- Belov, N. V. and Mokejeva, V. I. (1951) Crystal structure of ilvaite. *Akademi Nauk SSSR Doklady*, 81, 581–584. (in Russian).
- Bence, A. E. and Albee, A. L. (1968) Empirical correction factors for the electron microanalysis of silicates and oxides. *Journal of Geology*, 76, 382–403.
- Buddington, A. F. and Lindsley, D. H. (1964) Iron–titanium oxide minerals and synthetic equivalents. *Journal of Petrology*, 5, 310–357.
- Burt, D. M. (1971) Multisystems analysis of the relative stabilities of babingtonite and ilvaite. *Annual Report of the Director, Geophysical Laboratory, Carnegie Institution Year Book*, 70, 189–197.
- Champness, P. E. (1970) Nucleation and growth of iron oxides in olivine, (Mg, Fe)₂SiO₄. *Mineralogical Magazine*, 37, 790–800.
- Delvigne, J., Bisdorf, E. B. A., Sleeman, J., and Stoops, G. (1979) Olivines, their pseudomorphs and secondary products. *Pedologie*, 29, 247–309.
- Dietrich, V. (1972) Ilvait, Ferroantigorit und Greenalith als Begleiter oxidisch-sulfidischer Vererzungen in den Oberhalbsteiner Serpentiniten. *Schweizerische Mineralogische und Petrographische Mitteilungen*, 52, 57–74.
- Douglas, J. A. V. (1964) Geological investigations in East Greenland: Part VII the Basistoppen sheet—a differentiated basic intrusion into the upper part of the Skaergaard complex, East Greenland. *Meddelelser om Gronland*, 164, 1–66.
- Eugster, H. P. and Wones, D. R. (1962) Stability relations of the ferruginous biotite, annite. *Journal of Petrology*, 3, 82–125.
- Finger, L. W. (1974) The Geophysical Laboratory microprobe operating system—operator's manual. Carnegie Institute of Washington.

- Garryuchenkov, F. G. and Barabanov, V. F. (1972) Postmagmatic transformation of olivine (abst.). *Geochemistry International*, 8, 306.
- Gay, P. and LeMaitre, R. W. (1961) Some observations on iddingsite. *American Mineralogist*, 46, 92–111.
- Haggerty, S. E. and Baker, I. (1967) The alteration of olivine in basaltic and associated lavas; Part I: high temperature alteration. *Contributions to Mineralogy and Petrology*, 16, 233–257.
- Kohlstedt, D. L. and Vander Sande, J. B. (1975) An electron microscopy study of naturally occurring oxidation produced precipitates in iron-bearing olivines. *Contributions to Mineralogy and Petrology*, 53, 13–24.
- Lindsley, D. H., Brown, G. M. and Muir, I. D. (1969) Conditions of the ferrowollastonite–ferrohedenbergite inversion in the Skaergaard intrusion, East Greenland. *Mineralogical Society of America Special Paper* 2, 193–201.
- Moseley, D. (1981) Ilmenite exsolution in olivine. *American Mineralogist*, 66, 976–979.
- Naslund, H. R. (1980) Part I: Petrology of the Upper Border Group of the Skaergaard Intrusion, East Greenland, and Part II: An Experimental Study of Liquid Immiscibility in Iron-bearing Silicate Melts. Ph.D. Dissertation, University of Oregon.
- Norton, D. and Taylor, H. P., Jr. (1979) Quantitative simulation of the hydrothermal systems of crystallizing magmas on the basis of transport theory and oxygen isotope data: An analysis of the Skaergaard intrusion. *Journal of Petrology*, 20, 421–486.
- Ramdohr, P. (1969) *The Ore Minerals and Their Intergrowths*. Pergamon Press, Oxford, England.
- Schwarz, E. J., Coleman, L. C. and Cattroll, H. M. (1979) Paleomagnetic results from the Skaergaard intrusion, East Greenland. *Earth and Planetary Science Letters*, 42, 437–443.
- Streckeisen, A. (1976) To each plutonic rock its proper name. *Earth-Science Reviews*, 12, 1–33.
- Strunz, H. (1937) Über Kristallographie und chemische Zusammensetzung von Lawsonit und Lieverit. *Zeitschrift für Kristallographie*, 96, 504–506.
- Strunz, H. (1941) *Mineralogische Tabellen*. Akad. Verlagsgesellschaft Becker and Erler Kom. Ges., Leipzig.
- Takeuchi, Y. (1948) The structure of ilvaite. *X-Sen (X-Rays)*, 5, 8–14. (seen in *Chemical Abstracts* 44, 5766, 1950.)
- Taylor, H. P., Jr. and Forrester, R. W. (1979) An oxygen and hydrogen isotope study of the Skaergaard intrusion and its country rocks: a description of a 55-m.y. old fossil hydrothermal system. *Journal of Petrology*, 20, 355–419.
- Wager, L. R. and Brown, G. M. (1967) *Layered Igneous Rocks*. W. H. Freeman and Co., San Francisco, California.
- Williams, R. J. (1971) Reaction constants in the system Fe–MgO–SiO₂–O₂: intensive parameters in the Skaergaard intrusion, East Greenland. *American Journal of Science*, 271, 132–146.
- Wilshire, H. G. (1958) Alteration of olivine and orthopyroxene in basic lavas and shallow intrusions. *American Mineralogist*, 43, 120–147.

*Manuscript received, June 18, 1982;
accepted for publication, January 10, 1983.*