

Clinoenstatite-bearing lava from Népoui, New Caledonia

T. SAMESHIMA

Department of Geology, University of Auckland
Auckland, New Zealand

J.-P. PARIS

Bureau de Recherches Géologiques et Minières
B.P. 56, Nouméa, New Caledonia

PHILIPPA M. BLACK AND R. F. HEMING¹

Department of Geology, University of Auckland
Auckland, New Zealand

Abstract

A vitrophyric clinoenstatite-bearing lava associated with tholeiitic basalts has been found lying immediately below an overthrust peridotite sheet near Népoui, on the west coast of New Caledonia. The lava consists mainly of clinoenstatite phenocrysts (29%), sometimes enclosing euhedral chromite, a lesser amount of microphenocryst bronzite (20%) and a glassy mesostasis. Clinoenstatite phenocrysts are compositionally homogeneous ($\text{En}_{89}\text{Fs}_{10.5}\text{Wo}_{0.5}$) with polysynthetic twins on (100), a feature which is believed to indicate inversion from protoenstatite. Bronzite is variable in composition with $\text{En}_{74-84}\text{Fs}_{14-22}\text{Wo}_{1.5-4}$. Hypersthene ($\text{En}_{43-61}\text{Fs}_{36-52}\text{Wo}_{3-4}$), calcic and subcalcic augite and iron-rich hornblende occur as overgrowths on phenocrysts and as quench microlites.

The New Caledonian clinoenstatite boninite is similar chemically, mineralogically, texturally and in terms of age (Eocene–Oligocene) and tectonic setting (with ophiolite complexes) to the three other known clinoenstatite boninite localities (Cape Vogel, Papua New Guinea, Bonin Islands and Mariana Trench). Clinoenstatite boninite magmas are believed to result from extensive partial melting of depleted mantle peridotite under hydrous conditions and an abnormally high geothermal gradient.

Introduction

A vitrophyric clinoenstatite-bearing rock was found by one of the authors (J.-P. Paris) in a road cut a few tens of meters north of Riviere Blanche, on the main road (RT-1) north of Népoui on the west coast of New Caledonia and also *in situ* nearby in the bed of Riviere Blanche (Fig. 1).

The exposure in the road cut shows a mylonitic serpentinite sheet, an outlier of the ultramafic nappe overthrust on to the sialic basement of New Caledonia in the Upper Eocene (Paris *et al.*, 1979). The serpentinite overlies the "Formation des Basaltes" which is considered to be para-autochthonous and is composed mainly of basic submarine lavas (tholeiitic basalts and dolerites) with minor acid volcanics (andesite–rhyolite). The "Formation des Basaltes" was erupted during the Upper Creta-

ceous to Lower Eocene close to the site of the west Caledonian fault zone in a geotectonic setting believed by Paris (1981) to be an island arc environment. The clinoenstatite-bearing rock is possibly a flow and a member of the "Formation des Basaltes" but due to the poor exposure the precise relationship with other members of the formation is not clear.

The Népoui lava closely resembles the clinoenstatite-bearing volcanics described from Cape Vogel, Papua New Guinea (Dallwitz *et al.*, 1966; Dallwitz, 1968), the Mariana Trench (Dietrich *et al.*, 1978; Sharaskin and Dobretsov, 1979), and Mukojima and Chichijima, Bonin Islands (Shiraki, *et al.*, 1979; Komatsu, 1980). All of these rocks have petrochemical similarities and are often closely geographically associated with the nonfeldspathic bronzite andesite named "boninite" by Petersen (1891a,b), which, however, lacks clinoenstatite. Sharaskin *et al.* (1979) have proposed the name "marianite" for clinoenstatite-bearing nonfeldspathic andesites; but, if a

¹ Present address: Chevron Resources Company, P.O. Box 7140, San Francisco, California, 94120.

new name is to be applied to this whole group of clinoenstatite lavas, then it should be "vogelite" after the first discovery by Dallwitz *et al.* (1966). Other authors (*e.g.*, Jenner, 1981) prefer to use high magnesian andesites although this term could include feldspathic varieties. However, it seems desirable to use a name which will distinguish the feldspar-free lava types from the more common feldspathic igneous rocks so we have retained the term boninite because the clear mineralogical and chemical distinctions between boninites and the clinoenstatite (CE) rocks can be emphasized satisfactorily by the term "clinoenstatite boninite" (CE-boninite) which is adopted for this paper.

Petrology

The New Caledonia CE-boninite is strongly porphyritic and contains 29% by volume of polysynthetically-twinned clinoenstatite phenocrysts, usually 1–2 mm in length; 20% of orthopyroxene microphenocrysts (0.1 mm diameter) and small amounts of calcic clinopyroxene and hornblende occurring either as slender acicular crystals sometimes forming overgrowths on clinoenstatite and orthopyroxene phenocrysts or as feather-form quench microlites scattered through the glass. Clinoenstatite phenocrysts are sometimes aggregated into stellate clusters and rare centimeter-sized megacrysts are also present. Euhedral chromite grains are enclosed in the clinoenstatite. Tiny magnetite grains in the glass have frequently acted as nuclei for hornblende dendrites.

The glass is usually clear and pale brown in color but occasional spherical brown patches, about a centimeter in diameter, are altered to smectite. Isolated clinoenstatite megacrysts and round altered patches are also common in the Mukojima (Bonin Islands) CE-boninite (Shiraki *et al.*, 1979).

Electron microprobe analyses of representative minerals from the Népoui CE-boninite are listed in Table 1 and the compositional range of the pyroxenes and chromites are shown in Figures 2 and 3, together with comparative mineral analyses from other localities.

The Népoui clinoenstatites are chemically homogeneous with compositions clustered within the very narrow range $En_{87.5-90}Fs_{11-9}Wo_{0.5}$. All the Népoui clinoenstatites are polysynthetically twinned on (100) with habits and textures like those in the Cape Vogel rocks described and figured by Dallwitz *et al.* (1966). For the Cape Vogel clinoenstatite, Sadanaga and Okamura (1969) found that the crystals had the morphology of orthorhombic protoenstatite; heating at 1100°C for 3 hours changed the clinoenstatite into the proto-form.

The Népoui bronzite microphenocrysts show a wider compositional range ($En_{78-83}Fs_{25-22}Wo_{2-4}$) than the clinoenstatites, but are similar to orthopyroxene microphenocrysts in other CE-boninites (Fig. 2) with the orthopyroxenes always having higher Al, Ca and Fe values than the clinoenstatites. The orthopyroxenes in small elongated prisms dispersed throughout the glassy groundmass are




Fig. 1. Map of New Caledonia showing the position of the clinoenstatite boninite locality and its relationship to the tholeiitic "Formation des Basaltes" and the obducted peridotite nappe.

hypersthene. Calcic clinopyroxenes and amphiboles, whether rimming the clinoenstatite and bronzite or forming discrete prisms in the glass, are strongly zoned from Mg-rich cores (or from contacts with clinoenstatite) to Ca, Fe-enriched outer margins; undulose extinction suggests that they are quench crystals. Calcic clinopyroxene analyses plotted in Figure 2 have a large compositional scatter from calcic to subcalcic augite, reinforcing the suggestion that these are disequilibrium compositions formed during quenching.

The Népoui chromites are compositionally variable with crystals ranging from 49–59% Cr_2O_3 , 19–29% FeO^* , 10–12% MgO and less than 10% Al_2O_3 . The high 100 Cr/(Cr+Al) and low 100 Mg/(Mg+Fe) ratios are typical of all CE-boninite chromites in clear contrast to chromites from boninites and komatiites and from ultrabasic plutonic rocks and xenoliths (Crawford, 1980) as shown in Figure 3.

Geochemistry

Table 2 contains modal mineral, chemical and normative compositions of CE-boninites from New Caledonia, Cape Vogel (Dallwitz, 1968) and the Bonin Islands (Shiraki *et al.*, 1979) as well as the average of 21 CE-boninite chemical analyses from Cape Vogel (Jenner, 1981) and 5 from the Mariana Trench (Dietrich *et al.*, 1978). The chemical characteristics common to all these samples are high magnesia, and low values for alumina, lime, alkalis and titania relative to silica content. In comparison with other clinoenstatite rocks, the New Caledonian lavas are somewhat higher in silica and alkalis and slightly lower in iron oxides and lime (*cf.* Fig. 4).

Trace element data for the Népoui CE-boninite are

Table 1. Representative electron microprobe analyses of minerals in Népoui clinoenstatite boninite

	1	2	3	4	5	6	7	8	9	10	11
SiO ₂	56.14	56.04	57.07	54.59	50.84	51.47	49.62	40.72	39.97	0.91	1.48
TiO ₂	0.04	0.13	0.25	0.24	0.15	0.42	0.87	0.27	1.14	0.25	4.46
Al ₂ O ₃	0.14	0.97	1.01	5.01	2.35	3.45	4.18	13.24	6.23	5.53	5.56
Cr ₂ O ₃	0.12	0.31	0.28	nd	-	-	-	-	-	59.14	0.04
FeO*	7.60	9.38	11.96	23.81	16.36	12.75	15.56	23.27	21.73	19.72	79.25
MnO	0.16	0.25	0.26	0.31	0.44	0.25	0.22	0.16	0.30	0.32	0.21
NiO	0.07	-	-	-	-	-	-	-	-	-	-
HgO	35.72	31.02	27.69	10.97	22.06	17.50	10.73	9.06	14.34	11.47	1.23
CaO	0.25	0.83	1.30	1.51	8.28	12.18	18.14	7.31	13.68	0.02	0.35
Na ₂ O	0.01	nd	0.21	0.60	0.03	0.75	0.39	0.97	0.24	-	-
K ₂ O	0.01	nd	0.08	0.47	0.02	0.13	0.03	0.74	0.07	-	-
Total	100.53	98.90	100.11	97.57	100.53	98.90	99.74	95.74	97.75	97.36	92.58

1. clinoenstatite phenocryst $\alpha = 1.667(1)$; $\gamma = 1.676(1)$; $2V\gamma = 46.63^\circ$; $a = 9.61(1) \text{ \AA}$
 $b = 8.87(1) \text{ \AA}$; $c = 5.165(4)$ $\beta = 108.313(8)$

2. small bronzite crystal attached to clinoenstatite;
 3. bronzite microphenocryst
 4. long prismatic hypersthene
 5-7 calcic clinopyroxene
 8. acicular greenish hornblende
 9. brown feathery dendritic hornblende
 10. chromite inclusion in clinoenstatite
 11. magnetite, small groundmass crystal

listed in Table 3, showing the low Ti/Zr and high Cr/Ni and Zr/Y ratios typical of CE-boninite lavas. Compared with the Cape Vogel CE-boninites (Jenner, 1981) the Népoui rock has higher Y, Zr, Nb and La values and lower Ti/Zr and Zr/Y ratios.

Areas of clear glass in the New Caledonian CE-boninite were analyzed by electron probe microanalysis; to avoid surface damage and loss of alkalis the electron beam was defocused to a diameter of about 10 μm and moved over the surface while counting. The analysis, which totalled 92.3%, was recalculated to 100% and is listed in Table 2 together with an analysis of a glass from a clinoenstatite-free boninite from Kuroda and Shiraki (1975).




Fig. 2. Compositions of pyroxenes from clinoenstatite boninites. Pyroxenes with symbols on legend from literature or from Table 1. Small closed circles and x's are additional clinopyroxene and orthopyroxene analyses from the New Caledonian lava.




Fig. 3. Compositions of chromites from clinoenstatite boninites from New Caledonia (closed circles) and Bonin Islands (open squares; Shiraki and Kuroda, 1977; Shiraki *et al.*, 1979). Compositional fields for chromites in xenoliths in basic magmas, and from alpine and stratiform ultramafic complexes are indicated. B & B.K. is compositional fields of chromites from boninites and basaltic komatiites (Cameron *et al.*, 1979). The chromite field for Howque clinoenstatite-bearing pyroxenite is from Crawford (1980).

Table 2. Chemical, normative and modal analyses of clinoenstatite boninites

	New Caledonia Népoui		Papua-New Guinea Cape Vogel			Bonin Is. Muchojima Chichijima		Mariana Trench	CE Boninite Groundmass	Av Boninite	Boninite Glass
	1	2	3	4	5	6	7	8	9	10	11
SiO ₂	57.41	66.93	53.97	54.09	57.48	55.73	53.05	55.58	54.81	54.47	62.91
TiO ₂	0.22	0.29	0.23	0.30	0.25	0.19	0.19	0.19	0.26	0.24	0.08
Al ₂ O ₃	8.80	18.87	7.97	9.55	8.38	10.83	10.81	10.31	10.58	12.31	17.63
Fe ₂ O ₃	1.28		2.57	2.49		1.56	2.21	3.12	2.80	1.30	
FeO	5.79	2.90*	6.95	6.54	9.58	6.81	6.15	5.56	6.22	6.42	6.56
MnO	0.15	0.05	0.21	0.15	0.20	0.28	0.21	0.17	0.21	0.17	0.10
MgO	15.92	0.50	16.03	13.03	17.82	10.55	13.42	13.54	10.38	11.59	1.59
CaO	3.38	4.02	4.79	5.46	4.92	6.60	6.55	5.52	6.06	6.77	6.72
Na ₂ O	1.89	4.55	0.60	0.75	0.84	1.30	0.80	1.77	1.28	1.63	3.82
K ₂ O	0.62	1.89	0.35	0.41	0.33	0.53	0.25	0.75	0.51	0.51	0.56
P ₂ O ₅	0.06		0.06	0.07		0.01	0.01	0.03	0.05	0.04	
H ₂ O±	4.88		6.07	7.36		5.22	6.55		7.32	4.66	
Total	100.40	100.00	99.80	100.20		99.61	100.20		100.48	100.11	100.00
Q	9.52	21.64	11.48	13.93	7.62	13.39	10.67	9.02	14.70	8.39	17.04
Or	3.66	11.17	2.07	2.42	1.95	3.13	1.48	4.43	3.01	3.01	3.49
Ab	15.99	38.49	5.08	6.34	7.11	11.00	6.77	14.97	10.83	13.79	32.32
An	13.70	19.94	18.02	21.48	18.12	22.15	25.17	17.97	21.62	24.77	29.22
En	39.63	1.24	39.91	32.44	44.36	26.27	33.41	33.71	25.84	28.85	3.96
Fs	9.49	4.94	10.65	9.73	17.55	11.42	9.54	7.63	9.15	10.63	12.10
Wo	1.12	-	2.23	2.15	2.62	4.39	3.03	3.85	3.39	3.57	1.72
Mt	1.86	-	3.73	3.61		2.26	3.20	4.52	4.06	1.88	0.15
Il	0.42	0.55	0.44	0.57		0.36	0.36	0.36	0.46	0.45	-
Ap	0.14	-	0.14	0.17		0.02	0.02	0.07	0.12	0.09	-
C	-	2.03	-	-		-	-	-	-	-	-
Olivine	-						4.0				
Clinoenstatite	29.2		19	5		1.2	6.8				
Bronzite (Phene)	-		2	1		-	-				
Orthopyroxene	20.8					16.1	20.7				
Ca-Pyroxene	5.8		14	12		35.3	25.7				
Microlites	-		22	30		-	-				
Hornblende	5.9										
Chromite	0.1		0.3	0.3			0.4				
Zeolite	-		8	12			1.6				
Glass including altered glass	38.0		35	40		47.4	40.1				

1,2 whole rock (1) and groundmass glass (2)

3,4 whole rock analyses LB107 (3) and LB 105 (4) from Dallwitz (1968)

5 Average of 21 whole rock analyses listed by Jenner (1981)

6,7 Whole rock analyses from Shiraki et al. (1979)

8 Average of 5 clinoenstatite boninites listed by Dietrick et al. (1978)

9 Average CE boninite groundmass calculated by subtracting modal clinoenstatite from rock analyses 1,3,4,6 and 7.

10 Average boninite analyses from Bonin Islands (listed by Shiraki (1977)

11 Boninite glass from Kuroda and Shiraki (1975).

The water contents of the CE-boninite and boninite glasses have not been measured directly but can be estimated either from the electron probe analysis (the difference between the total and 100%) as 7.5 and 5.7% respectively for the CE-boninite and boninite glasses, or from the modal analysis and H₂O values of the bulk rock chemical analyses which gives values of 8.5% and 8.7%. The estimated H₂O contents of the glasses (6–9%) are compatible with the low refractive indices of the glasses:

1.520 for the New Caledonian CE-boninite and a range of 1.535–1.550 for a boninite from the Bonin Islands (Kuroda and Shiraki, 1975).

The composition of the CE-boninite glass is dacitic and extremely low in MgO; virtually all the Mg in the rock must be held in the chain silicates (as is also suggested by mineral analyses in Table 1). Compared with the boninite glass, the CE-boninite glass contains less iron and lime and more silica. Calculated groundmass compositions




Fig. 4. Plot of weight percent Al_2O_3 versus MgO and total iron as $\text{FeO} + \text{MgO}$ for all available analyses of clinoenstatite boninites and boninites from the Bonin Islands. Data for the New Caledonian clinoenstatite boninite are indicated by N. En = the average composition of clinoenstatite phenocrysts occurring in boninites.

obtained by subtracting the modal percentage of clinoenstatite from the bulk rock composition are similar to that of average boninites (Table 2).

Plots of wt.% Al_2O_3 versus MgO and $(\text{MgO} + \text{FeO}^*)$ (Fig. 4) for all available CE-boninite whole rock analyses and groundmass and glass compositions for CE-boninites and some data for boninites (Table 2) show strong linearity suggesting simple crystal fractionation and depletion of the liquid of an Fe-Mg phase (clinoenstatite) is the major factor controlling the chemical variation observed in the CE-boninites. The position of the average boninite composition and glass at the high Al_2O_3 , low FeO and MgO section of the plot suggests that addition of clinoenstatite to boninite liquids would change them towards CE-boninite compositions. The data seem to support the idea of Bloomer *et al.* (1979), and Howard and Stolper (1981) that there is a boninite series (including CE-boninite) which is controlled by Fe-Mg pyroxene and/or olivine and spinel fractionation.

Origin of the CE-boninites

Experimental work on the stability of MgSiO_3 includes the enstatite-protonstatite transition which has been determined at 985°C by Atlas (1952) and redetermined at $1042 \pm 3^\circ\text{C}$ by Sarver and Hummel (1962). The presence of the FeSiO_3 molecule apparently raises the inversion temperature (Boyd and Schairer, 1964; Kuno, 1966; Nakamura, 1971). Pressure also raises the inversion temperature and Boyd and England (1965) showed that the protoenstatite field disappears above 8 kbar. More recently, Chen and Presnall (1975) have extended the stability field to 15 kbar at temperatures approaching 1800°C . Sadanaga and Okamura (1969) inverted natural

Cape Vogel clinoenstatite into protoenstatite by heating at 1100°C for 3 hours.

The experimental results indicate that the conditions for crystallization of protoenstatite in a boninite magma would be at least 1100°C at a low pressure, perhaps less than a few kilobars. Dallwitz *et al.* (1966) suggested it was the high MgO content of the CE-boninite that was responsible for the crystallization of protoenstatite in the magma.

The geochemistry of boninites and CE-boninites indicates an origin depleted in incompatible elements and rich in refractory constituents. Melting of refractory mantle is facilitated by the presence of water, and the high water contents of CE-boninites plus the modal amphibole are evidence for hydrous conditions at the source.

Experimental work by Green (1973) has shown that high degrees of melting of pyrolite under water-saturated conditions and relatively low pressures (10 kbar) can yield magnesian, quartz normative basaltic andesites but the melt produced in his experiment had "normal" CaO and Al_2O_3 contents such that plagioclase should be a modal constituent. More recently, Kushiro and Takahashi (1981) have produced a liquid chemically similar to a boninite with about 13% MgO and 59% SiO_2 by partial melting of lherzolite under hydrous conditions at relatively low pressures (12–15 kbar) and high temperatures (1130°C).

It is clear from both the known stability relations of protoenstatite and the partial melting experiments producing a boninite-like liquid, that generation of a clinoen-

Table 3. Trace element compositions of clinoenstatite boninites

Népoûi New Caledonia	Cape Vogel, Papua New Guinea						
	HMA Type C	HMA Type E					
	1*	6	7	9	16	18	
Rb	13	8	9	6	4	1	11
Sr	180	274	96	69	63	59	110
Zr	90	30	62	32	28	42	24
Nb	8	0.88	3.2	1.7	1.3	2.2	1.3
Cr	1390	715	735	2225	2191	2178	1831
V	130	187	211	158	149	163	164
Ni	265	154	212	415	574	415	288
La	7	1.47	4.06	2.88	2.03	3.25	2.10
Ce	<5	3.51	9.74	5.35	4.35	6.45	4.30
Y	9	7	9	3	4	4	3
Rb/Sr	0.07	0.03	0.09	0.09	0.06	0.02	0.1
Cr/Ni	5.1	3.8	3.5	5.4	3.8	5.2	6.3
Ti/Zr	15	60			39		
Ti/Y	144	257			316		
Zr/Y	10	4.3			8.9		
Zr/Nb	11	34			19.4		
La/Nb	0.9	1.7			1.5		
La/Zr	0.09	0.05			0.078		

*Numbers refer to Analyses listed in Tables 2 and 3 of Jenner (1981)

statite boninite requires a hydrated, high temperature environment at shallow depths (less than 15 kbar). How water is introduced into refractory mantle at depths shallow enough to stabilize clinoenstatite, without introducing incompatible elements is an enigma, especially since hydrous metasomatic fluids are widely advocated by petrologists as a method of producing, for example nephelinite enriched in incompatible elements (*e.g.* Menzies and Murthy, 1980; Wass, 1980). Perhaps extensive mantle melting, causing extreme dilution of incompatible elements, would have the same effect (Shiraki *et al.*, 1980) but this requires an abnormally high heat input.

Anomalous heating of the upper mantle should be reflected by some corresponding unusual geological event. Cameron *et al.* (1979) have pointed out that boninites are almost exclusively associated with ophiolites. The CE-boninite occurrences are restricted to Eocene-Oligocene ophiolite associations along the western Pacific margin but in no locality is the relationship between the CE-boninites and associated ophiolites sufficiently clear to point to some common genetic event.

Ideas on the type of geological environment in which CE-boninites were erupted are various and contradictory. Shiraki *et al.* (1980) hypothesized that a possible site for generating clinoenstatite-bearing rocks may be at shallow depths below a spreading ridge or island arc after the main magmatic episode in an environment with an exceptionally steep geothermal gradient where water liberated from the underlying subducted oceanic crust could be introduced into the melting zone. Crawford *et al.* (1981) however, considered that the boninites (some clinoenstatite-bearing) associated with ophiolites and MORB tholeiites of the West Philippine-Mariana region and elsewhere were erupted in a fore-arc position after arc magmatism and before eruption of voluminous MORB tholeiites. In contrast to the above, Paris (1981) concludes that the "Formation des Basaltes" with which the New Caledonian CE-boninites are associated, were erupted in a back-arc expansion zone.

In conclusion, it can be said that CE-boninites are confined to the same tectonic environments as ophiolites and that CE-boninite magmas appear to result from extensive partial melting of depleted mantle peridotite under hydrous conditions and an abnormally high geothermal gradient. CE-boninites are not related, at least in any simple way, to geologic and petrographic processes that give rise to an island arc volcanic suite.

Acknowledgments

We are indebted to Professor R. N. Brothers and Dr. I. E. M. Smith for critical reading of the draft manuscript and their many helpful comments.

References

- Anderson, Olaf. (1915) The system anorthite-forsterite-silica. *American Journal of Science* (4th series), 39, 407-454.
- Atlas, Leon (1952) The polymorphism of $MgSiO_3$ and solid-state equilibria in the system $MgSiO_3$ - $CaMgSi_2O_6$. *Journal of Geology*, 60, 125-147.
- Bloomer, S., Melchior, J., Poreda, R., Hawkins, J. (1979) Mariana arc-trench studies: petrology of boninites and evidence for a boninite series. *Transactions of American Geophysical Union*, (EOS), 60, 968.
- Cameron, W. E., Nisbet, E. G. and Dietrich, V. J. (1979) Boninites komatiites and ophiolitic basalts. *Nature*, 280, 550-553.
- Chen, C-H and Presnall, D. C. (1975) The system Mg_2SiO_4 - SiO_2 at pressures up to 25 kilobars. *American Mineralogist*, 60, 398-406.
- Crawford, A. J. (1980) A clinoenstatite-bearing cumulate olivine pyroxenite from Howqua, Victoria. *Contributions to Mineralogy and Petrology*, 75, 353-368.
- Crawford, A. J., Beccaluva, L. and Serri, G. (1981) Tectonomagmatic evolution of the West Philippine-Mariana region and the origin of boninites. *Earth and Planetary Science Letters*, 54, 346-356.
- Dallwitz, W. B. (1968) Chemical composition and genesis of clinoenstatite-bearing volcanic rocks from Cape Vogel, Papua: a discussion. *Proceedings 23rd International Geological Congress* 2, 229-242.
- Dallwitz, W. B., Green, D. H. and Thompson, J. E. (1966) Clinoenstatite in a volcanic rock from the Cape Vogel area, Papua. *Journal of Petrology*, 7, 375-403.
- Daly, R. A. (1933) *Igneous Rocks and the Depth of the Earth*. McGraw Hill, New York.
- Dietrich, V., Emmermann, R., Oberhansli, R. and Puchelt, H. (1978) Geochemistry of basaltic and gabbroic rocks from the West Mariana basin and the Mariana trench. *Earth and Planetary Science Letters*, 39, 127-144.
- Green, D. H. (1973) Experimental melting studies on model upper mantle composition at high pressure under water-saturated and water undersaturated conditions. *Earth and Planetary Science Letters*, 19, 37-53.
- Greig, J. W. (1927) Immiscibility in silicate melts. *American Journal of Science* (5th series) 13, 1-44, 133-154.
- Howard, A. H. and Stolper, E. (1981) Experimental crystallisation of boninites from the Mariana Trench. *Transactions of the American Geophysical Union* (EOS), 62, 1091.
- Jenner, G. A. (1981) Geochemistry of high-Mg andesites from Cape Vogel, Papua New Guinea. *Chemical Geology*, 33, 307-332.
- Kikuchi, Yasushi (1980) On pyroxene components in certain volcanic rocks from Bonin Islands. (in Japanese). *Journal of Science Imperial University of Japan*, 3, 67-89.
- Komatsu, Masayuki (1980) Clinoenstatite in volcanic rocks from the Bonin Islands. *Contributions to Mineralogy and Petrology* 74, 329-338.
- Kuno, Hisashi (1966) Review of pyroxene relations in terrestrial rocks in the light of recent experimental work. *Mineralogical Journal*, 5, 21-43.
- Kuroda, N. and Shiraki, K. (1975) Boninite and related rocks of Chichijima, Bonin Islands, Japan. *Reports, Faculty of Science, Shizuoka University*, 10, 145-155.
- Kuroda, N., Shiraki, K. and Urano, H. (1978) Boninite as a possible calc-alkalic primary magma. *Bulletin Volcanologique*, 41, 563-575.
- Kushiro, I. and Sato, J. (1978) Origin of some calc-alkaline andesites in Japanese Islands. *Bulletin Volcanologique*, 41, 575-585.
- Kushiro, I. and Takahashi, E. (1981) Upper mantle materials in

- the oceanic and island arc regions and origin of abyssal basalts. (abstr.) Bulletin of the Volcanological Society of Japan, 26, 130.
- Menzies, M. and Murthy, Y. R. (1980) Enriched mantle: Nd and Sr isotopes in diopside from kimberlite nodules. *Nature*, 283, 634–636.
- Nakamura, Yasuo (1971) Equilibrium relations in Mg-rich part of the pyroxene quadrilateral. *Mineralogical Journal*, 6, 264–276.
- Paris, J. P. (1981) *Geology de la Nouvelle-Calédonie: un essai de synthèse*. Memoire Bureau de Recherches Geologiques et Minière No. 113.
- Paris, J. P., Andreieff, P. and Coudray, J. (1979) Sur l'âge éocène supérieur de la mise en place de la nappe ophiolitique de la Nouvelle-Calédonie, unité du charriage océanique périaustralien, déduit d'observations nouvelles sur la série de Népoui. *Compte Rendus Academy Science Paris* 288 D, 1659–1661.
- Petersen, Jon (1891a) Beitrage zur Petrographie von Sulphur Island, Peel Island, Hachijo und Mijakeshima. *Jahrb. Hamburg. Wiss. Anst.*, 8, 1–59.
- Petersen, Jon (1891b) Der Boninit von Peel Island. *Jahrb. Hamburg. Wiss. Anst.*, 8, 341–349.
- Sadanaga, R. and Okamura, F. P. (1969) X-ray study of the phase transformation of enstatite. *Proceedings of the Japanese Academy*, 45, 721–726.
- Sarver, J. F. and Hummel, F. A. (1962) Stability relations of magnesium metasilicate polymorphs. *Journal of the American Ceramic Society*, 45, 152–156.
- Sharaskin, A. Y., Dobretsov, N. L. and Sobolev, N. V. (1979) Marianites: the clinoenstatite bearing pillow-lavas associated with the ophiolite assemblage of Mariana trench. In A. Panamiotou, Ed., *Ophiolites*, p. 473–479. *Proceedings International Ophiolite Symposium Cyprus 1979*. Ministry of Agriculture and Natural Resources, Cyprus.
- Shiraki, K. and Kuroda, N. (1977) The boninite revisited. *Journal of Geography (Japan)*, 86, 71–72.
- Shiraki, K., Kuroda, N. and Urano, H. (1979) Clinoenstatite-bearing boninite of Muko-jima, Bonin Islands (in Japanese). *Journal of the Geological Society of Japan*, 85, 591–594.
- Shiraki, K., Kuroda, N., Urano, H. and Maruyama, S. (1980) Clinoenstatite in boninites from the Bonin Islands, Japan. *Nature*, 285, 31–32.
- Suzuki, K. and Shiraki, K. (1980) Chromite-bearing spessartite from Kasuga-mura, Japan, and their bearing on possible mantle origin andesite. *Contributions to Mineralogy and Petrology*, 71, 313–322.
- Wass, S. Y. (1980) Geochemistry and origin of xenolith-bearing and related alkali basaltic rocks from the Southern Highlands, N.S.W. Australia. *American Journal of Science*, 280-A Part 2, 639–694.

*Manuscript received, May 24, 1982;
accepted for publication, April 18, 1983.*