

Davanite, $K_2TiSi_6O_{15}$, in the Smoky Butte (Montana) lamproites: Discussion of X-ray powder data

GUNNAR RAADE

Mineralogisk-Geologisk Museum, Universitetet i Oslo, Sars' gate 1, N-0562 Oslo 5, Norway

Davanite, the Ti analogue of dalyite ($K_2ZrSi_6O_{15}$), was described as a new mineral from Murunski, USSR, by Lazebnik et al. (1984). Wagner and Velde (1986) have reported the second find of davanite, from Smoky Butte,

Montana. X-ray powder data for the Montana davanite are listed by Wagner and Velde (1986) together with data for the Russian mineral taken from the original description.

TABLE 1. X-ray powder-diffraction data for davanite

Synthetic $K_2TiSi_6O_{15}^*$			Murunskit†		Smoky Butte‡		Synthetic $K_2TiSi_6O_{15}^*$			Murunskit†		Smoky Butte‡	
hkl	d_{calc}	I_{calc}	d_{obs}	I_{obs}	d_{obs}	I_{obs}	hkl	d_{calc}	I_{calc}	d_{obs}	I_{obs}	d_{obs}	I_{obs}
100	6.377	12			6.375	w	220	2.051	10	2.057	1	2.053	w
001	5.931	6					231	2.036	3				
101	5.868	34	5.88	5	5.878	m	130	1.976	3				
111	4.285	32	4.31	1	4.290	w	112	1.973	5				
110	4.102	53	4.09	5	4.104	ms	022	1.925	4	1.929	<1	1.974	w
111	3.873	5			3.878	vw	310	1.878	11	1.886§	1	1.880	w
011	3.849	4					330	1.860	3			1.866	vw
101	3.604	47			3.605	w	141	1.857	5				
021	3.594	5					231	1.833	4			1.833	w
120	3.491	96	3.51	10	3.490	s	113	1.807	4				
020	3.435	26			3.423§	w	202	1.802	10	1.799	1	1.802	w
112	3.344	47	3.34	5	3.349	s	321	1.800	4				
210	3.293	18			3.305	w	323	1.766	9			1.766	w
121	3.183	29					233	1.762	9	1.762§	1		
012	3.176	32	3.19	5	3.179	ms	142	1.747	4				
211	3.023	100	3.022	10 b	3.022	vs	232	1.735	3				
002	2.966	34			2.970	w	214	1.719	3				
022	2.807	26					133	1.712	5				
220	2.790	13					124	1.684	4			1.684	w
212	2.785	20	2.789§	7	2.790	m b	410	1.667	3				
111	2.777	10			2.706	vw	204	1.664	5				
120	2.705	3					403	1.663	15	1.667	3 b	1.668	w
212	2.619	60	2.615	9 b	2.623	m b	332	1.652	8				
210	2.610	9					122	1.645	4				
021	2.591	19			2.588	m b	323	1.641	3				
211	2.576	18					241	1.622	3				
112	2.516	5			2.514	vw	141	1.612	6			1.613	w
031	2.458	10					143	1.606	4				
221	2.432	3					320	1.606	9	1.604	3	1.607	w
122	2.410	4					304	1.605	5				
131	2.403	8					340	1.588	3				
201	2.392	7					431	1.569	5				
311	2.384	15	2.391	3	2.391	w	242	1.541	2				
221	2.341	3			2.340	vw	143	1.539	2	1.542§	<1		
131	2.198	5					422	1.511	4				
312	2.197	6	2.215§	2	2.201§	w	213	1.511	3			1.510§	w
230	2.197	6					332	1.509	3				
222	2.143	13			2.144	w	233	1.497	12	1.498	4	1.500	w
013	2.116	14	2.119	2 b	2.115	w	023	1.494	2				
121	2.109	4					423	1.494	3				
132	2.091	18					151	1.492	3			1.496§	w
231	2.086	3					432	1.492	5				
023	2.070	4			2.076§	vw				+16 lines to $d = 1.011 \text{ \AA}$			

Note: vs = very strong, s = strong, ms = medium strong, w = weak, b = broad.

* Pattern calculated with the program LAZY PULVERIX (Yvon et al., 1977) from structural data of Gebert et al. (1983). Cell dimensions: $a = 7.250(2)$, $b = 7.474(2)$, $c = 6.909(3) \text{ \AA}$, $\alpha = 105.59(5)$, $\beta = 112.81(5)$, $\gamma = 99.28(5)^\circ$. All reflections with $I_{\text{calc}} \geq 3$ and $d > 1.490 \text{ \AA}$ are listed. Diffraction geometry: Debye-Scherrer. $\lambda_{\text{CuK}\alpha} = 1.5418 \text{ \AA}$.

† Lazebnik et al. (1984).

‡ Wagner and Velde (1986).

§ Not used for calculation of cell parameters.

Wagner and Velde (1986) remarked as an "interesting fact" that the strongest reflection in the two patterns could not be indexed satisfactorily. Again, there are unexplained differences in the indexing of the two patterns. Wagner and Velde claimed that the cell dimensions of the two davanites are not identical, the a parameter differing by 0.11 Å. However, no explanation is offered for this discrepancy. The chemical analyses reported for the two davanites are virtually identical (Table 2 in Wagner and Velde, 1986), and no such difference in the cell dimensions would actually be expected.

These apparent inconsistencies are quite simply explained by incorrect indexing of the powder patterns. The crystal structure of synthetic $K_2TiSi_6O_{15}$ was published by Gebert et al. (1983). They have also tabulated observed and calculated X-ray powder data, but have neglected to include the intensities. Again, some of the stronger reflections have somehow been omitted. Based on the structural data of Gebert et al. (1983), a calculated pattern for $K_2TiSi_6O_{15}$ is presented here (Table 1). It was computed with the program LAZY PULVERIX (Yvon et al., 1977). With this information, the patterns of the Murunski and Smoky Butte davanites can be easily indexed (Table 1).

Least-squares refinements of the cell parameters of the Murunski mineral based on 16 reflections gave $a = 7.26(1)$, $b = 7.51(2)$, $c = 6.92(1)$ Å, $\alpha = 105.6(2)^\circ$, $\beta = 112.9(1)^\circ$, $\gamma = 99.4(2)^\circ$, $V = 319.0(7)$ Å³. The Smoky Butte mineral has nearly identical cell parameters, based on 33 reflections: $a = 7.272(3)$, $b = 7.480(4)$, $c = 6.910(2)$ Å, $\alpha = 105.55(3)^\circ$, $\beta = 112.82(3)^\circ$, $\gamma = 99.42(3)^\circ$, $V = 318.2(1)$ Å³.

REFERENCES

- Gebert, W., Medenbach, O., and Flörke, O.W. (1983) Darstellung und Kristallographie von $K_2TiSi_6O_{15}$ —Isotyp mit Dalyit $K_2ZrSi_6O_{15}$. Tschermaks Mineralogische und Petrographische Mitteilungen, 31, 69–79.
Lazebnik, K.A., Lazebnik, Yu.D., and Makhotko, V.F. (1984) Davanite $K_2TiSi_6O_{15}$ —A new alkali titanosilicate (in Russian). Zapiski Vsesoyuznogo Mineralogicheskogo Obshchestva, 113, 95–97.
Wagner, C., and Velde, D. (1986) Davanite, $K_2TiSi_6O_{15}$, in the Smoky Butte (Montana) lamproites. American Mineralogist, 71, 1473–1475.
Yvon, K., Jeitschko, W., and Parthé, E. (1977). LAZY PULVERIX, a computer program, for calculating X-ray and neutron diffraction powder patterns. Journal of Applied Crystallography, 10, 73–74.

MANUSCRIPT RECEIVED APRIL 16, 1987

MANUSCRIPT ACCEPTED MAY 18, 1987