

## Thermochemistry of pyroxenes on the join $Mg_2Si_2O_6$ - $CaMgSi_2O_6$ : A revision of the model for pressures up to 30 kbar

PAULA M. DAVIDSON,\* DONALD H. LINDSLEY

Department of Earth and Space Sciences, State University of New York, Stony Brook, New York 11794-2100, U.S.A.

WILLIAM D. CARLSON

Department of Geological Sciences, University of Texas at Austin, Austin, Texas 78713, U.S.A.

Carlson and Lindsley (1988) presented a model for the thermodynamic properties of the  $Mg_2Si_2O_6$ - $CaMgSi_2O_6$  join that describes most of the known subsolidus phase relations up to 1600 °C and 60 kbar. They recognized that devising a model to cover this wide range required the adoption of a number of compromises, none of which violates the explicit data available for that join. The preferred values for their model parameters generate an extraordinarily complex  $P$ - $T$  diagram, especially at pressures below 5 kbar (Fig. 1). Much of that complexity is illustrated by the 1-atm  $T$ - $X$  diagram (Fig. 2a). The two coincidences of Pig and Opx compositions (one stable at 1387 °C, the other metastable at 1303 °C for 1 atm) result from complex intersections of the Opx and Cpx free-energy curves. Complexities involving PEn result from the choice to make it ideal, in contrast to the strong non-idealities of Opx and Cpx. The very wide field for Opx at high temperatures results from the relatively low temperature for the inversion of Di to "orthodiopside." Another effect of that low temperature is the predicted breakdown of Pig to Aug + Opx with increasing temperature at 1389 °C. They were aware of these complexities, but similar features arose in all variations on the model that satisfactorily replicated the experimental data over the entire range of pressures from 1 atm to 60 kbar. Because the goal of their modeling was to provide a thermodynamic framework encompassing the full range of experimental data on the Fe-free join, the complexities appearing at low pressure and high temperature were accepted as an unavoidable compromise. Those complexities were regarded as inconsequential to the principal goal of their model, inasmuch as the complexities occur predominantly above the solidus and therefore would not be stable.

Davidson and Lindsley (1989) attempted to remodel quadrilateral pyroxene phase relations, using the Di-En join of Carlson and Lindsley (1988) as a starting point. Davidson and Lindsley's model was quite successful in describing the known phase equilibria and calorimetric data for the quadrilateral, with one serious exception. That quadrilateral model was unable to duplicate the lower thermal stability of Pig in the range  $0.1 < Fe/(Fe + Mg)$

$< 0.3$ ; the calculated diagram (Fig. 3) shows a hiatus in that range, with no stability for Pig at any temperature. However, pigeonites in that range are found in rocks and simple experimental systems (Lindsley, 1983, and references therein) and in experiments on naturally occurring compositions (e.g., Grove and Juster, 1987). Inspection of Figures 2a and 3 makes it clear that the pigeonite hiatus is an artifact of the high-temperature breakdown of Pig required for the Di-En join by the Carlson and Linds-




Fig. 1.  $P$ - $T$  projection for the system  $Mg_2Si_2O_6$ - $CaMgSi_2O_6$  based on the Carlson and Lindsley (1988) model. This is an extension of Fig. 2 of Carlson and Lindsley. Melting relations are omitted, but the melting interval is approximated by the stippled area. Note log  $P$  scale. End-member phases are indicated as OEn, orthoenstatite; CEn, clinoenstatite; PEn, protoenstatite; Di, (clino)diopside; ODi, "orthodiopside." Solid solutions are abbreviated as O, orthopyroxene, some highly calcic; Pig, low-Ca clinopyroxene; A, high-Ca clinopyroxene; Pr, protopyroxene. Abbreviations in parentheses label metastable curves.  $S_1$ ,  $S_2$ , and  $S_3$  are singular points.

\* Present address: Department of Geological Sciences, University of Illinois at Chicago, Chicago, Illinois 60680, U.S.A.




Fig. 2. Phase relations calculated at 1 atm from the Carlson and Lindsley (1988) model (a) and the revised solution model presented in this work (b). The melting interval is indicated by the stippled area in (a). Abbreviations as in Fig. 1. The Carlson and Lindsley model correctly predicts the re-appearance of orthopyroxene at near-solidus temperatures (Longhi and Boudreau, 1980; Carlson, 1988), but the relative compositions of O

and Pig are not correct. Note that the predicted inversion of Di to "orthodiopside" lies just above the liquidus. The revised model (b) lacks any field for high- $T$  orthopyroxene, but otherwise describes the available experimental data in a much simpler fashion than does the Carlson and Lindsley model. The revised model predicts that "orthodiopside" will not be stable below 1754 °C—comfortably above the liquidus.

ley (1988) model: with the addition of Fe, a peculiar feature that is "harmless" in the Fe-free system propagates into the subsolidus region with deleterious effect.

It is clear that the pigeonite hiatus stems from the low predicted temperature for the inversion of Di to "orthodiopside" in the Carlson and Lindsley (1988) model. Although their modeling had been structured to maximize this temperature, the high-pressure constraints required a barely supersolidus value for it in order to fit the experimental data at all pressures. It is therefore not possible to modify their model so as to eliminate the pigeonite hiatus in the quadrilateral and retain that model's ability to reproduce all the experimental data. We conclude that a different Di-En model—one that makes no attempt to fit a subset of the experimental data—must be used as a starting point for quadrilateral pyroxene models. This discussion presents that revised Di-En model.

For several reasons, we exclude the data at 40 kbar and above from the new modeling. Most data on quadrilateral pyroxenes have been obtained below 30 kbar, so it is important to use a Di-En model appropriate for lower pressures. In addition, most natural pyroxenes with  $\text{Fe}/(\text{Fe} + \text{Mg}) > 0.15$  have crystallized at pressures well be-




Fig. 3. Stability relations of Pig predicted by a quadrilateral solution model that incorporates the Carlson and Lindsley (1988) model for Di-En. Melting relations are not shown. The predicted gap in the stability of Pig at  $0.10 < \text{Fe}/(\text{Fe} + \text{Mg}) < 0.30$  is incompatible with a variety of experimental data and natural occurrences for which the minimum stability of Pig is indicated by the dashed line.  $T(\text{cons})$  indicates the intersection of the calculated  $\text{Pig} = \text{Opx} + \text{Aug}$  reaction with the  $\text{Pig}$ - $\text{Aug}$  consolute temperature.

**TABLE 1.** Values of revised thermochemical parameters compared with those of Carlson and Lindsley (CL)

Parameter	CL	This work
$\Delta U_A^0$	4.261	6.111
$\Delta V_A^0$	0.05900	0.09754
$\Delta S_A^0$	2.721	4.1323
$\Delta U_B^0$	-35.92	-39.060
$\Delta V_B^0$	-1.753	-0.1375
$\Delta S_B^0$	-20.97	-17.295
$\Delta U_E^0$	0.9814	1.1632
$\Delta V_E^0$	-0.07954	-0.08199
$\Delta S_E^0$	0.09328	0.16801
$\Delta U_F^0$	-36.71	-6.129
$\Delta V_F^0$	0.0	0.0
$\Delta S_F^0$	2.444	10.2181
$W_{O1}(Cpx)$	26.23	26.125
$W_{O2}(Cpx)$	32.44	32.301
$W_{O1}(Opx)$	-0.02229	-0.03843
$W_{O2}(Opx)$	-0.08646	-0.00670
$W_A(Opx)$	28.60	20.0*
$W_B(Opx)$	-1.749	0.0*
$W_A(Proto)$	0.0*	20.0*

Note: Energy in kJ/mol; volume in kJ/(kbar·mol); entropy in J/(K·mol). Subscripts A, B, E, and F refer to relations given in Carlson and Lindsley (1988).  
\* Adopted values.

low 30 kbar. After exclusion of the high-pressure data (and a set at 1500 °C, 30 kbar), our approach to the modeling was virtually identical to that of Carlson and Lindsley (1988). We used linear programming and included in the objective function terms to maximize  $\Delta H$  and minimize  $\Delta S$  for the Di → ODi inversion. This combination makes the calculated temperature for that inversion as high as feasible. By omitting the 40 to 60-kbar data, we were also able to eliminate  $W_{Opx}^{Opx}$ , which has the implausibly large value of -1.749 J/(bar·mol) in the Carlson and Lindsley (1988) model. In preliminary fitting, we found that both  $W_{Opx}^{Opx}$  and  $W_{Opx}^{PEn}$  could vary from zero to more than 30 kJ. To minimize the complex interactions found in their model, we chose to make both values equal and reasonably close to the parameters for Aug. Because  $W_{Opx}^{Opx}$  varies antipathetically with  $\Delta H$  for the Di → ODi inversion we adopted compromise values of 20 kJ for both  $W_{Opx}^{Opx}$  and  $W_{Opx}^{PEn}$ . Values for the fitted parameters are compared to those of Carlson and Lindsley in Table 1. The revised value for  $\Delta U_A^0$  - 6.111 kJ—is marginally closer to that extrapolated from calorimetric measurements (Newton et al., 1979). The value for  $\Delta V_B^0$  is very much smaller than that of Carlson and Lindsley (1988), probably because we were able to eliminate  $W_{Opx}^{PEn}$ , which is highly correlated with  $\Delta V_B^0$ . More than three quarters of the parameters for the revised model lie outside the range of

maximum and minimum values derived by Carlson and Lindsley, a striking expression of the influence of the high-pressure data used in that model and omitted in the revised model.

Elimination of the high-pressure constraints also permits several improvements in the fit to the low-pressure data. The revised Di-En model yields 1466 °C for the coexistence of Opx + Pig + Di at 15 kbar—much closer to the experimental value of  $1465 \pm 10$  °C (Schweitzer, 1982) than the 1436 °C predicted by Carlson and Lindsley (1988). Except for the absence of a high-temperature field for orthopyroxene, this model fits the 1-atm data as well as does that of Carlson and Lindsley, and it avoids many of the complexities of their model at that pressure (Fig. 2b). Most important, the revised model avoids the second Pig = Opx + Aug reaction that led to the pigeonite hiatus in the quadrilateral model based on Carlson and Lindsley (1988). As a basis for fitting relations within the quadrilateral at pressures up to 30 kbar, the revised model is clearly superior.

What the revised Di-En model does *not* do nearly so well as that of Carlson and Lindsley (1988) is to predict the phase relations at simultaneous high pressures and temperatures—hardly a surprise since we were compelled to ignore the high-pressure data in deriving it. For compositions close to the Di-En join—especially for higher pressures—the Carlson and Lindsley 1988 model remains the model of choice.

#### ACKNOWLEDGMENTS

We thank S. K. Saxena for reviewing this manuscript. Portions of this work were supported by NSF grant EAR 87-20185 to D. H. Lindsley.

#### REFERENCES CITED

- Carlson, W.D. (1988) Subsolidus phase equilibria on the forsterite-saturated join  $Mg_2Si_2O_6$ - $CaMgSi_2O_6$  at atmospheric pressure. *American Mineralogist*, 73, 232-241.
- Carlson, W.D., and Lindsley, D.H. (1988) Thermochemistry of pyroxenes on the join  $Mg_2Si_2O_6$ - $CaMgSi_2O_6$ . *American Mineralogist*, 73, 242-252.
- Davidson, P.M., and Lindsley, D.H. (1989) Thermodynamic analysis of pyroxene-olivine-quartz equilibria in the system CaO-MgO-FeO-SiO<sub>2</sub>. *American Mineralogist*, 74, in press.
- Grove, T.L., and Juster, T.C. (1987) Orthopyroxene and pigeonite stability relations at 1-atm in natural andesite and basalt systems. *Geological Society of America Abstracts with Programs*, 19, 685.
- Lindsley, D.H. (1983) Pyroxene thermometry. *American Mineralogist*, 68, 477-493.
- Longhi, J., and Boudreau, A.E. (1980) The orthoenstatite liquidus field in the system forsterite-diopside-silica. *American Mineralogist*, 65, 563-573.
- Newton, R.C., Charlu, P.A., Anderson, P.A.M., and Kleppa, O.J. (1979) Thermochemistry of synthetic clinopyroxenes in the join  $CaMgSi_2O_6$ - $Mg_2Si_2O_6$ . *Geochimica et Cosmochimica Acta*, 43, 55-60.
- Schweitzer, E. (1982) The reaction pigeonite = diopside<sub>ss</sub> + enstatite<sub>ss</sub> at 15 kbars. *American Mineralogist*, 67, 54-58.

MANUSCRIPT RECEIVED JANUARY 29, 1988

MANUSCRIPT ACCEPTED AUGUST 1, 1988