

Table 6. Atomic coordinates ($\text{\AA} \times 104$) and equivalent isotropic $U(\text{eq})$ and anisotropic displacement parameters ($\text{\AA}^2 \times 103$) for 120853 n.5. $U(\text{eq})$ is defined as one third of the trace of the orthogonalized U_{ij} tensor. The anisotropic displacement factor exponent takes the form: $-2 \pi^2 [h^2 a^{*2} U_{11} + \dots + 2 h k a^* b^* U_{12}]$

U13	x U12	y	z	U(eq)	U11	U22	U33	U23
O(11)	3631(1)	3392(1)	1203(2)	7(1)	5(1)	8(1)	8(1)	0(1)
1(1)	0(1)							
O(12)	3613(1)	1780(1)	6488(2)	7(1)	5(1)	8(1)	7(1)	0(1)
1(1)	0(1)							
O(21)	6149(1)	5099(1)	3090(2)	9(1)	10(1)	6(1)	9(1)	-1(1)
2(1)	-2(1)							
O(22)	6061(1)	9982(1)	8052(2)	10(1)	11(1)	7(1)	11(1)	1(1)
3(1)	2(1)							
O(31)	6066(1)	2668(1)	40(2)	7(1)	6(1)	9(1)	7(1)	-
2(1)	2(1)	0(1)						
O(32)	5976(1)	2403(1)	4970(2)	8(1)	7(1)	9(1)	7(1)	2(1)
2(1)	0(1)							
Si(1)	5392(1)	3479(1)	2265(1)	5(1)	4(1)	5(1)	6(1)	0(1)
1(1)	0(1)							
Si(2)	5372(1)	1628(1)	7305(1)	5(1)	4(1)	5(1)	6(1)	0(1)
1(1)	0(1)							
M11	2500	3477(1)	7500	6(1)	5(1)	5(1)	7(1)	0
1(1)	0							
M1	2500	1597(1)	2500	7(1)	6(1)	7(1)	7(1)	0
1(1)	0							
M2	2500	5520(1)	2500	10(1)	11(1)	8(1)	8(1)	0
0(1)	0							
M21	2500	9506(1)	7500	10(1)	13(1)	6(1)	9(1)	0
-2(1)	0							