

Comparative planetary mineralogy: Pyroxene major- and minor-element chemistry and partitioning of vanadium between pyroxene and melt in planetary basalts

JIM KARNER,* J.J. PAPIKE, AND C.K. SHEARER

Institute of Meteoritics, Department of Earth and Planetary Sciences, University of New Mexico, Albuquerque, New Mexico 87131-1126, U.S.A.

DEPOSIT ART INFORMATION (4 pages)




FIGURE A-1. Fe³⁺ vs. X_{Fe} for all of the terrestrial and martian pyroxene. See Figure A-2 caption for explanation of abbreviations.




FIGURE A-2. Na vs. X_{Fe} variation diagram for pyroxene grains from all of the planetary basalt samples. KEW=Keweenawan, IA=Island Arc, CP=Columbia Plateau, HAW=Hawaiian, and TP=Taos Plateau. Lunar samples are listed by thin section number. Martian samples are DaG476, NWA1195, NWA1110, Shergotty and Yamato 980459 (Y98). The sample from 4 Vesta is Pasamonte.




FIGURE A-3. Cr vs. X_{Fe} for pyroxene grains from all of the planetary basalt samples.




FIGURE A-4. Mn vs. Fe²⁺ in atoms per formula unit (afu) for pyroxene analyses from each thin section from the different basalt suites.




FIGURE A-5. V pyroxene/V rock (V*) vs. X_{Fe} for pyroxene grains from select basalt suites from the four different planets.




FIGURE A-6. V* vs. Ca (afu) for the same pyroxene analyses as in Figure A-5.




FIGURE A-7. Ti vs. X_{Fe} for pyroxene grains from all the planetary basalt samples.




FIGURE A-8. Al vs. X_{Fe} for pyroxene grains from all the planetary basalt samples.