

data\_vm\_x1\_190k\_abs

\_audit\_creation\_method SHELXL-97  
\_chemical\_name\_systematic  
;  
?  
;  
\_chemical\_name\_common ?  
\_chemical\_melting\_point ?  
\_chemical\_formula\_moiety ?  
\_chemical\_formula\_sum  
'H B Ca O5 Si'  
\_chemical\_formula\_weight 159.99

loop\_  
\_atom\_type\_symbol  
\_atom\_type\_description  
\_atom\_type\_scatter\_dispersion\_real  
\_atom\_type\_scatter\_dispersion\_imag  
\_atom\_type\_scatter\_source  
'H' 'H' 0.0000 0.0000  
'International Tables Vol C Tables 4.2.6.8 and 6.1.1.4'  
'B' 'B' 0.0013 0.0007  
'International Tables Vol C Tables 4.2.6.8 and 6.1.1.4'  
'O' 'O' 0.0106 0.0060  
'International Tables Vol C Tables 4.2.6.8 and 6.1.1.4'  
'Si' 'Si' 0.0817 0.0704  
'International Tables Vol C Tables 4.2.6.8 and 6.1.1.4'  
'Ca' 'Ca' 0.2262 0.3064  
'International Tables Vol C Tables 4.2.6.8 and 6.1.1.4'

\_symmetry\_cell\_setting ?  
\_symmetry\_space\_group\_name\_H-M ?

loop\_  
\_symmetry\_equiv\_pos\_as\_xyz  
'x, y, z'  
'-x, y+1/2, -z+1/2'  
'-x, -y, -z'  
'x, -y-1/2, z-1/2'

\_cell\_length\_a 4.83080(10)  
\_cell\_length\_b 7.60110(10)  
\_cell\_length\_c 9.6290(2)  
\_cell\_angle\_alpha 90.00  
\_cell\_angle\_beta 90.176(2)  
\_cell\_angle\_gamma 90.00  
\_cell\_volume 353.569(11)  
\_cell\_formula\_units\_Z 4  
\_cell\_measurement\_temperature 463(2)  
\_cell\_measurement\_reflns\_used ?  
\_cell\_measurement\_theta\_min ?

```

_cell_measurement_theta_max    ?

_exptl_crystal_description     ?
_exptl_crystal_colour         ?
_exptl_crystal_size_max       ?
_exptl_crystal_size_mid       ?
_exptl_crystal_size_min       ?
_exptl_crystal_density_meas    ?
_exptl_crystal_density_diffn   3.006
_exptl_crystal_density_method  'not measured'
_exptl_crystal_F_000           320
_exptl_absorpt_coefficient_mu   2.002
_exptl_absorpt_correction_type ?
_exptl_absorpt_correction_T_min ?
_exptl_absorpt_correction_T_max ?
_exptl_absorpt_process_details ?

_exptl_special_details
;
?
;

_diffrn_ambient_temperature    190(2)
_diffrn_radiation_wavelength   0.71073
_diffrn_radiation_type         MoK\alpha
_diffrn_radiation_source       'fine-focus sealed tube'
_diffrn_radiation_monochromator graphite
_diffrn_measurement_device_type ?
_diffrn_measurement_method     ?
_diffrn_detector_area_resol_mean ?
_diffrn_standards_number       ?
_diffrn_standards_interval_count ?
_diffrn_standards_interval_time ?
_diffrn_standards_decay_%      ?
_diffrn_reflns_number          8520
_diffrn_reflns_av_R_equivalents 0.0315
_diffrn_reflns_av_sigmaI/netI  0.0247
_diffrn_reflns_limit_h_min     -7
_diffrn_reflns_limit_h_max     8
_diffrn_reflns_limit_k_min     -13
_diffrn_reflns_limit_k_max     13
_diffrn_reflns_limit_l_min     -16
_diffrn_reflns_limit_l_max     17
_diffrn_reflns_theta_min       4.22
_diffrn_reflns_theta_max       40.10
_reflns_number_total           2200
_reflns_number_gt              1942
_reflns_threshold_expression    >2sigma(I)

_computing_data_collection     ?
_computing_cell_refinement     ?
_computing_data_reduction      ?
_computing_structure_solution   ?
_computing_structure_refinement 'SHELXL-97 (Sheldrick, 1997)'

```

\_computing\_molecular\_graphics ?  
\_computing\_publication\_material ?

\_refine\_special\_details

;

Refinement of  $F^2$  against ALL reflections. The weighted R-factor wR and goodness of fit S are based on  $F^2$ , conventional R-factors R are based on F, with F set to zero for negative  $F^2$ . The threshold expression of  $F^2 > 2\sigma(F^2)$  is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on  $F^2$  are statistically about twice as large as those based on F, and R-factors based on ALL data will be even larger.

;

\_refine\_ls\_structure\_factor\_coef Fsqd  
\_refine\_ls\_matrix\_type full  
\_refine\_ls\_weighting\_scheme calc  
\_refine\_ls\_weighting\_details  
'calc w=1/[\s^2^(Fo^2^)+(0.0100P)^2^+0.0000P] where P=(Fo^2^+2Fc^2^)/3'  
\_atom\_sites\_solution\_primary direct  
\_atom\_sites\_solution\_secondary difmap  
\_atom\_sites\_solution\_hydrogens geom  
\_refine\_ls\_hydrogen\_treatment mixed  
\_refine\_ls\_extinction\_method SHELXL  
\_refine\_ls\_extinction\_coef 0.035(2)  
\_refine\_ls\_extinction\_expression  
'Fc^\*=kFc[1+0.001xFc^2^/l^3^/sin(2\q)]^-1/4^'  
\_refine\_ls\_number\_reflns 2200  
\_refine\_ls\_number\_parameters 78  
\_refine\_ls\_number\_restraints 0  
\_refine\_ls\_R\_factor\_all 0.0262  
\_refine\_ls\_R\_factor\_gt 0.0212  
\_refine\_ls\_wR\_factor\_ref 0.0444  
\_refine\_ls\_wR\_factor\_gt 0.0439  
\_refine\_ls\_goodness\_of\_fit\_ref 1.457  
\_refine\_ls\_restrained\_S\_all 1.457  
\_refine\_ls\_shift/su\_max 0.001  
\_refine\_ls\_shift/su\_mean 0.000

loop\_

\_atom\_site\_label  
\_atom\_site\_type\_symbol  
\_atom\_site\_fract\_x  
\_atom\_site\_fract\_y  
\_atom\_site\_fract\_z  
\_atom\_site\_U\_iso\_or\_equiv  
\_atom\_site\_adp\_type  
\_atom\_site\_occupancy  
\_atom\_site\_symmetry\_multiplicity  
\_atom\_site\_calc\_flag  
\_atom\_site\_refinement\_flags  
\_atom\_site\_disorder\_assembly  
\_atom\_site\_disorder\_group

Ca Ca 0.99179(3) 0.106263(19) 0.336047(16) 0.00445(4) Uani 1 1 d . . .

Si Si 0.46940(4) 0.26645(3) 0.08429(2) 0.00310(5) Uani 1 1 d . . .  
B B 0.56919(18) 0.41076(11) 0.34073(9) 0.00410(13) Uani 1 1 d . . .  
O1 O 0.24095(12) 0.40161(7) 0.03728(6) 0.00537(10) Uani 1 1 d . . .  
O2 O 0.67107(12) 0.30114(7) 0.45756(6) 0.00477(9) Uani 1 1 d . . .  
O3 O 0.67594(12) 0.33462(7) 0.21057(6) 0.00464(9) Uani 1 1 d . . .  
O4 O 0.31449(12) 0.08821(7) 0.14588(6) 0.00514(10) Uani 1 1 d . . .  
O5 O 0.25875(12) 0.41335(8) 0.33644(7) 0.00545(10) Uani 1 1 d . . .  
H H 0.203(3) 0.4463(18) 0.4099(17) 0.024(4) Uiso 1 1 d . . .

loop\_

\_atom\_site\_aniso\_label  
\_atom\_site\_aniso\_U\_11  
\_atom\_site\_aniso\_U\_22  
\_atom\_site\_aniso\_U\_33  
\_atom\_site\_aniso\_U\_23  
\_atom\_site\_aniso\_U\_13  
\_atom\_site\_aniso\_U\_12

Ca 0.00466(6) 0.00485(6) 0.00385(6) -0.00010(4) 0.00033(4) -0.00031(4)  
Si 0.00375(9) 0.00321(8) 0.00234(9) -0.00017(6) 0.00009(7) 0.00020(6)  
B 0.0044(3) 0.0042(3) 0.0037(3) 0.0001(2) 0.0001(2) -0.0001(2)  
O1 0.0059(2) 0.0053(2) 0.0049(2) -0.00026(17) -0.00043(18) 0.00156(17)  
O2 0.0053(2) 0.0055(2) 0.0035(2) 0.00147(17) 0.00061(17) 0.00046(17)  
O3 0.0051(2) 0.0060(2) 0.0028(2) -0.00111(17) 0.00009(17) 0.00003(17)  
O4 0.0058(2) 0.0037(2) 0.0059(2) 0.00063(17) 0.00179(18) 0.00009(16)  
O5 0.0041(2) 0.0076(2) 0.0047(2) -0.00088(18) 0.00026(18) 0.00055(17)

\_geom\_special\_details

;

All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

;

loop\_

\_geom\_bond\_atom\_site\_label\_1  
\_geom\_bond\_atom\_site\_label\_2  
\_geom\_bond\_distance  
\_geom\_bond\_site\_symmetry\_2  
\_geom\_bond\_publ\_flag  
Ca O1 2.2760(6) 2\_645 ?  
Ca O1 2.2788(6) 4\_666 ?  
Ca O4 2.4124(6) 1\_655 ?  
Ca O2 2.4444(6) . ?  
Ca O5 2.5221(6) 2\_645 ?  
Ca O3 2.6056(6) . ?  
Ca O3 2.6543(6) 2\_745 ?  
Ca O5 2.6668(6) 1\_655 ?  
Si O1 1.5733(6) . ?  
Si O2 1.6461(6) 4\_565 ?  
Si O3 1.6538(6) . ?  
Si O4 1.6582(6) . ?

B O4 1.4668(10) 2\_655 ?  
B O3 1.4750(11) . ?  
B O2 1.4827(10) . ?  
B O5 1.5002(10) . ?  
O1 Ca 2.2760(6) 2\_655 ?  
O1 Ca 2.2788(6) 4\_465 ?  
O2 Si 1.6461(6) 4\_566 ?  
O3 Ca 2.6543(6) 2\_755 ?  
O4 B 1.4668(10) 2\_645 ?  
O4 Ca 2.4124(6) 1\_455 ?  
O5 Ca 2.5221(6) 2\_655 ?  
O5 Ca 2.6668(6) 1\_455 ?

loop\_

\_geom\_angle\_atom\_site\_label\_1  
\_geom\_angle\_atom\_site\_label\_2  
\_geom\_angle\_atom\_site\_label\_3  
\_geom\_angle  
\_geom\_angle\_site\_symmetry\_1  
\_geom\_angle\_site\_symmetry\_3  
\_geom\_angle\_publ\_flag  
O1 Ca O1 77.69(2) 2\_645 4\_666 ?  
O1 Ca O4 133.58(2) 2\_645 1\_655 ?  
O1 Ca O4 107.64(2) 4\_666 1\_655 ?  
O1 Ca O2 80.96(2) 2\_645 . ?  
O1 Ca O2 86.72(2) 4\_666 . ?  
O4 Ca O2 144.08(2) 1\_655 . ?  
O1 Ca O5 73.66(2) 2\_645 2\_645 ?  
O1 Ca O5 142.73(2) 4\_666 2\_645 ?  
O4 Ca O5 77.11(2) 1\_655 2\_645 ?  
O2 Ca O5 111.34(2) . 2\_645 ?  
O1 Ca O3 114.48(2) 2\_645 . ?  
O1 Ca O3 135.96(2) 4\_666 . ?  
O4 Ca O3 93.713(19) 1\_655 . ?  
O2 Ca O3 56.437(19) . . ?  
O5 Ca O3 78.629(19) 2\_645 . ?  
O1 Ca O3 81.89(2) 2\_645 2\_745 ?  
O1 Ca O3 78.771(19) 4\_666 2\_745 ?  
O4 Ca O3 55.589(18) 1\_655 2\_745 ?  
O2 Ca O3 159.53(2) . 2\_745 ?  
O5 Ca O3 74.074(18) 2\_645 2\_745 ?  
O3 Ca O3 142.497(9) . 2\_745 ?  
O1 Ca O5 146.84(2) 2\_645 1\_655 ?  
O1 Ca O5 76.562(19) 4\_666 1\_655 ?  
O4 Ca O5 74.765(19) 1\_655 1\_655 ?  
O2 Ca O5 77.064(19) . 1\_655 ?  
O5 Ca O5 137.809(13) 2\_645 1\_655 ?  
O3 Ca O5 72.546(18) . 1\_655 ?  
O3 Ca O5 112.855(18) 2\_745 1\_655 ?  
O1 Si O2 114.00(3) . 4\_565 ?  
O1 Si O3 115.34(3) . . ?  
O2 Si O3 106.60(3) 4\_565 . ?  
O1 Si O4 108.63(3) . . ?  
O2 Si O4 106.19(3) 4\_565 . ?

O3 Si O4 105.37(3) . . ?  
O4 B O3 107.49(7) 2\_655 . ?  
O4 B O2 108.90(6) 2\_655 . ?  
O3 B O2 107.94(6) . . ?  
O4 B O5 111.90(6) 2\_655 . ?  
O3 B O5 109.50(6) . . ?  
O2 B O5 110.96(7) . . ?  
Si O1 Ca 129.74(3) . 2\_655 ?  
Si O1 Ca 126.56(3) . 4\_465 ?  
Ca O1 Ca 102.31(2) 2\_655 4\_465 ?  
B O2 Si 122.82(5) . 4\_566 ?  
B O2 Ca 100.76(5) . . ?  
Si O2 Ca 122.90(3) 4\_566 . ?  
B O3 Si 122.41(5) . . ?  
B O3 Ca 94.19(4) . . ?  
Si O3 Ca 118.90(3) . . ?  
B O3 Ca 92.96(4) . 2\_755 ?  
Si O3 Ca 118.90(3) . 2\_755 ?  
Ca O3 Ca 104.068(19) . 2\_755 ?  
B O4 Si 127.56(5) 2\_645 . ?  
B O4 Ca 103.53(5) 2\_645 1\_455 ?  
Si O4 Ca 121.22(3) . 1\_455 ?  
B O5 Ca 120.23(5) . 2\_655 ?  
B O5 Ca 118.16(4) . 1\_455 ?  
Ca O5 Ca 106.07(2) 2\_655 1\_455 ?

\_diffn\_measured\_fraction\_theta\_max 0.994  
\_diffn\_reflns\_theta\_full 40.00  
\_diffn\_measured\_fraction\_theta\_full 0.994  
\_refine\_diff\_density\_max 0.541  
\_refine\_diff\_density\_min -0.358  
\_refine\_diff\_density\_rms 0.106