

WHAT LURKS IN THE MARTIAN ROCKS AND SOIL? INVESTIGATIONS OF SULFATES, PHOSPHATES, AND PERCHLORATES
Stability and spectroscopy of Mg sulfate minerals: Role of hydration on sulfur isotope partitioning†

EMA BOBOCIOIU^{1,*} AND RAZVAN CARACAS¹

¹Laboratoire de Géologie de Lyon (LGLTPE) CNRS UMR 5276, Université Claude Bernard Lyon 1, Ecole Normale Supérieure de Lyon 46, allée d'Italie, 69364 Lyon, France

ABSTRACT

We study hydrated magnesium sulfate minerals from first-principles calculations based on density-functional theory. We determine the heat of hydration for $\text{MgSO}_4 \cdot n\text{H}_2\text{O}$, compute the Raman and infrared spectra for several phases and calculate the S isotope partitioning as a function of hydration. We find that epsomite and meridianiite with, respectively, $n = 7$ and $n = 11$ water molecules per MgSO_4 unit are particularly stable with respect to other individual or combinations of hydration states. The Raman spectra of all phases present clear SO_4 features that are easily identifiable. We use this to show one can use the vibrational spectroscopic information as an identification tool in a remote environment, like the martian surface. We discuss the character and atomic displacement pattern of all vibration modes and compute the $^{34}\text{S}/^{32}\text{S}$ partitioning; this work shows that hydration favors enrichment in the lighter S isotope ^{32}S with respect to the heavier ^{34}S , which is accumulated in the less hydrous structures. We show for the first time that the signature of $^{34}\text{S}/^{32}\text{S}$ partitioning could be observed by in situ spectroscopy on the surface of Mars. Moreover this can be related to the diurnal cycle of hydration and dehydration and hence it can improve the modeling of the water circulation on Mars.

Keywords: MgSO_4 , hydration, S isotope partitioning, density-functional theory