

## **Trace and minor elements in galena: A reconnaissance LA-ICP-MS study**

**LUKE GEORGE<sup>1</sup>, NIGEL J. COOK<sup>1</sup>, CRISTIANA L. CIOBANU<sup>1,\*</sup> AND BENJAMIN P. WADE<sup>2</sup>**

<sup>1</sup>Centre for Tectonics, Resources and Exploration (TRaX), School of Earth and Environmental Sciences, University of Adelaide, North Terrace, South Australia 5005, Australia

<sup>2</sup>Adelaide Microscopy, University of Adelaide, North Terrace, South Australia 5005, Australia

### **ABSTRACT**

Minor and trace elements can substitute into the crystal lattice of galena at various concentrations. In situ LA-ICP-MS analysis and trace element mapping of a range of galena specimens from different deposit types are used to obtain minor/trace element data, aimed at achieving insight into factors that control minor/trace element partitioning. The previously recognized coupled substitution  $\text{Ag}^{+}+(\text{Bi,Sb})^{3+} \leftrightarrow 2\text{Pb}^{2+}$  is confirmed. However, the poorer correlation between Ag and (Bi+Sb) when the latter elements are present at high concentrations ( $\sim 2000$  ppm), suggests that site vacancies may come into play:  $[2(\text{Bi,Sb})^{3+} + \square \leftrightarrow 3\text{Pb}^{2+}]$ . Galena is the primary host of Tl in all mapped mineral assemblages. Along with Cu, Tl is likely incorporated into galena via the coupled substitution:  $(\text{Ag,Cu,Tl})^{+}+(\text{Bi,Sb})^{3+} \leftrightarrow 2\text{Pb}^{2+}$ . Tin can reach significant concentrations in galena ( $>500$  ppm). Cd and minor Hg can be incorporated into galena; the simple isovalent substitution  $(\text{Cd,Hg})^{2+} \leftrightarrow \text{Pb}^{2+}$  is inferred. This paper shows for the first time, oscillatory and sector compositional zoning of minor/trace elements (Ag, Sb, Bi, Se, Te, Tl) in galena from two epithermal ores. Zoning is attributed to slow crystal growth into open spaces within the vein at relatively low temperatures.

The present data show that galena can host a broader range of elements than previously recognized. For many measured elements, the data sets generated display predictable partitioning patterns between galena and coexisting minerals, which may be dependent on temperature or other factors. Trace element concentrations in galena and their grain-scale distributions may also have potential in the identification of spatial and/or temporal trends within individual metallogenic belts, and as markers of ore formation processes in deposits that have undergone superimposed metamorphism and deformation. Galena trace element geochemistry may also display potential to be used as a trace/minor element vector approach in mineral exploration, notably for recognition of proximal-to-distal trends within a given ore system.

**Keywords:** Galena, trace elements, laser ablation-inductively coupled plasma-mass spectrometry, compositional zoning, substitution mechanisms