

Carbon speciation in silicate-C-O-H melt and fluid as a function of redox conditions: An experimental study, in situ to 1.7 GPa and 900 °C

BJORN MYSEN^{1,*}

¹Geophysical Laboratory, Carnegie Institution of Washington, 5251 Broad Branch Road NW, Washington, D.C. 20015, U.S.A.

ABSTRACT

Carbon speciation in and partitioning among silicate-saturated C-O-H fluids and (C-O-H)-saturated melts have been determined ~1.7 GPa and 900 °C under reducing and oxidizing conditions. The measurements were conducted in situ while the samples were at the conditions of interest. The solution equilibria were (1) $2\text{CH}_4 + \text{Q}^n = 2\text{CH}_3 + \text{H}_2\text{O} + \text{Q}^{n+1}$ and (2) $2\text{CO}_3^{2-} + \text{H}_2\text{O} + 2\text{Q}^{n+1} = \text{HCO}_3^- + 2\text{Q}^n$, under reducing and oxidizing conditions, and where the superscript, n , in the Q^n -species denotes number of bridging oxygen in the silicate species (Q-species). The abundance ratios, CH_3/CH_4 and $\text{HCO}_3^-/\text{CO}_3^{2-}$, increase with temperature. The enthalpy change associated with the species transformation differs for fluids and melts and also for oxidized and reduced carbon [Reducing: $\Delta H_{(1)}^{\text{fluid}} = 16 \pm 5$ kJ/mol, $\Delta H_{(1)}^{\text{melt}} = 50 \pm 5$ kJ/mol; oxidizing $\Delta H_{(2)}^{\text{fluid}} = 81 \pm 14$ kJ/mol]. For the exchange equilibrium of CH_4 and CH_3 species between fluid and melt, the temperature-dependent equilibrium constant, $(X_{\text{CH}_4}/X_{\text{CH}_3})^{\text{fluid}}/(X_{\text{CH}_4}/X_{\text{CH}_3})^{\text{melt}}$, yields $\Delta H = 34 \pm 3$ kJ/mol.

Increased abundance ratios, CH_4/CH_3 and $\text{HCO}_3^-/\text{CO}_3^{2-}$, lead to increased polymerization of silicate+(C-O-H) melt. Because of such relations, melt transport properties (e.g., viscosity) and element partition coefficients between magmatic liquids, C-O-H fluids, and crystalline phases can vary by more than 100% with speciation changes of C-bearing volatiles upper mantle. These structure effects are more pronounced the higher the pressure and the more mafic the magma.

Keywords: Redox, COH volatiles, melt structure, melt properties, fluid structure