

## Fayalite oxidation processes in Obsidian Cliffs rhyolite flow, Oregon

AUDREY M. MARTIN<sup>1,2,\*</sup>, ETIENNE MÉDARD<sup>3</sup>, BERTRAND DEVOURARD<sup>3,4</sup>, LINDSAY P. KELLER<sup>1</sup>, KEVIN RIGHTER<sup>1</sup>, JEAN-LUC DEVIDAL<sup>3</sup> AND ZIA RAHMAN<sup>1</sup>

<sup>1</sup>NASA Johnson Space Center, Mailcode KT, 2101 NASA Parkway, Houston, Texas 77058, U.S.A.

<sup>2</sup>Earth, Environmental and Planetary Sciences, Case Western Reserve University, 10900 Euclid Avenue, Cleveland, Ohio 44118, U.S.A.

<sup>3</sup>Laboratoire Magmas et Volcans, Université Blaise Pascal-CNRS-IRD, OPGC, 5 rue Kessler, 63038 Clermont-Ferrand, France

<sup>4</sup>CEREGE, AMU-CNRS-IRD (UM34), Technopôle Arbois-Méditerranée, BP 80, 13545 Aix-en-Provence, France

### ABSTRACT

This study investigates the oxidation of fayalite  $\text{Fe}_2^{2+}\text{SiO}_4$  that is present in lithophysae from a rhyolite flow (Obsidian Cliffs, Oregon). Textural, chemical, and structural analyses of the successive oxidation zones are used to constrain: (1) the oxidation processes of olivine, and (2) the role of temperature, chemical diffusion, and meteoric infiltration. Petrologic analyses and thermodynamic modeling show that the rhyolite flow emplaced at 800–950 °C. Fayalite-bearing lithophysae formed only in the core of the lava flow. Variations in the gas composition inside the lithophysae induced the oxidation of fayalite to a laihunite-1M zone  $\text{Fe}_1^{2+}\text{Fe}_2^{3+}\square_1(\text{SiO}_4)_2$ . This zone is made of nano-lamellae of amorphous silica  $\text{SiO}_2$  and laihunite-3M  $\text{Fe}_{1.6}^{2+}\text{Fe}_{1.6}^{3+}\square_{0.8}(\text{SiO}_4)_2 + \text{hematite Fe}_2\text{O}_3$ . It probably formed by a nucleation and growth process in the fayalite fractures and defects and at fayalite crystal edges. The laihunite-1M zone then oxidized into an “oxyfayalite” zone with the composition  $\text{Fe}_{0.52}^{2+}\text{Fe}_{2.32}^{3+}\square_{1.16}(\text{SiO}_4)_2$ . This second oxidation zone is made of lamellae of amorphous silica  $\text{SiO}_2$  and hematite  $\text{Fe}_2\text{O}_3$ , with a possible small amount of ferrosilite  $\text{Fe}^{2+}\text{SiO}_3$ . A third and outer zone, composed exclusively of hematite, is also present. The successive oxidation zones suggest that there may be a mineral in the olivine group with higher  $\text{Fe}^{3+}$  content than laihunite-1M. The transformation of laihunite-1M to this “oxyfayalite” phase could occur by a reaction such as



This would imply that  $\text{Fe}^{3+}$  can also be incorporated in the M1 site of olivine.

**Keywords:** Olivine, fayalite, laihunite, oxyfayalite, rhyolite, lithophysae, oxidation