

Dissolved Cl, oxygen fugacity, and their effects on Fe behavior in a hydrous rhyodacitic melt

AARON S. BELL^{1,*} AND JAMES D. WEBSTER²

¹Institute of Meteoritics, University of New Mexico, Albuquerque, New Mexico 87109, U.S.A.

²Department of Earth and Planetary Science, American Museum of Natural History, New York, New York 10024, U.S.A.

ABSTRACT

We have conducted a series of experiments to evaluate the intrinsic effects of dissolved chlorine on $\text{Fe}^{3+}/\Sigma\text{Fe}$ and magnetite solubility in hydrous chloride-rich rhyodacitic liquids. The addition of Cl to the melt appears to have two prominent effects on iron in the melt: (1) dissolved Cl appears to perturb the magnetite-melt equilibrium, such that greater $\text{FeO}^{\text{total}}$ contents are required to support magnetite saturation in Cl-bearing melts than in Cl-free melts of equivalent bulk compositions; and (2) a systematic and progressive decrease of the measured $\text{Fe}^{3+}/\Sigma\text{Fe}$ as f_{O_2} is increased. These two intimately related effects each have important implications for redox processes occurring in Cl-enriched arc magmas.

Keywords: Iron, oxygen fugacity, chlorine, XANES