

OUTLOOKS IN EARTH AND PLANETARY MATERIALS

Normal to inverse transition in martian spinel: Understanding the interplay between chromium, vanadium, and iron valence state partitioning through a crystal-chemical lens

JAMES J. PAPIKE¹, PAUL V. BURGER¹, AARON S. BELL¹, CHARLES, K. SHEARER¹, LOAN LE² AND JOHN JONES³

¹Institute of Meteoritics, Department of Earth and Planetary Sciences, University of New Mexico, Albuquerque, New Mexico 87131, U.S.A.

²JSC Engineering, Technology and Science (JETS), NASA Johnson Space Center, Houston, Texas 77058, U.S.A.

³NASA Johnson Space Center, Houston, Texas 77058, U.S.A.

ABSTRACT



Spinel is a very important rock-forming mineral that is found in basalts from Earth, Mars, the Earth's Moon, and basaltic meteorites. Spinel can be used as a sensitive indicator of petrologic and geochemical processes that occur in its host rock. This paper highlights the role of increasing f_{O_2} (from IW-1 to FMQ+2) in converting a >90% normal spinel to an ~25% magnetite (inverse) spinel, the trajectory of $D_{\text{V}}^{\text{spinel/melt}}$ as it relates to the ratio of $\text{V}^{3+}/\text{V}^{4+}$ in the melt, and the crystal chemical attributes of the spinel that control the intrinsic compatibility of both V^{3+} and V^{4+} . This work examines the nuances of the V partitioning and provides a crystal chemical basis for understanding Fe³⁺, Cr, and V substitution into the octahedral sites of spinel. Understanding this interplay is critical for using spinels as both indicators of planetary parentage and reconstructing the redox history of magmatic systems on the terrestrial planets. Three potential examples for this use are provided. In addition, this work helps explain the ubiquitous miscibility gap between spinels with changing ulvospinel contents.

Keywords: Spinel, crystal chemistry, partitioning, vanadium, chromium, Invited Centennial article