

CHEMISTRY AND MINERALOGY OF EARTH'S MANTLE

Ca-Al-silicate inclusions in natural moissanite (SiC)

SIMONPIETRO DI PIERRO^{1,*} AND EDWIN GNOS²

¹UMR CNRS 5570, Laboratoire de Sciences de la Terre, ENS Lyon, France

²Natural History Museum of Geneva, route de Malagnou 1, CP 6434, 1211 Geneve 6, Switzerland

ABSTRACT

Hundred-micrometer-sized calcium-aluminum-silicates (CAS) inclusions occur in moissanite-4H, moissanite-15R, and moissanite-6H from Turkey. These inclusions commonly consist of tabular exsolution lamellae of two different minerals. By combined electron microprobe and Raman spectroscopy analysis, at least eight different, essentially Mg- and Fe-free Ca-Al-silicate or Al-silicate phases have been discerned. The most common phase is dmisteinbergite, a hexagonal modification of $\text{CaAl}_2\text{Si}_2\text{O}_8$, occurring in association with lamellae of $\text{Ca}_x(\text{Al,Si})_{1-x}\text{O}_3$ or $\text{Ca}_{1-x}(\text{Al,Si})_{2+x}\text{O}_5$ compositions. All three phases contain significant amounts of BaO (up to 2 mol% of celsiane component in dmisteinbergite), SrO, SO_3 , and light rare earth elements (LREE). In particular, $\text{Ca}_{1-x}(\text{Al,Si})_{2+x}\text{O}_5$ contains up to 2.1 wt% of LREE, 3.9 wt% of F, and significant traces of Cl, while it is also associated to osbornite (TiN). Pure ghlenite, $\text{Ca}_2\text{Al}_2\text{SiO}_7$, and three additional compositions, namely $\text{CaAl}_{4-x}\text{Si}_x\text{O}_7$, $\text{Ca}_{1-x}(\text{Al,Si})_{3+x}\text{O}_6$, and $\text{Ca}_{3-x}(\text{Al,Si})_{6+x}\text{O}_{14}$ have been found, either occurring as single grains or forming exsolution lamellae. They also contain significant amounts of BaO, SrO, SO_3 , and LREE. One last intriguing phase is composed in average of 65.9 wt% SiO_2 , 17.4% Al_2O_3 , 3.0% alkalis, 6.0% BaO, 2.0% $\text{CaO}+\text{MgO}$, 0.9% ZrO_2 , and up to 0.5% LREE. Dmisteinbergite and ghlenite show Raman peaks in very good agreement with literature data, $\text{Ca}_x(\text{Al,Si})_{1-x}\text{O}_3$ shows main Raman modes at 416 and 1009 cm^{-1} , $\text{Ca}_{1-x}(\text{Al,Si})_{3+x}\text{O}_6$ at 531 and 1579 cm^{-1} while $\text{Ca}_{3-x}(\text{Al,Si})_{6+x}\text{O}_{14}$ has a strong peak at 553 cm^{-1} . $\text{CaAl}_{4-x}\text{Si}_x\text{O}_7$ shows a weak Raman pattern, while $\text{Ca}_{1-x}(\text{Al,Si})_{2+x}\text{O}_5$ has no detectable Raman modes. Since the association moissanite-CAS is thermodynamically not stable at ambient pressure and moissanite crystals hosting the CAS phases have $\delta^{13}\text{C}$ values typical of deep-mantle origin, we interpret the CAS inclusions as partially retrogressed HP minerals. Striking analogies exist between observed CAS compositions and experimentally obtained HP-HT mineralogy. For instance, $\text{Ca}_x(\text{Al,Si})_{1-x}\text{O}_3$ contains up to 25 mol% of Al_2O_3 , which is considered as the upper limit of alumina solubility in Ca-perovskite. The study confirms that CAS phases are an important mantle depository for large ion lithophile elements (LILE) and LREE.

Keywords: Moissanite, dmisteinbergite, ghlenite, unknown CAS mineral, Raman spectra, mineral composition