

## **FTIR spectroscopy of D<sub>2</sub>O and HDO molecules in the *c*-axis channels of synthetic beryl**

**RUDOLF I. MASHKOVTSSEV<sup>1,\*</sup>, VIKTOR G. THOMAS<sup>1</sup>, DMITRY A. FURSENKO<sup>1</sup>, ELENA S. ZHUKOVA<sup>2,3</sup>,  
VLADIMIR V. USKOV<sup>2</sup>, AND BORIS P. GORSHUNOV<sup>2,3</sup>**

<sup>1</sup>V.S. Sobolev Institute of Geology and Mineralogy SB RAS, 630090 Novosibirsk, Russia

<sup>2</sup>Moscow Institute of Physics and Technology, Dolgoprudny, 141700 Moscow Region, Russia

<sup>3</sup>A.M. Prokhorov General Physics Institute RAS, 119991 Moscow, Russia

### **ABSTRACT**

This paper presents the results of Fourier transform infrared (FTIR) spectroscopy of a synthetic beryl, containing D<sub>2</sub>O molecules in its *c*-axis channels, which we synthesized under hydrothermal conditions at 600 °C and 1.5 kbar. The frequencies of absorbance bands in the range of the stretching vibrations and their overtones and combination modes for D<sub>2</sub>O and HDO molecules have been assigned for the first time. On the basis of our assignments, the absorbance bands observed for the natural beryl in the range of the OD stretching vibrations have been explained.

**Keywords:** Beryl; emerald; FTIR spectroscopy; H<sub>2</sub>O, D<sub>2</sub>O, and HDO molecules