

## **Gaussian thermoluminescence in long-range disordered K-feldspar**

**LUIS SÁNCHEZ-MUÑOZ<sup>1,\*</sup>, JAVIER GARCÍA-GUINEA<sup>2</sup>, PETER D. TOWNSEND<sup>3</sup>, TJIPTO JUWONO<sup>4</sup>,  
AND ANA CREMADES<sup>5</sup>**

<sup>1</sup>Instituto de Cerámica y Vidrio (CSIC), Kelsen 5, 28049 Madrid, Spain

<sup>2</sup>Museo Nacional Ciencias Naturales (CSIC), José Gutiérrez Abascal 2, 28006 Madrid, Spain

<sup>3</sup>School of Engineering, University of Sussex, Brighton, U.K.

<sup>4</sup>Center for Econophysics, Surya University, Tangerang, Banten, Indonesia

<sup>5</sup>Dipartimento, Física de Materiales, Fac. Físicas, University Complex Madrid, 28040 Madrid, Spain

### **ABSTRACT**

The thermoluminescence behavior of long-range ordered crystals is usually explained by the band structure model, using first- and second-order kinetics. However, feldspars have order-disorder phenomena and twinning, and consequently these mathematical descriptions are not helpful in most cases. In this work, the thermally stimulated intrinsic blue luminescence at 440 nm from X-ray induced defects of the K-rich feldspars is used to show a progressive behavior change along the order-disorder series. It is observed a gradual conversion of the TL signal from a very asymmetric peak with exponential rise and power law decay in microcline and orthoclase, where a  $\tau$  coefficient in log-log plots decreases with twin/domain size, to a more symmetric signal in a partially disordered sanidine, up to reach a completely symmetric Gaussian peak in fully disordered sanidine. These results are compatible with the Bäessler's model of disorder, which suggest that atomic disorder involves the transformation of delocalized bands first into band tails as the source of electron traps, and later in localized density of states following a Gaussian distribution.

**Keywords:** K-feldspars, thermoluminescence, order-disorder series, density of states