

SPECIAL COLLECTION: WATER IN NOMINALLY HYDROUS AND ANHYDROUS MINERALS

## Hydrogen incorporation mechanisms in forsterite: New insights from $^1\text{H}$ and $^{29}\text{Si}$ NMR spectroscopy and first-principles calculation

XIANYU XUE<sup>1,\*</sup>, MASAMI KANZAKI<sup>1</sup>, DOREEN TURNER<sup>2</sup>, AND DOMINIK LOROCH<sup>3</sup>

<sup>1</sup>Institute for Planetary Materials, Okayama University, Misasa, Tottori 682-0193 Japan

<sup>2</sup>Institute of Geoscience, University of Jena, Burgweg 11, Jena 07749 Germany

<sup>3</sup>Institute for Mineralogy, University of Muenster, Corrensstrasse 24, Muenster, 48149 Germany

### ABSTRACT

The presence of water (hydrogen) in nominally anhydrous mantle minerals may have profound effects on their physical properties (e.g., electrical conductivity, diffusivity, rheology), and these effects are expected to depend on how water is incorporated in the crystal structure. For olivine, the most abundant upper mantle mineral, despite extensive studies, mostly using vibrational spectroscopy, the interpretations are still not well constrained. To provide better understanding on this issue, we carried out a comprehensive  $^1\text{H}$  and  $^{29}\text{Si}$  NMR study on an  $\text{Mg}_2\text{SiO}_4$  forsterite sample containing about 0.5 wt%  $\text{H}_2\text{O}$  synthesized at 12 GPa and 1200 °C, complemented by Raman measurement and first-principles calculation of the geometry, stability, and NMR parameters of model structures. The Raman spectra contain relatively sharp O-H stretching bands near 3612, 3579, and 3567  $\text{cm}^{-1}$  and a broader band near 3547  $\text{cm}^{-1}$ , similar to previous reports. The  $^1\text{H}$  static and MAS NMR data revealed that there are two main populations of protons in the hydrous forsterite structure, one experiencing strong  $^1\text{H}$ - $^1\text{H}$  homonuclear dipolar couplings and contributing to a broad peak near 2.4 ppm, and another with weaker dipolar couplings and contributing to a narrower peak near 1.2 ppm in the MAS NMR spectrum at 30 kHz. Two-dimensional  $^1\text{H}$  CRAMPS-MAS NMR measurements confirmed that the two proton components belong to the same phase and the contrast in MAS NMR peak width is largely due to difference in the strength of  $^1\text{H}$ - $^1\text{H}$  homonuclear dipolar couplings. In addition, there is also a very weak, narrow  $^1\text{H}$  MAS NMR peak near 7.3 ppm (contributing to <0.1% of the total intensity) due to protons that are more remote from the two main components. First-principles calculation confirmed that the two main proton components can be attributed to the hydrogarnet-like substitution mechanism of four H ions for one Si [(4H)<sub>Si</sub>] in a tetrahedral site of olivine, but unlike hydrogarnet with one of the protons pointing away from the tetrahedral center and located in an adjacent interstitial site, thus experiencing weaker dipolar couplings than those in the vicinity of the vacant tetrahedron; the very weak narrow peak near 7.3 ppm can be attributed to the substitution mechanism of two H ions for one Mg in an M1 site [(2H)<sub>M1</sub>] of forsterite. The  $^1\text{H}$ - $^{29}\text{Si}$  CP-MAS NMR spectra revealed both a broad peak encompassing the position for OH defect-free forsterite (−61.7 ppm) and a narrower peak at higher frequency (−60.9 ppm). First-principles calculation indicates that these peaks are accountable by the same models as for the  $^1\text{H}$  NMR data. Thus, this study has provided unambiguous evidence supporting that hydrogen is incorporated in forsterite at relatively high-pressure dominantly as (4H)<sub>Si</sub> defects, with (2H)<sub>M1</sub> defects playing only a very minor role. The much larger  $^1\text{H}$  chemical shift for protons associated with the latter (than the former) is correlated with stronger hydrogen bonding for the latter, which in turn reflects difference in bonding environments of the OH groups (with the latter bonded to a Si, and the former only bonded to Mg). Similar correlation applies to the O-H stretching frequency. The (4H)<sub>Si</sub> defects are responsible for the observed high-frequency O-H stretching bands (>3450  $\text{cm}^{-1}$ ), and the (2H)<sub>M1</sub> defects give lower frequencies (undetected here due to low abundance, but most likely near 3160–3220  $\text{cm}^{-1}$  as previously reported) in vibrational spectra. These results can serve as a guide for (re-)interpretation of infrared and Raman spectroscopic data on hydrous olivine produced under different pressure and silica activity conditions, and require reconsideration of any models for the effects of water on physical properties of olivine based on different interpretations of such data. This study also demonstrated the usefulness of the combined solid-state NMR and first-principles calculation approach in unraveling the hydrogen incorporation mechanisms in nominally anhydrous minerals.

**Keywords:** Forsterite, water, structure, Si vacancy, Mg vacancy, NMR, first-principles calculation