

Carbocernaite from Bear Lodge, Wyoming: Crystal chemistry, paragenesis, and rare-earth fractionation on a microscale

ANTON R. CHAKHMOURADIAN^{1,*}, MARK A. COOPER¹, EKATERINA P. REGUIR¹, AND MEGHAN A. MOORE²

¹Department of Geological Sciences, University of Manitoba, Winnipeg, Manitoba R3T 2N2, Canada

²4016 West Lennox Loop, Coeur d'Alene, Idaho 83815, U.S.A.

ABSTRACT

Zoned crystals of carbocernaite occur in hydrothermally reworked burbankite-fluorapatite-bearing calcite carbonatite at Bear Lodge, Wyoming. The mineral is paragenetically associated with pyrite, strontianite, barite, ancylite-(Ce), and late-stage calcite, and is interpreted to have precipitated from sulfate-bearing fluids derived from an external source and enriched in Na, Ca, Sr, Ba, and rare-earth elements (REE) through dissolution of the primary calcite and burbankite. The crystals of carbocernaite show a complex juxtaposition of core-rim, sectoral, and oscillatory zoning patterns arising from significant variations in the content of all major cations, which can be expressed by the empirical formula $(\text{Ca}_{0.43-0.91}\text{Sr}_{0.40-0.69}\text{REE}_{0.18-0.59}\text{Na}_{0.18-0.53}\text{Ba}_{0-0.08})_{\Sigma 1.96-2.00}(\text{CO}_3)_2$. Interelement correlations indicate that the examined crystals can be viewed as a solid solution between two hypothetical end-members, $\text{CaSr}(\text{CO}_3)_2$ and $\text{NaREE}(\text{CO}_3)_2$, with the most Na-REE-rich areas in pyramidal (morphologically speaking) growth sectors representing a probable new mineral species. Although the Bear Lodge carbocernaite is consistently enriched in light REE relative to heavy REE and Y (chondrite-normalized $\text{La/Er} = 500\text{--}4200$), the pyramidal sectors exhibit a greater degree of fractionation between these two groups of elements relative to their associated prismatic sectors. A sample approaching the solid-solution midline $[(\text{Ca}_{0.57}\text{Na}_{0.42})_{\Sigma 0.99}(\text{Sr}_{0.50}\text{REE}_{0.47}\text{Ba}_{0.01})_{\Sigma 0.98}(\text{CO}_3)_2]$ was studied by single-crystal X-ray diffraction and shown to have a monoclinic symmetry [space group $P11m$, $a = 6.434(4)$, $b = 7.266(5)$, $c = 5.220(3)$ Å, $\gamma = 89.979(17)^\circ$, $Z = 2$] as opposed to the orthorhombic symmetry (space group $Pb2_1m$) proposed in earlier studies. The symmetry reduction is due to partial cation order in sevenfold-coordinated sites occupied predominantly by Ca and Na, and in tenfold-coordinated sites hosting Sr, REE, and Ba. The ordering also causes splitting of carbonate vibrational modes at 690–740 and 1080–1100 cm^{-1} in Raman spectra. Using Raman micro-spectroscopy, carbocernaite can be readily distinguished from burbankite- and ancylite-group carbonates characterized by similar energy-dispersive spectra.

Keywords: Carbocernaite, carbonatite, isotopic composition, rare earth elements, crystal structure, zoning, Bear Lodge, Wyoming