

## **Radiation damage in sulfides: Radioactive galena from burning heaps, after coal mining in the Lower Silesian basin (Czech Republic)**

**MICHAL ČURDA<sup>1,2,\*</sup>, VIKTOR GOLIÁŠ<sup>1</sup>, MARIANA KLEMENTOVÁ<sup>3</sup>, LADISLAV STRNAD<sup>4</sup>,  
ZDENĚK MATĚJ<sup>5</sup>, AND RADEK ŠKODA<sup>6</sup>**

<sup>1</sup>Institute of Geochemistry, Mineralogy and Mineral Resources, Charles University, Faculty of Science, Albertov 6, 128 43 Prague 2, Czech Republic

<sup>2</sup>Czech Geological Survey, Geologická 6, 152 00 Prague 5, Czech Republic

<sup>3</sup>Institute of Inorganic Chemistry, Czech Academy of Sciences, 250 68 Husinec-Řež, Czech Republic

<sup>4</sup>Laboratories of the Geological Institutes, Faculty of Science, Charles University, Albertov 6, 128 43 Prague 2, Czech Republic

<sup>5</sup>Department of Condensed Matter Physics, Faculty of Mathematics and Physics, Charles University, Ke Karlovu 5, 121 16 Prague 2, Czech Republic

<sup>6</sup>Institute of Earth Sciences, Faculty of Science, Masaryk University, Kotlářská 2, 611 37 Brno, Czech Republic

### **ABSTRACT**

The isotopic composition of lead ( $^{207}\text{Pb}/^{206}\text{Pb}$ ,  $^{208}\text{Pb}/^{206}\text{Pb}$ , and  $^{210}\text{Pb}$ ) in a recently formed galena from burning heaps after coal mining in Radvanice, Markoušovice, and Rybníček, the Lower Silesian basin, Czech Republic, was studied in detail.  $^{210}\text{Pb}$  activity in galena varied from  $135 \pm 9$  to  $714 \pm 22$  Bq/g and calculated integral doses ranged from  $2.21 \times 10^{11}$  to  $6.11 \times 10^{11}$   $\alpha/\text{g}$ . The radioactivity of the galena causes micro-deformations in its crystal structure as indicated by the Williamson-Hall graphs, showing that the level of micro-strain depends on the length of time that galena samples were exposed to the radiation. However, the crystal structure of galena is affected very inhomogeneously; according to TEM investigations there are domains of fully crystalline, polycrystalline, and fully metamict galena within one crystal. Inductively coupled plasma-mass spectrometry (ICP-MS) was used to determine the isotopic composition of the studied galena. The stable isotope ratios of Pb varied for  $^{207}\text{Pb}/^{206}\text{Pb}$  from 0.8402 to 0.8435 and for  $^{208}\text{Pb}/^{206}\text{Pb}$  from 2.0663 to 2.0836. The average ratios  $^{207}\text{Pb}/^{206}\text{Pb} = 0.8312$  and  $^{208}\text{Pb}/^{206}\text{Pb} = 2.0421$  were obtained for coal from the same localities. These isotope ratios show that there is no isotopic fractionation taking place during the coal burning and subsequent galena crystallization from hot gases.

**Keywords:** Galena, radiation, lead-isotopes, radiation effects, metamict state