

## High-pressure phase behavior and equations of state of ThO<sub>2</sub> polymorphs

BETHANY A. CHIDESTER<sup>1,\*</sup>, OLIVIA S. PARDO<sup>1,†</sup>, REBECCA A. FISCHER<sup>1,‡</sup>, ELIZABETH C. THOMPSON<sup>1</sup>,  
DION L. HEINZ<sup>1</sup>, CLEMENS PRESCHER<sup>2,§</sup>, VITALI B. PRAKAPENKA<sup>2</sup>, AND ANDREW J. CAMPBELL<sup>1</sup>

<sup>1</sup>Department of the Geophysical Sciences, University of Chicago, Chicago, Illinois 60637, U.S.A.

<sup>2</sup>GeoSoilEnviroCARS, University of Chicago, Argonne National Laboratory, Argonne, Illinois 60439, U.S.A.

### ABSTRACT

ThO<sub>2</sub> is an important material for understanding the heat budget of Earth's mantle, as well as the stability of nuclear fuels at extreme conditions. We measured the in situ high-pressure, high-temperature phase behavior of ThO<sub>2</sub> to ~60 GPa and ~2500 K. It undergoes a transition from the cubic fluorite-type structure (thorianite) to the orthorhombic  $\alpha$ -PbCl<sub>2</sub> cotunnite-type structure between 20 and 30 GPa at room temperature. Prior to the transition at room temperature, an increase in unit-cell volume is observed, which we interpret as anion sub-lattice disorder or pre-transformation “melting” (Bouffeffel et al. 2006). The thermal equation of state parameters for both thorianite [ $V_0 = 26.379(7)$ ,  $K_0 = 204(2)$ ,  $\alpha K_T = 0.0035(3)$ ] and the high-pressure cotunnite-type phase [ $V_0 = 24.75(6)$ ,  $K_0 = 190(3)$ ,  $\alpha K_T = 0.0037(4)$ ] are reported, holding  $K'_0$  fixed at 4. The similarity of these parameters suggests that the two phases behave similarly within the deep Earth. The lattice parameter ratios for the cotunnite-type phase change significantly with pressure, suggesting a different structure is stable at higher pressure.

**Keywords:** XRD data, ThO<sub>2</sub>, Raman spectroscopy, ThO<sub>2</sub>, phase transition, high-pressure studies, diamond-anvil cell, high-temperature studies, laser-heating