

LETTER

UHP Ti-chondrodite in the Zermatt-Saas serpentinite: Constraints on a new tectonic scenario

PIETRO LUONI^{1,*}, GISELLA REBAY², MARIA IOLE SPALLA¹, AND DAVIDE ZANONI¹

¹Dipartimento di Scienze della Terra ‘A. Desio’, Università degli Studi di Milano, Via Mangiagalli, 34-20133 Milano, Italy

²Dipartimento di Scienze della Terra e dell’Ambiente, Università degli Studi di Pavia, Via Ferrata, 1-27100 Pavia, Italy

ABSTRACT

We focus on the key role of different Ti-humite minerals in subducted serpentinites as possible indicators of extreme pressure conditions. The occurrence of Ti-chondrodite and/or Ti-clinohumite assemblages in the eclogitized serpentinites of the Zermatt-Saas Zone (ZSZ) of the Western Alps allows the recrystallization of such rocks at UHP conditions ($P = 2.8\text{--}3.5$ GPa, $T = 600\text{--}670$ °C) to be determined. Such conditions are similar to those registered by the nearby Cignana unit, a main Alpine area for UHP metamorphism, where coesite and microdiamond have been found. In ZSZ serpentinites, the new UHP assemblage predates the previously recognized HP-UHP paragenesis, which was recently dated at 65 Ma. This finding opens up a new interpretation for the petrologically and structurally well-constrained HP/UHP records, especially because all other ages for HP-UHP metamorphism in the ZSZ are much younger, and for the size of UHP units. Our findings suggest that ophiolites in the axial zone of collisional belts are a mosaic of oceanic lithosphere slices that recorded contrasted thermal and mechanical evolutions during their physical trajectories in the subduction wedge.

Keywords: Ti-clinohumite and Ti-chondrodite assemblages, integrated mineralogical and structural analysis, Alpine subduction, Western Alps