

Solution mechanisms of COHN fluids in melts to upper mantle temperature, pressure, and redox conditions

BJORN MYSEN^{1,*}

¹Geophysical Laboratory, Carnegie Institution of Washington, Washington, D.C. 20015, U.S.A.

ABSTRACT

We wished to advance the knowledge of speciation among volatiles during melting and crystallization in the Earth's interior; therefore, we explored the nature of carbon-, nitrogen-, and hydrogen-bearing species as determined in COHN fluids and dissolved in coexisting aluminosilicate melts. Micro-Raman characterization of fluids and melts were conducted in situ while samples were at a temperature up to 825 °C and pressure up to ~1400 MPa under redox conditions controlled with the Ti-TiO₂-H₂O hydrogen fugacity buffer. The fluid species are H₂O, H₂, NH₃, and CH₄. In contrast, under oxidizing conditions, the species are H₂O, N₂, and CO₂.

The equilibria among silicate structures (Q-species) and reduced carbon and nitrogen species are, $2\text{NH}_3 + 4\text{Q}^n \rightleftharpoons 2\text{Q}^{n-1}(\text{NH}_2) + 2\text{Q}^{n-1}(\text{OH})$, and $2\text{CH}_4 + 4\text{Q}^n \rightleftharpoons 2\text{Q}^{n-1}(\text{CH}_3) + 2\text{Q}^{n-1}(\text{OH})$. The Qⁿ and Qⁿ⁻¹ denote silicate species with, respectively, n and n-1 bridging O atoms. The formulation in parentheses, (NH₂), (CH₃), and (OH), is meant to indicate that those functional groups replace one or more oxygen in the silicate tetrahedra. There is no evidence for O-NH₂ or O-CH₃ bonding. Therefore, a solution of reduced C- and N-species in the COHN system results in depolymerization of silicate melts. The ΔH values derived from the $X_{\text{NH}_2}/X_{\text{NH}_3}$ and $X_{\text{CH}_3}/X_{\text{CH}_4}$ evolution with temperature, respectively, were 8.1 ± 2.3 kJ/mol and between -4.9 ± 1.0 and -6.2 ± 2.2 kJ/mol.

The fluid/melt partition coefficients, $K^{\text{fluid/melt}}$, of the reduced species, H₂O, H₂, NH₃, and CH₄, remain above unity at all temperatures. For example, for carbon it is in the 6–15 range with a $\Delta H = -13.4 \pm 2.4$ kJ/mol. These values compare with a 0.8–3 range with $\Delta H = -19 \pm 2.4$ kJ/mol in N-free silicate-CO systems. The $K^{\text{fluid/melt}}$ values for reduced nitrogen and molecular hydrogen are in the 6–10 and 6–12 range with ΔH values of -5.9 ± 0.9 and 8 ± 6 kJ/mol, respectively.

A change in redox conditions during melting and crystallization in the Earth sufficient to alter oxidized to reduced carbon- and nitrogen-bearing species will affect all melt properties that depend on melt polymerization. This suggestion implies that changing redox conditions during melting of a COHN-bearing mantle can have a profound effect on physical and chemical properties of melts and on melting and melt aggregation processes.

Keywords: Volatiles, COHN, Redox, pressure, structure, mantle