

## **Quantitative measurement of olivine composition in three dimensions using helical-scan X-ray micro-tomography**

**MATTHEW J. PANKHURST<sup>1,2,3,4,\*</sup>, NGHIA T. VO<sup>5</sup>, ALAN R. BUTCHER<sup>6,7</sup>, HAILI LONG<sup>6</sup>,  
HONGCHANG WANG<sup>7</sup>, SARA NONNI<sup>4</sup>, JASON HARVEY<sup>3</sup>, GUÐMUNDUR GUÐFINNSSON<sup>8</sup>, RONALD FOWLER<sup>9</sup>,  
ROBERT ATWOOD<sup>4,5</sup>, RICHARD WALSHAW<sup>10</sup>, AND PETER D. LEE<sup>4,11</sup>**

<sup>1</sup>Instituto Tecnológico y de Energías Renovables (ITER), 38900 Granadilla de Abona, Tenerife, Canary Islands, Spain

<sup>2</sup>Instituto Volcanológico de Canarias (INVOLCAN), 38400 Puerto de la Cruz, Tenerife, Canary Islands, Spain

<sup>3</sup>School of Earth and Environment, University of Leeds, Leeds, LS2 9JT, U.K.

<sup>4</sup>Research Complex at Harwell, Rutherford Appleton Laboratories, Didcot, OX11 0FA, U.K.

<sup>5</sup>Diamond Light Source Ltd., Didcot, OX11 0DE, U.K.

<sup>6</sup>FEI, Stiklestadveien 1, 7041 Trondheim, Norway

<sup>7</sup>Geological Survey of Finland, FI-02151 Espoo, Finland

<sup>8</sup>Institute of Earth Sciences, University of Iceland, Reykjavik 101, Iceland

<sup>9</sup>Scientific Computing Department, Science and Technology Facilities Council, Rutherford Appleton Laboratory, Harwell Campus, OX11 0QX, U.K.

<sup>10</sup>Leeds Electron Microscopy and Spectroscopy Centre, University of Leeds, Leeds, LS2 9JT, U.K.

<sup>11</sup>Mechanical Engineering, University College London, Gower Street, London, WC1E 6BT, U.K.

### **ABSTRACT**

Olivine is a key constituent in the silicate Earth; its composition and texture informs petrogenetic understanding of numerous rock types. Here we develop a quantitative and reproducible method to measure olivine composition in three dimensions without destructive analysis, meaning full textural context is maintained. The olivine solid solution between forsterite and fayalite was measured using a combination of three-dimensional (3D) X-ray imaging techniques, 2D backscattered electron imaging, and spot-analyses using wavelength-dispersive electron probe microanalysis. The linear attenuation coefficient of natural crystals across a range of forsterite content from ~73–91 mol% were confirmed to scale linearly with composition using 53, 60, and 70 kV monochromatic beams at I12-JEEP beamline, Diamond Light Source utilizing the helical fly-scan acquisition. A polychromatic X-ray source was used to scan the same crystals, which yielded image contrast equivalent to measuring the mol% of forsterite with an accuracy of <1.0%. X-ray tomography can now provide fully integrated textural and chemical analysis of natural samples containing olivine, which will support 3D and 3D+time petrologic modeling. The study has revealed >3 mm domains within a large crystal of San Carlos forsterite that varies by ~2 Fo mol%. This offers a solution to an outstanding question of inter-laboratory standardization, and also demonstrates the utility of 3D, non-destructive, chemical measurement. To our knowledge, this study is the first to describe the application of XMT to quantitative chemical measurement across a mineral solid solution. Our approach may be expanded to calculate the chemistry of other mineral systems in 3D, depending upon the number, chemistry, and density of end-members.

**Keywords:** Micro-computed tomography, densitometry, olivine, chemical composition