

Geochronology and trace element mobility in rutile from a Carboniferous syenite pegmatite and the role of halogens

GEORGIA PE-PIPER^{1,*}, JUSTIN NAGLE¹, DAVID J.W. PIPER², AND CHRIS R.M. MCFARLANE³

¹Department of Geology, Saint Mary's University, Halifax, Nova Scotia B3H 3C3, Canada; ORCID 0000-0002-7670-7358

²Natural Resources Canada, Geological Survey of Canada (Atlantic), Bedford Institute of Oceanography, P.O. Box 1006, Dartmouth, Nova Scotia B2Y 4A2, Canada; ORCID 0000-0002-6276-7560

³Department of Earth Sciences, University of New Brunswick, 2 Bailey Drive, Fredericton, New Brunswick E3B 5A3, Canada

ABSTRACT

This study investigates Ti mobility in the presence of halogens, as shown by the hydrothermal alteration of magmatic rutile in syenite. The syenite pegmatite studied intrudes gabbro, is preserved as a tectonic block in a major strike-slip fault zone, and formed in a back-arc environment in which there was widespread A-type granite plutonism. Rutile was studied by SEM and Raman spectroscopy, trace elements were analyzed by LA-ICP-MS, and age was determined by in situ U-Pb analysis. Magmatic rutile in the syenite forms millimetric-scale crystals rimmed by magmatic titanite and magnetite and also occurs as smaller interstitial crystals. Hydrothermal alteration occurred preferentially along crystal margins and fractures by a layer-by-layer dissolution-reprecipitation process resulting in high Zr contents (~5000 ppm) in the rutile, together with enrichment in U and depletion in high field strength elements. The magmatic emplacement age of the syenite was ~360 Ma (dated rutile G) and no younger than 353.9 ± 5.7 Ma (mean Concordia age of interstitial rutile). The syenite was synchronous with the later phases of regional A-type granite plutonism. Most magmatic rutile has REE patterns either (1) with 1–50 times chondrite enrichment, LREE > HREE and a Eu anomaly, resulting from felsic melt inclusions, or (2) flat patterns with 0.1–10 times chondrite enrichment, present in ilmenite exsolution lamellae or inclusions. Later hydrothermal halogen-rich fluids, derived from dissolution of halite, produced widespread metasomatic scapolite in the syenite. These fluids also leached Ti and other HFSE, together with REE, from large fractured rutile crystals. Such fluids resulted in local dissolution-reprecipitation of Ti and Zr and resetting of the U-Pb system in the altered rutile, at 337.4 ± 3.5 Ma. Normalized REE abundances in the hydrothermal rutile show a U-shaped pattern, with the greatest depletion in the MREE. Variations in dissolution and transport of Zr and Ti by halogen-rich fluids affect the Zr in rutile geothermometer, which yields unrealistic temperatures when applied in this study. More generally, the complexities of rutile chemistry in this hydrothermal setting could be reproduced in deeper subduction settings as a result of variations in halogen content of fluids released by prograde metamorphism.

Keywords: Rutile, halogens, zirconium, hydrothermal, dissolution-reprecipitation