

LETTER

Discovery of asimowite, the Fe-analog of wadsleyite, in shock-melted silicate droplets of the Suizhou L6 and the Quebrada Chimborazo 001 CB3.0 chondrites

LUCA BINDI^{1,*}, FRANK E. BRENKER², FABRIZIO NESTOLA³, TAMARA E. KOCH², DAVID J. PRIOR⁴, KAT LILLY⁴, ALEXANDER N. KROT⁵, MARTIN BIZZARRO⁶, AND XIANDE XIE^{7,8}

¹Dipartimento di Scienze della Terra, Università degli Studi di Firenze, Via G. La Pira 4, I-50121 Firenze, Italy

²Goethe University Frankfurt, NanoGeoscience, Altenhoferallee 1, 60438 Frankfurt, Germany

³Dipartimento di Geoscienze, Università di Padova, Via Gradenigo, 6, 35131 Padova, Italy

⁴University of Otago, Department of Geology, Dunedin 9054, New Zealand

⁵University of Hawaii at Mānoa, Hawaii Institute of Geophysics and Planetology, 1680 East-West Road, Honolulu, Hawaii 96822, U.S.A.

⁶StarPlan—Centre for Star and Planet Formation, Natural History Museum of Copenhagen University of Copenhagen, Øster Voldgade 5-7, DK-1350, Copenhagen, Denmark

⁷Key Laboratory of Mineralogy and Metallogeny, Guangzhou Institute of Geochemistry, Chinese Academy of Sciences, Guangzhou 510640, China

⁸Guangdong Provincial Key Laboratory of Mineral Physics and Materials, Guangzhou 510640, China

ABSTRACT

We report the first natural occurrence and single-crystal X-ray diffraction study of the Fe-analog of wadsleyite [$a = 5.7485(4)$, $b = 11.5761(9)$, $c = 8.3630(7)$ Å, $V = 556.52(7)$ Å³; space group *Imma*], spinelloid-structured Fe₂SiO₄, a missing phase among the predicted high-pressure polymorphs of ferroan olivine, with the composition (Fe_{1.10}Mg_{0.80}Cr_{0.04}Mn_{0.02}Ca_{0.02}Al_{0.02}Na_{0.01})_{Σ2.01}(Si_{0.97}Al_{0.03})_{Σ1.00}O₄. The new mineral was approved by the International Mineralogical Association (No. 2018-102) and named asimowite in honor of Paul D. Asimow, the Eleanor and John R. McMillan Professor of Geology and Geochemistry at the California Institute of Technology. It was discovered in rare shock-melted silicate droplets embedded in Fe,Ni-metal in both the Suizhou L6 chondrite and the Quebrada Chimborazo (QC) 001 CB3.0 chondrite. Asimowite is rare, but the shock-melted silicate droplets are very frequent in both meteorites, and most of them contain Fe-rich wadsleyite (Fa_{30–45}). Although the existence of such Fe-rich wadsleyite in shock veins may be due to the kinetic reasons, new theoretical and experimental studies of the stability of (Fe,Mg)₂SiO₄ at high temperature (>1800 K) and pressure are clearly needed. This may also have a significant impact on the temperature and chemical estimates of the mantle's transition zone in Earth.

Keywords: Wadsleyite, iron, spinelloid, chondrite, meteorite, crystal structure, microprobe analysis, Earth's transition zone