

Nixonite, Na₂Ti₆O₁₃, a new mineral from a metasomatized mantle garnet pyroxenite from the western Rae Craton, Darby kimberlite field, Canada

CHIARA ANZOLINI^{1,*}, FEI WANG², GARRETT A. HARRIS¹, ANDREW J. LOCOCK¹, DONGZHOU ZHANG³, FABRIZIO NESTOLA⁴, LUCA PERUZZO⁵, STEVEN D. JACOBSEN², AND D. GRAHAM PEARSON¹

¹Department of Earth and Atmospheric Sciences, 1-26 Earth Sciences Building, University of Alberta, Edmonton, Alberta T6G 2E3, Canada

²Department of Earth and Planetary Sciences, Northwestern University, Evanston, Illinois 60208, U.S.A.

³School of Ocean and Earth Science and Technology, Hawai'i Institute of Geophysics and Planetology, University of Hawaii at Manoa, Honolulu, Hawaii 966822, U.S.A. Orcid 0000-0002-6679-892X

⁴Department of Geosciences, University of Padova, Via G. Gradenigo 6, 35131 Padova, Italy. Orcid 0000-0002-4875-5125

⁵CNR, Istituto di Geoscienze e Georisorse, Sezione di Padova, Via G. Gradenigo 6, 35131 Padova, Italy

ABSTRACT

Nixonite (IMA 2018-133), ideally Na₂Ti₆O₁₃, is a new mineral found within a heavily metasomatized pyroxenite xenolith from the Darby kimberlite field, beneath the west-central Rae Craton, Canada. It occurs as microcrystalline aggregates, 15 to 40 μm in length. Nixonite is isostructural with jeppeite, K₂Ti₆O₁₃, with a structure consisting of edge- and corner-shared titanium-centered octahedra that enclose alkali-metal ions. The Mohs hardness is estimated to be between 5 and 6 by comparison to jeppeite, and the calculated density is 3.51(1) g/cm³. Electron microprobe wavelength-dispersive spectroscopic analysis (average of 6 points) yielded: Na₂O 6.87, K₂O 5.67, CaO 0.57, TiO₂ 84.99, V₂O₃ 0.31, Cr₂O₃ 0.04, MnO 0.01, Fe₂O₃ 0.26, SrO 0.07, total 98.79 wt%. The empirical formula, based on 13 O atoms, is: (Na_{1.24}K_{0.67}Ca_{0.06})_{Σ1.97}(Ti_{5.96}V_{0.023}Fe_{0.018})_{Σ6.00}O₁₃ with minor amounts of Cr and Mn. Nixonite is monoclinic, space group *C2/m*, with unit-cell parameters *a* = 15.3632(26) Å, *b* = 3.7782(7) Å, *c* = 9.1266(15) Å, β = 99.35(15)°, and *V* = 522.72(1) Å³, *Z* = 2. Based on the average of seven integrated multi-grain diffraction images, the strongest diffraction lines are [*d*_{obs} in Å (*I* in %) (*hkl*)]: 3.02 (100) (310), 3.66 (75) (110), 7.57 (73) (200), 6.31 (68) (20 $\bar{1}$), 2.96 (63) (31 $\bar{1}$), 2.96 (63) (20 $\bar{3}$), and 2.71 (62) (402). The five main Raman peaks of nixonite, in order of decreasing intensity, are at 863, 280, 664, 135, and 113 cm⁻¹. Nixonite is named after Peter H. Nixon, a renowned scientist in the field of kimberlites and mantle xenoliths. Nixonite occurs within a pyroxenite xenolith in a kimberlite, in association with rutile, priderite, perovskite, freudenbergite, and ilmenite. This complex Na-K-Ti-rich metasomatic mineral assemblage may have been produced by a fractionated Na-rich kimberlitic melt that infiltrated a mantle-derived garnet pyroxenite and reacted with rutile during kimberlite crystallization.

Keywords: Nixonite, new mineral, crystal structure, jeppeite, mantle xenolith, kimberlite, Rae Craton