

## **Melting in the Fe-FeO system to 204 GPa: Implications for oxygen in Earth's core**

**KENTA OKA<sup>1,\*</sup>, KEI HIROSE<sup>1,2</sup>, SHOH TAGAWA<sup>1</sup>, YUTO KIDOKORO<sup>1</sup>, YOICHI NAKAJIMA<sup>3,\*</sup>,  
YASUHIRO KUWAYAMA<sup>1</sup>, GUILLAUME MORARD<sup>4</sup>, NICOLAS COUDURIER<sup>4</sup>, AND GUILLAUME FIQUET<sup>4</sup>**

<sup>1</sup>Department of Earth and Planetary Science, The University of Tokyo, 7-3-1 Hongo, Bunkyo, Tokyo 113-0033, Japan

<sup>2</sup>Earth-Life Science Institute, Tokyo Institute of Technology, 2-12-1 Ookayama, Meguro, Tokyo 152-8550, Japan

<sup>3</sup>Department of Physics, Kumamoto University, 2-39-1 Kurokami, Chuo-ku, Kumamoto, Kumamoto 860-8555, Japan

<sup>4</sup>Sorbonne Université, Muséum National d'Histoire Naturelle, UMR CNRS 7590, Institut de Minéralogie, de Physique des Matériaux et de Cosmochimie, IMPMC, 75005 Paris, France

### **ABSTRACT**

We performed melting experiments on Fe-O alloys up to 204 GPa and 3500 K in a diamond-anvil cell (DAC) and determined the liquidus phase relations in the Fe-FeO system based on textural and chemical characterizations of recovered samples. Liquid-liquid immiscibility was observed up to 29 GPa. Oxygen concentration in eutectic liquid increased from >8 wt% O at 44 GPa to 13 wt% at 204 GPa and is extrapolated to be about 15 wt% at the inner core boundary (ICB) conditions. These results support O-rich liquid core, although oxygen cannot be a single core light element. We estimated the range of possible liquid core compositions in Fe-O-Si-C-S and found that the upper bounds for silicon and carbon concentrations are constrained by the crystallization of dense inner core at the ICB.

**Keywords:** Core, oxygen, Fe-FeO system, eutectic melting, high pressure