

Spinel-Anorthosites on the Moon: Impact Melt Origins Suggested by Enthalpy Constraints

Allan H. Treiman, Michael J. Kulis, and Allen F. Glazner

Appendix D: Equilibria Between Peridotite and Anorthosite in the System Fo-An-Sil

Extending our results toward realistic lunar compositions, we consider the system Fo-An with additional SiO₂, the system Fo-An-Sil (forsterite-anorthite-silica), Figure D1. Its ternary liquidus diagrams are shown in Figure D1a as experimentally determined by Andersen (1915) and Irvine (1975), and Figure D1b as calculated with FactSage. The topographies of the experimental and calculated diagrams are similar with a significant exception. The fields of liquidus pyroxene are nearly identical, although the calculated diagram includes several varieties of MgSiO₃ pyroxene (protopyroxene, space group Pbcn; pigeonite, P2₁/c; and orthopyroxene, Pbcn), which are rarely distinguished in experimental studies. The significant difference is that FactSage calculations yield a significantly larger spinel liquidus field than determined in experiments. The spinel-liquidus field is so enlarged that the calculated diagram has no L + Ol + Pl field (or line, Fig. D1b), which is not consistent with natural occurrences and laboratory experiments.

Rather than investigate the whole ternary system, we focus on the formation of lunar spinel by selecting a binary join in Fo-An-Sil that is relevant to hypotheses that invoke anorthosite assimilation into picritic or Mg-suite magmas (Morgan et al., 2006; Gross and Treiman, 2011; Prissel et al., 2016). The join between anorthite and a model peridotite composition of 88% forsterite by mass and 12% SiO₂ (Figure 3a, see Appendix B) passes close to the projections of the Apollo 14 B ‘green glass’ picrite, Table 1 of main text (Delano, 1986; Elkins-Tanton et al., 2003), and the suggested composition of possible parent magma compositions for the Mg-suite plutonic rocks (Longhi et al., 2010; Prissel et al., 2016). The A14B green glass is appropriate here because it is among the most picritic (i.e., richest in normative olivine; Fig. 3a) and most magnesian (highest Mg#) of the lunar magma compositions.

Figures D1c-f show equilibrium phase relations along that peridotite-anorthite join (at 1 bar), both as T-X and ΔH^* -X diagrams.

Interpretation 2: Peridotite-Anorthosite in Fo-An-Sil

Spinel stability is a problem in the FactSage calculated diagrams, as its stability fields are significantly larger than those constrained by experiments (Figs. D1a, b). Of most importance, the calculated T-X diagram shows a liquidus field of Opx + Sp + L. This difference is significant because Opx + Sp + L is not stable at low pressure (i.e., 1 bar as in the FactSage calculation); instead, Ol + Pl + L are stable together at low pressures in lunar, terrestrial, and experimental systems. The assemblage Opx + Sp + L is stable only at high pressures, i.e. > 0.3 GPa (Presnall et al., 1978), and the effect of pressure on spinel stability underlies several hypotheses for its occurrence on the Moon (Herzberg and Baker, 1980; McCallum and Schwartz, 2001; Prissel et al., 2014). This mismatch has little effect on calculated liquidus equilibria for this binary join (Figs. D1c-f), but is important at sub-liquidus conditions because it implies that spinel could form at temperatures (and in phase assemblages) different from those constrained by experiments. In Figures D1d & f, note that there are calculated stability fields for Px + Sp + Ol + L, Px + Sp + Pl + L, and Px + Sp + Ol + Pl + L; although these fields are small, they are not consistent with experimental data (Fig. 3a) and diagrams derived from them (Figs. 3c, e).




Figure D1. Phase relations in the ternary system Fo-An-Sil ($\text{Mg}_2\text{SiO}_4 - \text{CaAl}_2\text{Si}_2\text{O}_8 - \text{SiO}_2$) at 1-bar pressure.

a. Ternary liquidus surface, as constrained by laboratory experiments (Andersen, 1915; Irvine, 1975). Note that the liquidus fields of olivine and spinel touch, i.e. there are compositions and temperatures where both olivine and anorthite are on the liquidus together, and that spinel and pyroxene are never on the liquidus together. Blue line connects anorthite with a peridotite composition (Perid14B'') such that mixing along it yields model compositions like the Apollo 14 green glass (A14B'') and the Mg-suite parent magma (MGSPM'').

b. Ternary liquidus surface, as calculated in FactSage. Note that the liquidus field of spinel is larger than in part **a**, and implies that spinel and pyroxene can be liquidus phases together. This is contrary to results of laboratory experiments (see Fig. D1a).

c. Temperature-composition phase relations along the join PeridA14B'' – Anorthite, (shown as blue lines in Figures D1a & b) calculated in FactSage. Pig = pigeonite pyroxene; Ppx = protopyroxene; Opx = orthopyroxene. The field denoted by * is for L+Sp+Ol+An. Complex relations near the PeridA14B'' axis are omitted for clarity.

d. ΔH^* -X diagram for the pseudo-binary join Perid14B''-An calculated with FactSage (Bale et al., 2009; Bale et al., 2016) consistent with Fig. D1d. Pink field have phases consistent with remote sensing detection as spinel-peridotite. Pig = pigeonite pyroxene; Ppx = protopyroxene; field marked as '*' is for L+Sp+Opx+Ol+An; field marked as '**' is for L+Opx+Pig+Ol+An. The complex phase relations at lower T and lower An content (mostly involving pyroxenes) are in Figure A4.

References cited

- Andersen, O. (1915) The system anorthite-forsterite-silica. *American Journal of Science*, 232, 407-454.
- Delano, J.W. (1986) Pristine lunar glasses: Criteria, data, and implications. *Journal of Geophysical Research: Solid Earth*, 91(B4), 201-213.
- Elkins-Tanton, L.T., Chatterjee, N., and Grove, T.L. (2003) Experimental and petrological constraints on lunar differentiation from the Apollo 15 green picritic glasses. *Meteoritics & Planetary Science*, 38(4), 515-527.
- Gross, J., and Treiman, A.H. (2011) Unique spinel - rich lithology in lunar meteorite ALHA 81005: Origin and possible connection to M³ observations of the farside highlands. *Journal of Geophysical Research: Planets*, 116, E10009.
- Herzberg, C.T., and Baker, M.B. (1980) The cordierite-to spinel-cataclasite transition - Structure of the lunar crust. *Proceedings of the Conference on the Lunar Highlands Crust*, p. 113-132. Pergamon, New York.
- Irvine, T. (1975) Olivine-pyroxene-plagioclase relations in the system Mg₂SiO₄--CaAl₂Si₂O₈ - KAlSi₃O₈--SiO₂ and their bearing on the differentiation of stratiform intrusions. *Yearbook, Carnegie Institution of Washington*, 74, 492-500.
- Longhi, J., Durand, S.R., and Walker, D. (2010) The pattern of Ni and Co abundances in lunar olivines. *Geochimica et Cosmochimica Acta*, 74(2), 784-798.
- McCallum, I.S., and Schwartz, J.M. (2001) Lunar Mg suite: Thermobarometry and petrogenesis of parental magmas. *Journal of Geophysical Research: Planets*, 106(E11), 27969-27983.
- Morgan, Z., Liang, Y., and Hess, P. (2006) An experimental study of anorthosite dissolution in lunar picritic magmas: implications for crustal assimilation processes. *Geochimica et Cosmochimica Acta*, 70(13), 3477-3491.
- Presnall, D., Dixon, S.A., Dixon, J.R., O'Donnell, T., Brenner, N., Schrock, R., and Dycus, D. (1978) Liquidus phase relations on the join diopside-forsterite-anorthite from 1 atm to 20 kbar: their bearing on the generation and crystallization of basaltic magma. *Contributions to Mineralogy and Petrology*, 66(2), 203-220.

Prissel, T.C., Parman, S., Jackson, C., Rutherford, M., Hess, P., Head, J., Cheek, L., Dhingra, D., and Pieters, C. (2014) Pink Moon: The petrogenesis of pink spinel anorthosites and implications concerning Mg-suite magmatism. *Earth and Planetary Science Letters*, 403, 144-156.

Prissel, T.C., Parman, S.W., and Head, J.W. (2016) Formation of the lunar highlands Mg-suite as told by spinel. *American Mineralogist*, 101(7), 1624-1635.