

Synthesis of calcium orthocarbonate, Ca_2CO_4 -*Pnma* at *P-T* conditions of Earth's transition zone and lower mantle

Jannes Binck^{1,*,\dagger}, Dominique Laniel², Lkhamsuren Bayarjargal¹, Saiana Khandarkhaeva³, Timofey Fedotenko², Andrey Aslandukov², Konstantin Glazyrin⁴, Victor Milman⁵, Stella Chariton⁶, Vitali B. Prakapenka⁶, Natalia Dubrovinskaia², Leonid Dubrovinsky³, and Björn Winkler¹

¹Institut für Geowissenschaften, Goethe-Universität Frankfurt, Altenhöferallee 1, 60438 Frankfurt am Main, Germany

²Laboratory of Crystallography, University of Bayreuth, 95440 Bayreuth, Germany

³Bayerisches Geoinstitut, University of Bayreuth, 95440, Bayreuth, Germany

⁴Photon Science, Deutsches Elektronen-Synchrotron, Notkestrasse 85, 22607, Hamburg, Germany

⁵BIOVIA Dassault Systèmes, 334 Science Park, Cambridge CB4 0WN, U.K.

⁶Center for Advanced Radiation Sources, The University of Chicago, Chicago, Illinois 60637, U.S.A.

Abstract

We show, by single-crystal diffraction studies in laser-heated diamond-anvil cells, that Ca_2CO_4 orthocarbonate, which contains CO_4^{4-} tetrahedra, can be formed already at ~ 20 GPa at ~ 1830 K, i.e., at much lower pressures than other carbonates with sp^3 -hybridized carbon. Ca_2CO_4 can also be formed at ~ 89 GPa and ~ 2500 K. This very broad *P-T* range suggests the possible existence of Ca_2CO_4 in the Earth's transition zone and in most of the lower mantle. Raman spectroscopy shows the typical bands associated with tetrahedral CO_4^{4-} -groups. DFT-theory based calculations reproduce the experimental Raman spectra and indicate that at least in the athermal limit the phase assemblage of $\text{Ca}_2\text{CO}_4 + 2\text{SiO}_2$ is more stable than $2\text{CaSiO}_3 + \text{CO}_2$ at high pressures.

Keywords: Carbonate, Ca_2CO_4 , structure, X-ray diffraction, Raman spectroscopy, density functional theory; Volatile Elements in Differentiated Planetary Interiors