

Characterization of carbon phases in Yamato 74123 ureilite to constrain the meteorite shock history

ANNA BARBARO^{1,*}, FABRIZIO NESTOLA^{2,3}, LIDIA PITTARELLO⁴, LUDOVIC FERRIÈRE⁴, MARA MURRI⁵, KONSTANTIN D. LITASOV⁶, OLIVER CHRIST², MATTEO ALVARO¹, AND M. CHIARA DOMENEGHETTI¹

¹Department of Earth and Environmental Sciences, University of Pavia, Via A. Ferrata 1, I-27100 Pavia, Italy

²Department of Geosciences, University of Padova, Via Gradenigo 6, 35131 Padova, Italy

³Geoscience Institute, Goethe-University Frankfurt, Altenhöferallee 1, 60323 Frankfurt, Germany

⁴Department of Mineralogy and Petrography, Natural History Museum, Burgring 7, 1010 Vienna, Austria

⁵Department of Earth and Environmental Sciences, University of Milano-Bicocca, I-20126 Milano, Italy

⁶Vereshchagin Institute for High Pressure Physics RAS, Troitsk, Moscow, 108840 Russia

ABSTRACT

The formation and shock history of ureilite meteorites, a relatively abundant type of primitive achondrites, has been debated for decades. For this purpose, the characterization of carbon phases can provide further information on diamond and graphite formation in ureilites, shedding light on the origin and history of this meteorite group. In this work, we present X-ray diffraction and micro-Raman spectroscopy analyses performed on diamond and graphite occurring in the ureilite Yamato 74123 (Y-74123). The results show that nano- and microdiamonds coexist with nanographite aggregates. This, together with the shock-deformation features observed in olivine, such as mosaicism and planar fractures, suggest that diamond grains formed by a shock event (≥ 15 GPa) on the ureilitic parent body (UPB). Our results on Y-74123 are consistent with those obtained on the NWA 7983 ureilite and further support the hypothesis that the simultaneous formation of nano- and microdiamonds with the assistance of a Fe-Ni melt catalysis may be related to the heterogeneous propagation and local scattering of the shock wave. Graphite geothermometry revealed an average recorded temperature (T_{\max}) of 1314 °C (± 120 °C) in agreement with previously estimated crystallization temperatures reported for graphite in Almahata Sitta ureilite.

Keywords: Carbon phases, diamond, graphite, ureilite meteorites, shock, impact event