

Measurements of the Lamb-Mössbauer factor at simultaneous high-pressure-temperature conditions and estimates of the equilibrium isotopic fractionation of iron

**DONGZHOU ZHANG^{1,4,*}, JENNIFER M. JACKSON², WOLFGANG STURHAHN², JIYONG ZHAO³,
E. ERCAN ALP³, AND MICHAEL Y. HU³**

¹Hawaii Institute of Geophysics and Planetology, University of Hawaii at Manoa, Honolulu, Hawaii 96822, U.S.A.

²Seismological Laboratory, California Institute of Technology, Pasadena, California 91125, U.S.A.

³Advanced Photon Source, Argonne National Laboratory, Argonne, Illinois 60439, U.S.A.

⁴GSECARS, University of Chicago, Argonne, Illinois 60439, U.S.A.

ABSTRACT

Isotopic fractionation has been linked to the lattice vibrations of materials through their phonon spectra. The Lamb-Mössbauer factor (f_{LM}) has the potential to provide information about the lattice vibrations in materials. We constrain the temperature evolution of the f_{LM} of γ - and ϵ -Fe at in situ high- P - T conditions between 1650 K and the melting point. We find that the vibrations of γ - and ϵ -Fe can be described using a quasiharmonic model with a pressure- and temperature-dependent Debye temperature computed from the measured f_{LM} . From the Debye temperature, we derive the equilibrium isotopic fractionation β -factor of iron. Our results show that the quasiharmonic behavior of metallic iron would lower the value of $\ln\beta_{\text{Fe}}^{57/54}$ by 0.1‰ at 1600–2800 K and 50 GPa when compared to the extrapolation of room temperature nuclear resonant inelastic X-ray scattering data. Our study suggests that anharmonicity may be more prevalent in Fe metal than in lower mantle minerals at 2800 K and 50 GPa, a relevant condition for the core formation, and the silicate mantle may be isotopically heavy in iron.

Keywords: Iron isotope fractionation, high pressure-temperature, Mossbauer spectroscopy, anharmonicity