

Oxidation state of iron and Fe-Mg partitioning between olivine and basaltic martian melts

ANDREW K. MATZEN^{1,*}, ALAN WOODLAND², JOHN R. BECKETT³, AND BERNARD J. WOOD^{1,†}

¹Department of Earth Sciences, University of Oxford, Oxford, OX1 3AN, U.K.

²Goethe-Universität Institut für Geowissenschaften, Frankfurt, D-60438, Germany

³California Institute of Technology, MC 170-25, Pasadena, California 91125, U.S.A.

ABSTRACT

We performed a series of experiments at 1 atm pressure and temperatures of 1300–1500 °C to determine the effect of oxygen fugacity on the oxidation state of Fe in a synthetic martian basalt. Ferric-ferrous ratios were determined on the quenched glasses using Mössbauer spectroscopy. Following the conventional doublet assignments in the spectrum, we obtain a $\text{Fe}^{3+}/\Sigma\text{Fe}$ value of 0.19 at 1450 °C and an oxygen fugacity corresponding to the QFM buffer. If we apply the Berry et al. (2018) assignments the calculated $\text{Fe}^{3+}/\Sigma\text{Fe}$ drops to 0.09, and the slope of $\log(X_{\text{FeO}_{1.5}}^{\text{melt}}/X_{\text{FeO}}^{\text{melt}})$ vs. $\log(f_{\text{O}_2})$ changes from 0.18 to 0.26.

Combining oxidation state data together with results of one additional olivine-bearing experiment to determine the appropriate value(s) for the olivine (Ol)-liquid (liq) exchange coefficient, $K_{\text{D,Fe}^{2+}\text{-Mg}} = (\text{FeO}/\text{MgO})^{\text{Ol}}/(\text{FeO}/\text{MgO})^{\text{liq}}$ (by weight), suggests a $K_{\text{D,Fe}^{2+}\text{-Mg}}$ of 0.388 ± 0.006 (uncertainty is one median absolute deviation) using the traditional interpretation of Mössbauer spectroscopy and a value of 0.345 ± 0.005 following the Mössbauer spectra approach of Berry et al. (2018).

We used our value of $K_{\text{D,Fe}^{2+}\text{-Mg}}$ to test whether any of the olivine-bearing shergottites represent liquids. For each meteorite, we assumed a liquid composition equal to that of the bulk and then compared that liquid to the most Mg-rich olivine reported. Applying a $K_{\text{D,Fe}^{2+}\text{-Mg}}$ of ~ 0.36 leads to the possibility that bulk Yamato 980459, NWA 5789, NWA 2990, Tissint, and EETA 79001 (lithology A) represent liquids.

Keywords: Olivine-phyric shergottites, Fe^{2+} -Mg partitioning, olivine, basalt, Mars