

## Pliniusite, $\text{Ca}_5(\text{VO}_4)_3\text{F}$ , a new apatite-group mineral and the novel natural ternary solid-solution system pliniusite–svabite–fluorapatite

IGOR V. PEKOV<sup>1,\*</sup>, NATALIA N. KOSHYAKOVA<sup>1</sup>, NATALIA V. ZUBKOVA<sup>1</sup>, ARKADIUSZ KRZĄTAŁA<sup>2,†</sup>,  
DMITRY I. BELAKOVSKIY<sup>3</sup>, IRINA O. GALUSKINA<sup>2</sup>, EVGENY V. GALUSKIN<sup>2</sup>, SERGEY N. BRITVIN<sup>4</sup>,  
EVGENY G. SIDOROV<sup>5,†</sup>, YEVEGENY VAPNIK<sup>6</sup>, AND DMITRY YU PUSHCHAROVSKY<sup>1</sup>

<sup>1</sup>Faculty of Geology, Moscow State University, Vorobievy Gory, 119991 Moscow, Russia

<sup>2</sup>Institute of Earth Sciences, Faculty of Natural Sciences, University of Silesia, Będzińska 60, 41-200 Sosnowiec, Poland

<sup>3</sup>Fersman Mineralogical Museum of the Russian Academy of Sciences, Leninsky Prospekt 18-2, 119071 Moscow, Russia

<sup>4</sup>Department of Crystallography, St. Petersburg State University, University Emb. 7/9, 199034 St. Petersburg, Russia

<sup>5</sup>Institute of Volcanology and Seismology, Far Eastern Branch of Russian Academy of Sciences, Piip Boulevard 9,  
683006 Petropavlovsk-Kamchatsky, Russia

<sup>6</sup>Department of Geological and Environmental Sciences, Ben-Gurion University of the Negev, POB 653, Beer-Sheva 84105, Israel

### ABSTRACT

The new apatite-group mineral pliniusite, ideally  $\text{Ca}_5(\text{VO}_4)_3\text{F}$ , was found in fumarole deposits at the Tolbachik volcano, Kamchatka, Russia, and in a pyrometamorphic rock of the Hatrurim Complex, Israel. Pliniusite, together with fluorapatite and svabite, forms a novel and almost continuous ternary solid-solution system characterized by wide variations of  $T^{5+} = \text{P, As, and V}$ . In paleo-fumarolic deposits at Mountain 1004 (Tolbachik), members of this system, including the holotype pliniusite, are associated with hematite, tenorite, diopside, andradite, kainotrope, baryte and supergene volborthite, brochantite, gypsum and opal. In sublimates of the active Arsenatnaya fumarole (Tolbachik), pliniusite–svabite–fluorapatite minerals coexist with anhydrite, diopside, hematite, berzeliite, schäferite, calciojohillerite, forsterite, enstatite, magnesioferrite, ludwigite, rhabdobarite-group fluoroborates, powellite, baryte, udinaite, arsenudinaite, paraberzeliite, and spinel. At Nahal Morag, Negev Desert, Israel, the pliniusite cotype and V-bearing fluorapatite occur in schorlomite-gehlenite paralava with rankinite, walstromite, zadovite-aradite series minerals, magnesioferrite, hematite, khesinite, barioferrite, perovskite, gurimite, baryte, tenorite, delafossite, wollastonite, and cuspidine. Pliniusite forms hexagonal prismatic crystals up to  $0.3 \times 0.1$  mm and open-work aggregates up to 2 mm across (Mountain 1004) or grains up to 0.02 mm (Nahal Morag and Arsenatnaya fumarole). Pliniusite is transparent to semitransparent, colorless or whitish, with a vitreous luster. The calculated density is  $3.402 \text{ g/cm}^3$ . Pliniusite is optically uniaxial (–),  $\omega = 1.763(5)$ ,  $\varepsilon = 1.738(5)$ . The empirical formulas of pliniusite type specimens calculated based on 13 anions (O+F+Cl) per formula unit are  $(\text{Ca}_{4.87}\text{Na}_{0.06}\text{Sr}_{0.03}\text{Fe}_{0.02})_{\Sigma 4.98}(\text{V}_{1.69}\text{As}_{0.66}\text{P}_{0.45}\text{S}_{0.12}\text{Si}_{0.09})_{\Sigma 3.01}\text{O}_{11.97}\text{F}_{1.03}$  (Mountain 1004) and  $(\text{Ca}_{4.81}\text{Sr}_{0.12}\text{Ba}_{0.08}\text{Na}_{0.05})_{\Sigma 5.06}(\text{V}_{2.64}\text{P}_{0.27}\text{S}_{0.07}\text{Si}_{0.03})_{\Sigma 3.01}\text{O}_{12.15}\text{F}_{0.51}\text{Cl}_{0.34}$  (Nahal Morag). Pliniusite has a hexagonal structure with space group  $P6_3/m$ ,  $a = b = 9.5777(7)$ ,  $c = 6.9659(5)$  Å,  $V = 553.39(7)$  Å<sup>3</sup>, and  $Z = 2$ . The structure was solved using single-crystal (holotype) X-ray diffraction,  $R = 0.0254$ . The mineral was named in honor of the famous Roman naturalist Pliny the Elder, born Gaius Plinius Secundus (AD 23–79). It is suggested that the combination of high temperature, low pressure, and high oxygen fugacity favors the incorporation of  $\text{V}^{5+}$  into calcium apatite-type compounds, leading to the formation of fluorovanadates.

**Keywords:** Pliniusite, apatite group, new mineral, calcium fluoride vanadate, fluorapatite, svabite, crystal structure, X-ray diffraction, Raman spectroscopy, electron microprobe analysis