

## **On the origin of fluorine-poor apatite in chondrite parent bodies**

**FRANCIS M. McCUBBIN<sup>1,\*;‡</sup>, JONATHAN A. LEWIS<sup>1</sup>, JESSICA J. BARNES<sup>2</sup>, JEREMY W. BOYCE<sup>1</sup>,  
JULIANE GROSS<sup>1,3</sup>, MOLLY C. McCANTA<sup>4,‡</sup>, POORNA SRINIVASAN<sup>5,6</sup>, BRENDAN A. ANZURES<sup>7</sup>,  
NICOLE G. LUNNING<sup>1</sup>, STEPHEN M. ELARDO<sup>8</sup>, LINDSAY P. KELLER<sup>1</sup>, TABB C. PRISSEL<sup>9</sup>,  
AND CARL B. AGE<sup>5,6</sup>**

<sup>1</sup>NASA Johnson Space Center, Mailcode XI, 2101 NASA Parkway, Houston, Texas 77058, U.S.A.

<sup>2</sup>Lunar and Planetary Laboratory, University of Arizona, 1629 E University Boulevard, Tucson, Arizona 85721, U.S.A.

<sup>3</sup>Department of Earth and Planetary Sciences, Rutgers University, Piscataway, New Jersey 08854, U.S.A.

<sup>4</sup>Department of Earth and Planetary Sciences, University of Tennessee, Knoxville, Tennessee 37996, U.S.A.

<sup>5</sup>Institute of Meteoritics, University of New Mexico, 200 Yale Boulevard SE, Albuquerque, New Mexico 87131, U.S.A.

<sup>6</sup>Department of Earth and Planetary Sciences, University of New Mexico, 200 Yale Boulevard SE, Albuquerque, New Mexico 87131, U.S.A.

<sup>7</sup>Lunar and Planetary Institute, 3600 Bay Area Boulevard, Houston, Texas 77058, U.S.A.

<sup>8</sup>Department of Geological Sciences, University of Florida, Gainesville, Florida 32611, U.S.A.

<sup>9</sup>Jacobs, NASA Johnson Space Center, 2101 NASA Parkway, Houston, Texas 77058, U.S.A.

### **ABSTRACT**

We conducted a petrologic study of apatite within one LL chondrite, six R chondrites, and six CK chondrites. These data were combined with previously published apatite data from a broader range of chondrite meteorites to determine that chondrites host either chlorapatite or hydroxylapatite with  $\leq 33$  mol% F in the apatite X-site (unless affected by partial melting by impacts, which can cause F-enrichment of residual apatite). These data indicate that either fluorapatite was not a primary condensate from the solar nebula or that it did not survive lower temperature nebular processes and/or parent body processes. Bulk-rock Cl and F data from chondrites were used to determine that the solar system has a Cl/F ratio of  $10.5 \pm 1.0$  ( $3\sigma$ ). The Cl/F ratios of apatite from chondrites are broadly reflective of the solar system Cl/F value, indicating that apatite in chondrites is fluorine poor because the solar system has about an order of magnitude more Cl than F. The Cl/F ratio of the solar system was combined with known apatite-melt partitioning relationships for F and Cl to predict the range of apatite compositions that would form from a melt with a chondritic Cl/F ratio. This range of apatite compositions allowed for the development of a crude model to use apatite X-site compositions from achondrites (and chondrite melt rocks) to determine whether they derive from a volatile-depleted and/or differentiated source, albeit with important caveats that are detailed in the manuscript. This study further highlights the utility of apatite as a mineralogical tool to understand the origin of volatiles (including H<sub>2</sub>O) and the diversity of their associated geological processes throughout the history of our solar system, including at its nascent stage.

**Keywords:** CK chondrite, R chondrite, phosphate, volatiles, chlorine, fluorine, water; Experimental Halogens in Honor of Jim Webster