

# Grain-scale zircon Hf isotope heterogeneity inherited from sediment-metasomatized mantle: Geochemical and Nd-Hf-Pb-O isotopic constraints on Early Cretaceous intrusions in central Lhasa Terrane, Tibetan Plateau

MING-JIAN LI<sup>1,2</sup>, YUN-CHUAN ZENG<sup>1,\*</sup>, MASSIMO TIEPOLO<sup>2</sup>, FEDERICO FARINA<sup>2</sup>, JI-FENG XU<sup>1</sup>,  
FENG HUANG<sup>1</sup>, XI-JUN LIU<sup>3</sup>, QIN CHEN<sup>4</sup>, AND YUAN MA<sup>5</sup>

<sup>1</sup>State Key Laboratory of Geological Processes and Mineral Resources, School of Earth Science and Resources, China University of Geosciences, Beijing 100083, China

<sup>2</sup>Dipartimento di Scienze della Terra “A. Desio”, Università degli Studi di Milano, Via Botticelli 23, Milano 20133, Italy

<sup>3</sup>Guangxi Key Laboratory of Hidden Metallic Ore Deposits Exploration, College of Earth Sciences, Guilin University of Technology, Guilin 541004, China

<sup>4</sup>The Second Geological Exploration Institute of China Metallurgical Geology Bureau, Fuzhou 350108, China

<sup>5</sup>China Railway Design Corporation, Tianjin 300142, China

## ABSTRACT

Clarifying the mechanism of recycling of pre-existing continental crustal materials into the source of mantle-derived magma is a challenging effort that can be of great value to improving our understanding of mantle processes and continental crust growth. This study presents an integrated investigation of whole-rock and mineral geochemical and Nd-Hf-O-Pb isotopic data for dolerites and diorites intruded in the central Lhasa Terrane of Tibetan Plateau at ca. 120 Ma (zircon U-Pb ages). These intrusions have similar distributions of trace elements that are characterized by depletion in Nb-Ta relative to Th, Ba, and U, and moderately negative whole-rock  $\epsilon_{Nd}(t)$  (–5.0 to –1.7) values. Magmatic zircon shows dramatically variable  $\epsilon_{Hf}(t)$  values (from –5.0 to +13.7 in the same rock, including up to 12 epsilon unit variability in single grains). On the other hand, the zircon  $\delta^{18}O$  values are relatively uniform (+6.0‰ to +7.7‰). The constant  $^{208}Pb/^{206}Pb$  values of clinopyroxene crystallized at ca. 500–900 MPa suggest no contamination with lower continental crust. The lack of covariation between Hf and O isotopes from the same grains, and the lack of relationship between Hf isotopes and trace elements (e.g., Hf, Th/U, and Yb/Gd) in the magmatic zircons, together with the absence of ancient zircon xenocrysts, imply limited upper crustal contamination. In combination with high-whole-rock Th/La (>0.29) ratios, we interpret the zircon Hf isotope heterogeneity as inherited from a depleted asthenospheric mantle with the addition of 1–4% Hf from isotopically heterogeneous sediments. Our study therefore emphasizes the need for caution when using complex Hf isotopic zonation in zircon as an argument for intracrustal hybridization of two end-member magmas derived from distinct reservoirs. In addition, the high-Zr/Y ratios and no negative Zr-Hf anomalies of the Aruo intrusions imply a high surface temperature of the down going slab that was able to fully dissolve zircons in the subducted sediments. This requires a special geodynamic condition that was most likely related to the steepening of flatly subducted Neo-Tethyan lithosphere at ca. 120 Ma according to a synthesis of regional tectonic-magmatic-sedimentary records.

**Keywords:** Zircon Hf heterogeneity, sedimentary recycling, mantle metasomatism, magma mixing, Lhasa Terrane