

**Table A1.** Atomic coordinates, equivalent isotropic displacement parameters ( $\text{\AA}^2$ ), and occupancies for the sample La-K-Pym analyzed at 273(2) K.

Atom	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> <sub>eq</sub>	Occ.
<b>Pb1</b>	2/3	1/3	0.49478(4)	0.01495(6)	Pb <sub>0.88(1)</sub> La <sub>0.06(2)</sub> K <sub>0.06(2)</sub>
<b>Pb2</b>	0.75085(2)	0.74497(2)	3/4	0.01481(4)	Pb <sub>1.00</sub>
<b>P</b>	0.40974(13)	0.37862(13)	3/4	0.0086(2)	P <sub>1.00</sub>
<b>O1</b>	0.3419(4)	0.4846(4)	3/4	0.0191(9)	O <sub>1.00</sub>
<b>O2</b>	0.5893(4)	0.4743(4)	3/4	0.0167(9)	O <sub>1.00</sub>
<b>O3</b>	0.3601(3)	0.2728(3)	0.5805(4)	0.0194(6)	O <sub>1.00</sub>
<b>Cl</b>	1	1	1	0.0123(4)	Cl <sub>1.00</sub>

**Table A2.** Experimental details and crystallographic characteristics of La-K-Pym analyzed at two different temperatures, 180(2) K and 273(2) K.

	La-K-Pym analyzed at 180(2) K*	La-K-Pym analyzed at 273(2) K
Diffractometer	Rigaku Oxford Diffraction	Bruker AXS
X-ray radiation	MoK $\alpha$ ( $\lambda = 0.71073 \text{ \AA}$ )	MoK $\alpha$ ( $\lambda = 0.71073 \text{ \AA}$ )
Temperature	180(2) K	273(2) K
Space group	<i>P6<sub>3</sub>/m</i>	<i>P6<sub>3</sub>/m</i>
Unit-cell parameters		
<i>a</i> ( $\text{\AA}$ )	10.00129(8)	10.0011(5)
<i>c</i> ( $\text{\AA}$ )	7.29865(8)	7.2986(5)
<i>V</i> ( $\text{\AA}^3$ )	632.244(12)	632.22(8)
<i>Z</i>	2	2
Absorption coefficient	64.024 mm <sup>-1</sup>	66.43 mm <sup>-1</sup>
F(000)	1113	1132
$\theta$ range	2.35 to 33.78°	2.35 to 28.99°
Index ranges	-15 $\leq$ h $\leq$ 15, -15 $\leq$ k $\leq$ 15, -10 $\leq$ l $\leq$ 11	-13 $\leq$ h $\leq$ 13, -13 $\leq$ k $\leq$ 13, -9 $\leq$ l $\leq$ 9
Collected reflections / unique reflections	45590 / 893	7818 / 584
Refinement method	Full-matrix least-squares on F <sup>2</sup>	Full-matrix least-squares on F <sup>2</sup>
Refined parameters	43	43
R1, Fo > 4sig(Fo)	0.0225	0.0165
R1, all unique data	0.0238	0.0184
wR2	0.0448	0.0351
GooF	1.189	1.175
Extinction coefficient	0.00006(7)	0.00018(4)
Largest difference peaks (e <sup>-</sup> $\text{\AA}^{-3}$ )	4.679 and -1.960	3.065 and -1.008

\* Data repeated in Table 2 and selected for discussion.

**Table A3.** Comparison of selected bond lengths (Å), bond angles (°) and twist angles (°) for La-K-Pym analyzed at two different temperatures, 180(2) K and 273(2) K.

	La-K-Pym analyzed at 180(2) K*	La-K-Pym analyzed at 273(2) K
Pb1–O1 (×3)	2.581(4)	2.582(3)
O2 (×3)	2.674(4)	2.673(3)
O3 (×3)	2.881(5)	2.882(3)
Mean	2.712	2.712
Polyhedral volume (Å <sup>3</sup> )	38.777	38.804
Pb2–O1	3.079(4)	3.074(3)
O2	2.356(6)	2.359(3)
O3 (×2)	2.625(4)	2.624(3)
O3 (×2)	2.636(5)	2.637(3)
Cl (×2)	3.1131(2)	3.11261(18)
Mean	2.773	2.773
Polyhedral volume (Å <sup>3</sup> )	37.019	37.0037
P–O1	1.519(6)	1.518(4)
O2	1.557(6)	1.557(4)
O3 (×2)	1.539(4)	1.540(3)
Mean	1.536	1.539
Polyhedral volume (Å <sup>3</sup> )	1.866	1.865
O1–P–O2	110.5(4)	110.6(2)
O1–P–O3 (×2)	112.3(2)	112.14(14)
O2–P–O3 (×2)	107.3(2)	107.40(14)
O3–P–O3	106.9(4)	106.9(2)
Twist angle	17.91	18.17

\* Data repeated in Table 4 and selected for discussion.