

High-pressure single-crystal X-ray diffraction and infrared spectroscopic studies of the $C2/m$ - $P2_1/m$ phase transition in cummingtonite

HEXIONG YANG, ROBERT M. HAZEN, CHARLES T. PREWITT, LARRY W. FINGER,
REN LU, AND RUSSELL J. HEMLEY

Geophysical Laboratory and Center for High Pressure Research, Carnegie Institution of Washington,
5251 Broad Branch Road, NW, Washington, D.C. 20015-1305, U.S.A.

ABSTRACT

The structural changes associated with the $C2/m$ - $P2_1/m$ phase transition in cummingtonite with $(\text{Fe} + \text{Mn})/(\text{Fe} + \text{Mn} + \text{Mg}) \approx 0.50$ have been studied with single-crystal X-ray diffraction at various pressures up to 7.90 GPa and infrared spectroscopy up to 8.63 GPa. With increasing pressure, the crystal transforms from $C2/m$ to $P2_1/m$ symmetry at ~ 1.21 GPa, as determined by the appearance of reflections violating the $C2/m$ space group. Infrared spectra provide additional evidence for the phase transition: A distinct splitting of OH stretching bands results from an increase from one to two nonequivalent OH positions. The $C2/m$ - $P2_1/m$ transition is of weakly displacive first-order or tricritical character with apparent slope changes in the plots of the axial ratios a/b and a/c as a function of pressure. The unit-cell compression is considerably anisotropic with the a dimension in both $C2/m$ and $P2_1/m$ phases being the most compressible. Major structural changes for the $C2/m$ - $P2_1/m$ transition include: (1) One crystallographically distinct silicate chain becomes two discontinuously, coupled by the splitting of the M4-O5 bond, as well as M4-O6, into two nonequivalent bonds, and (2) the M4-cation coordination increases from sixfold to sevenfold. More importantly, we observed a change in the sense of rotation for the A chain while the crystal structure maintains $P2_1/m$ symmetry: It is O rotated, as the B chain, at 1.32 GPa, but S-rotated at 2.97 GPa and higher pressures. As pressure increases from 1.32 to 7.90 GPa, there is a switching of the nearest bridging O atoms coordinated with the M4 cation: The M4-O5B distance contracts from 2.944 to 2.551 Å, whereas the M4-O6B distance increases from 2.754 to 2.903 Å. Compression mechanisms for the low- and high-pressure polymorphs appear to be slightly different. In the $C2/m$ phase, the behavior of the A and M4 sites controls the compression of the structure, whereas the response of the M1, M2, and M3 octahedra to pressure also plays a role in determining the compression of the $P2_1/m$ structure. The phase transition is regarded as primarily driven by the differential compression between the M4 and T sites, and the symmetry breaking provides a necessary tighter coordination for the M4 site. Based on our data, the obvious changes in the hyperfine parameters of ^{57}Fe in grunerite between 1.0 and 3.4 GPa, observed by Zhang and Hafner (1992), are likely to result from the $C2/m$ - $P2_1/m$ structural transformation.