

## **Experimental phase-equilibrium study of Al- and Ti-contents of calcic amphibole in MORB—A semiquantitative thermobarometer**

**W.G. ERNST\* AND JUN LIU**

Department of Geological and Environmental Sciences, Stanford University, Stanford, California 94305-2115, U.S.A.

### **ABSTRACT**

Calcic amphiboles were synthesized from a natural mid-ocean ridge basalt (MORB) in 39 experiments representing 24 sets of pressure-temperature ( $P$ - $T$ ) conditions ranging from 650–950 °C, 0.8–2.2 GPa, at  $f_{\text{O}_2}$  controlled by the fayalite-magnetite-quartz (FMQ) buffer, and  $P_{\text{aqueous fluid}} = P_{\text{total}}$ . Experiments lasted up to 1630 h at low temperatures; in all cases, synthesized hornblendes were coarse-grained ( $5\text{--}7 \times 10\text{--}15 \mu\text{m}$ ) and chemically homogeneous. Over the investigated pressure range, Ca-amphibole coexisting with phases rich in Al and Ti gradually changes composition from sodic-calcic, Si-rich at low temperatures to calcic, Si-poor at high temperatures: it is barroisite at 650 °C, edenite at 700 °C, and pargasite at 800–950 °C. Electron microprobe data were combined with 41 comparable analyses from the literature for Ca-amphiboles synthesized from MORBs at intermediate  $f_{\text{O}_2}$  in order to erect a petrogenetic grid for the experimental range 0.0–2.2 GPa, 450–1050 °C. Isoleths for  $\text{Al}_2\text{O}_3$  in Ca-amphibole exhibit markedly negative  $P$ - $T$  slopes, indicating increasing  $\text{Al}_2\text{O}_3$  contents with both  $P$  and  $T$ . In contrast,  $\text{TiO}_2$  isopleths are nearly independent of  $P$ , demonstrating that  $\text{TiO}_2$  in Ca-amphibole increases almost exclusively as a function of  $T$ . For natural metabasaltic assemblages that contain coexisting Al-rich and Ti-rich phases, and closely approached chemical equilibrium under crustal or uppermost mantle conditions, this semiquantitative petrogenetic grid allows the simultaneous assignment of attendant  $P$  and  $T$  employing Ca-amphibole  $\text{Al}_2\text{O}_3$  and  $\text{TiO}_2$  contents. However, during slow cooling, natural Ca-amphiboles may exsolve  $\text{TiO}_2$  as rutile, titanite, and/or ilmenite, but in general do not redistribute  $\text{Al}_2\text{O}_3$ , so this thermobarometer must be applied with caution to inhomogeneous specimens.