

Iron in kornerupine: A ^{57}Fe Mössbauer spectroscopic study and comparison with single-crystal structure refinement

**EDWARD S. GREW,^{1,*} GÜNTHER J. REDHAMMER,² GEORG AMTHAUER,² MARK A. COOPER,³
FRANK C. HAWTHORNE,³ AND KARL SCHMETZER⁴**

¹Department of Geological Sciences, University of Maine, 5790 Bryand Center, Orono, Maine 04469, U.S.A.

²Institut für Mineralogie, Universität Salzburg, Hellbrunnerstrasse, 34 A-5020 Salzburg, Austria

³Department of Geological Sciences, University of Manitoba, Winnipeg, Manitoba R3T 2N2, Canada

⁴Marbacherstrasse, 22b, D-85238 Petershausen, Germany

ABSTRACT

Iron is an important constituent of kornerupine, $(\square, \text{Mg}, \text{Fe})(\text{Al}, \text{Mg}, \text{Fe})_9(\text{Si}, \text{Al}, \text{B})_5\text{O}_{21}(\text{OH}, \text{F})$. We obtained Mössbauer spectra at 300 K on twelve samples with $\Sigma\text{Fe} = 0.30\text{--}1.30$ atoms per formula unit (apfu) and $\text{Fe}^{3+}/\Sigma\text{Fe} = 0\text{--}0.31$; several samples were also run at 77 and 430 K. Models allowing unequivocal refinement of the spectra and determination of site occupancies were developed only when single-crystal refinement (SREF) of six of the samples constrained the number of possibilities. The spectra could then be fitted to three Fe^{2+} doublets and one Fe^{3+} doublet. The Fe^{2+} doublets have nearly identical isomer shifts: $\delta = 1.14\text{--}1.19$ mm/s for the octahedral M1 and M2 sites and $1.12\text{--}1.20$ mm/s for the irregular, eightfold-coordinated X site (relative to $\alpha\text{-Fe}$ at 300 K). However, they differ to a variable extent in quadrupole splitting, $\Delta E_Q \approx 1.06\text{--}1.80, 1.83\text{--}2.27, \text{ and } 2.14\text{--}3.41$ mm/s, respectively, to the M1, M2, and X sites. The Fe^{3+} doublet corresponds to the M4 site. The Mössbauer and SREF occupancies are in excellent agreement for the six samples.

The M1 doublet is split in B-bearing kornerupine and the proportion of Fe corresponding to each doublet, as well as quadrupole splitting, varies with B content. Similarly, the X doublet is split in F-bearing kornerupine, and quadrupole splitting of the X site increases with increasing F content. In contrast to most silicates, resolution of the spectra improves with increasing temperature. Quadrupole splitting of the X, M1, and M2 sites decreases with temperature, the X site at a lesser rate consistent with its being the most distorted site.

To a first approximation, the $\text{Fe}^{3+}/\Sigma\text{Fe}$ ratio in kornerupine determined by SREF and Mössbauer spectroscopy increases with increasing Fe_2O_3 and $\text{Fe}^{3+}/\Sigma\text{Fe}$ ratio of the associated sillimanite, sapphirine, and ilmenite-hematite, i.e., the measured $\text{Fe}^{3+}/\Sigma\text{Fe}$ ratios are related to the oxygen fugacity at which the kornerupine crystallized.