

^{57}Fe nuclear forward scattering of synchrotron radiation in hedenbergite $\text{CaFeSi}_2\text{O}_6$ at hydrostatic pressures up to 68 GPa

**L. ZHANG,¹ J. STANEK,² S.S. HAFNER,^{1,*} H. AHSBAHS,¹ H.F. GRÜNSTEUDEL,³ J. METGE,³
AND R. RÜFFER³**

¹Scientific Center of Materials Sciences and Institute of Mineralogy, University of Marburg, 35032 Marburg, Germany

²Institute of Physics, Jagiellonian University, 30-059 Cracow, Poland

³European Synchrotron Radiation Facility, 38043 Grenoble Cedex, France

ABSTRACT

The ^{57}Fe nuclear forward scattering (NFS) of synchrotron radiation and the use of diamond anvils with helium as pressure medium allowed study of the electronic state of Fe^{2+} in the chain silicate hedenbergite $\text{CaFeSi}_2\text{O}_6$ at pressures up to 68 GPa. Characteristics of NFS time spectra were compared with those of conventional Mössbauer spectra.

NFS time spectra of ^{57}Fe in hedenbergite revealed a reversible phase transition between 53 and 68 GPa at room temperature, which is probably a transition from the paramagnetic phase at low pressures to a magnetic phase at high pressures. If this interpretation is correct, the Néel temperature T_N of hedenbergite depends critically on pressure ($T_N = 45$ K at 1 atm).