

## **Order-disorder kinetics in orthopyroxene with exsolution products**

**MICHELE ZEMA,<sup>1,\*</sup> M. CHIARA DOMENEGHETTI,<sup>2,†</sup> AND VITTORIO TAZZOLI<sup>1</sup>**

<sup>1</sup>Dipartimento di Scienze della Terra, Via Ferrata 1, 27100 Pavia, Italy

<sup>2</sup>CNR–Centro di Studio per la Cristallografia e la Cristallografia, Dipartimento di Scienze della Terra, via Ferrata 1, 27100 Pavia, Italy

### **ABSTRACT**

The equilibrium behavior and kinetics of the Fe-Mg intracrystalline exchange reaction in an orthopyroxene sample containing exsolution products were studied by X-ray diffraction (XRD). Isothermal annealing experiments were performed on an orthopyroxene crystal from the Johnstown diogenite, which shows coherent (100) augite lamellae and Guinier-Preston zones. The kinetic experiments were carried out at 700, 800, and 850 °C until Fe-Mg exchange equilibrium was reached. Oxygen fugacity was controlled by the WI buffer. Equilibrium conditions were also confirmed by “reversal” experiments. After each annealing run single-crystal XRD data were collected, and the orthopyroxene phase (*Pbca*) was refined after subtraction of the contribution of the exsolved *C2/c* phase to the observed structure factors. The fraction of augite was ~2%. The low values of the disordering rate constants  $K^+$ , calculated using Mueller’s equation, and the surprisingly high value of the activation energy (102.3 kcal/mol) for the Fe-Mg disordering reaction are ascribed to the presence of exsolution products in the orthopyroxene. Exsolution products seem not to affect the equilibrium behavior.